

MATH 2934 – Additional problem assigned on 2/2/16

Additional problem.

Let $\mathbf{r}(t) = f(t)\mathbf{i} + g(t)\mathbf{j} + h(t)\mathbf{k}$. In your homework you showed that

$$\frac{d}{dt}|\mathbf{r}(t)| = \frac{\mathbf{r}'(t) \cdot \mathbf{r}(t)}{|\mathbf{r}(t)|} \quad (1)$$

(this is Exercise **13.2/53** from the book). In this problem you will derive several other expressions for derivatives related to vector functions.

(a) Show that

$$\frac{d}{dt} \frac{1}{|\mathbf{r}(t)|} = -\frac{1}{|\mathbf{r}(t)|^2} \frac{d}{dt}|\mathbf{r}(t)| = -\frac{\mathbf{r}'(t) \cdot \mathbf{r}(t)}{|\mathbf{r}(t)|^3} .$$

Hint: Use (1) and the identity

$$\frac{d}{dt} \frac{1}{\phi(t)} = -\frac{1}{\phi(t)^2} \frac{d}{dt}\phi(t)$$

for a function ϕ of one variable like in Calculus I (the identity for ϕ follows directly from the Chain Rule for a function of one variable).

(b) Use your result from part (a) to show that

$$\frac{d}{dt} \frac{\mathbf{r}(t)}{|\mathbf{r}(t)|} = -\frac{\mathbf{r}'(t) \cdot \mathbf{r}(t)}{|\mathbf{r}(t)|^3} \mathbf{r}(t) + \frac{\mathbf{r}'(t)}{|\mathbf{r}(t)|} .$$

(c) The vector $\mathbf{u}(t) := \frac{\mathbf{r}(t)}{|\mathbf{r}(t)|}$ is a unit vector in the direction of $\mathbf{r}(t)$. We know from Example 4 in Section 13.2 that if a vector $\mathbf{u}(t)$ has constant length, then the vector is perpendicular to its derivative, i.e., $\mathbf{u}(t) \cdot \mathbf{u}'(t) = 0$. Use your result from part (b) to show by a direct calculation that

$$\frac{\mathbf{r}(t)}{|\mathbf{r}(t)|} \cdot \frac{d}{dt} \frac{\mathbf{r}(t)}{|\mathbf{r}(t)|} = 0 ,$$

as it should be.