

# Symbolic powers in rings of positive characteristic

Daniel Smolkin

Algebra and Algebraic Geometry Seminar  
Madison, Wisconsin  
January 25, 2019

## 1 Learning objectives

1. We can understand the symbolic powers of (positive characteristic) rings by closely studying the maps  $R^{1/p^e} \rightarrow R$ .
2. For certain rings (e.g. Toric varieties, Hibi rings) the study of these maps boils down to (hard!) combinatorics

## 2 Symbolic and ordinary powers of ideals

We assume, for simplicity:

**Global assumptions:**  $R$  is a normal domain finitely generated over a perfect field  $k$ .

Though everything works even if  $k$  is not perfect and  $R$  is just reduced.

3. **Definition:** if  $\mathfrak{p} \in \text{Spec } R$ , we define  $\mathfrak{p}^{(n)} :=$  \_\_\_\_\_
4. Remark: these are larger than ordinary powers, i.e.  $\mathfrak{p}^{(n)} \supseteq \mathfrak{p}^n$ . Rarely an equality.
5. **Exercise:** let  $R = k[x, y, z]/(xy - z^5)$  and  $\mathfrak{p} = (x, z)$ . Then  $\mathfrak{p}^{(5)} =$  \_\_\_\_\_  $\supsetneq \mathfrak{p}^5$ .
6. **Exercise:** Let  $\mathfrak{m} \subseteq R$  be a maximal ideal. Then  $\mathfrak{m}^{(n)} = \mathfrak{m}^n$  for all  $n$ .
7. Intuition:  $\mathfrak{p}^{(n)}$  is the set of regular functions on  $\text{Spec } R$  that \_\_\_\_\_  
\_\_\_\_\_ (cf. Zariski-Nagata theorem).

**Main question:** How does  $\mathfrak{p}^{(n)}$  relate to  $\mathfrak{p}^n$ ? More precisely, for which  $a, b \in \mathbb{N}$  do we have \_\_\_\_\_ ?

8. In 2000, Ein, Lazarsfeld, and Smith gave a striking answer to this question:

**Theorem 1** ([ELS01]). *Let  $R$  be a regular ring over an algebraically closed field of characteristic 0. Then  $\mathfrak{p}^{(hn)} \subseteq \mathfrak{p}^n$  for all prime ideals  $\mathfrak{p}$  of height  $h$ .*

9. We will talk about weakening the regularity assumption in this theorem.
10. Remark: in particular, if  $\dim R = d$ , we see that  $\mathfrak{p}^{(dn)} \subseteq \mathfrak{p}^n$  for all  $\mathfrak{p}$  and all  $n$ . Because this number  $d$  depends only on the ring  $R$  (and not on the primes  $\mathfrak{p}$ ) we say these rings have the *Uniform Symbolic Topology Property*, or USTP for short.

### 3 Commutative algebra mod $p$

11. To prove something like Theorem 1, it actually suffices to work with rings of positive characteristic, using standard “reduction mod  $p$ ” techniques. For instance, to show that

$$R = \frac{\mathbb{C}[x, y, z]}{(x^3 - 5y^2 + 7z^3)}$$

has USTP it suffices to show that its reductions mod  $p$ ,

$$R_p = \frac{\mathbb{C}[x, y, z]}{(x^3 - 5y^2 + 7z^3)}$$

have USTP for all  $p \gg 0$ . In general, there’s a rich theory saying that many properties of a ring in characteristic 0 can be checked mod  $p$  sufficiently large. See [HH99, Chapter 2] for details.

12. **Exercise:** How would one define the reduction of a ring such as

$$S = \frac{\mathbb{C}[x, y, z]}{(\sqrt{2}x^3 - \pi y^2 + \frac{i}{7}z^3)}$$

modulo  $p$ ?

13. **Now let  $R$  have characteristic  $p > 0$ .** Consider the  $R$ -module,  $R^{1/p}$  defined by  $R^{1/p} :=$

**Key idea:** We can learn a lot about  $R$  by studying the  $R$ -module structure of  $R^{1/p^e}$  for  $e > 0$ . Note that  $R^{1/p^e}$  is always a finitely generated module in our setting.

14. For instance, a theorem of Kunz says that  $R$  is regular if and only if  $R^{1/p^e}$  is a flat  $R$ -module for some (all)  $e > 0$  [Kun69].
15. Example from number theory: these modules can be used to detect whether an elliptic curve in positive characteristic is “ordinary” or “supersingular” [BS15].
16. Recall: our goal is to weaken the regularity hypothesis in Theorem 1. The crux of Ein–Lazarsfeld–Smith’s proof<sup>1</sup> is the following chain of containments:

$$\mathfrak{p}^{(hn)} \subseteq \sum_{e>0} \sum_{\varphi: R^{1/p^e} \rightarrow R} \varphi \left( (\mathfrak{p}^{(hn)})^{1/p^e} \right) \subseteq \sum_{e>0} \sum_{\varphi: R^{1/p^e} \rightarrow R} \varphi \left( (\mathfrak{p}^{(hn)})^{\lfloor p^e/n \rfloor / p^e} \right)^n \subseteq \mathfrak{p}^n$$

The second containment breaks if  $R$  is not regular! So we make the sum on the left a little smaller:

**Theorem 2** ([Smo18]). *Let  $R$  be a normal domain finitely generated over a perfect field  $k$  of characteristic  $p$ . Then, for all ideals  $\mathfrak{a}$  of  $R$ , we have<sup>2</sup>*

$$\sum_{e>0} \sum_{\varphi \in \mathcal{D}_e^{(n)}(R)} \varphi \left( \mathfrak{a}^{1/p^e} \right) \subseteq \sum_{e>0} \sum_{\varphi: R^{1/p^e} \rightarrow R} \varphi \left( \mathfrak{a}^{\lfloor p^e/n \rfloor / p^e} \right)^n,$$

<sup>1</sup>At least, the positive-characteristic analog of their proof. The original proof uses *multiplier ideals* which are, fascinatingly, a close analog of these test ideals that works in characteristic 0. Constructing multiplier ideals requires resolution of singularities, which is not known in positive characteristic.

<sup>2</sup>For the experts: I’m sacrificing precision for clarity by omitting test elements in the sums below.

where  $\mathcal{D}_e^{(n)}(R) \subseteq \text{Hom}_R(R^{1/p^e}, R)$  is the set of maps admitting a lifting to the  $n$ -fold tensor product:

$$\begin{array}{ccc} (R^{\otimes_k n})^{1/p^e} & \dashrightarrow & R^{\otimes_k n} \\ \downarrow & & \downarrow \\ R^{1/p^e} & \xrightarrow{\varphi} & R \end{array}$$

17. I won't explain how this works in this talk, but here's the key take-away I want you to have from this discussion:

**Key idea:** This set of maps  $\mathcal{D}_e^{(n)}(R)$  is a correction term that accounts for our ring  $R$  not being regular. If the correction term is not too bad, then the conclusion of Theorem 1 still holds. [CS18, Theorem 4.1]

18. **Definition:** If  $\mathcal{D}_e^{(n)}(R)$  is big enough for the argument to work (for some  $e$ ), then  $R$  is called  $n$ -Diagonally  $F$ -Regular ( $n$ -DFR). If this is true for all  $n > 0$ , we say  $R$  is Diagonally  $F$ -Regular (DFR).

19. **Aside for experts:** Concretely, we need the test ideal of  $\mathcal{D}^{(n)}(R)$  to be all of  $R$ , i.e.

$$\sum_e \sum_{\varphi \in \mathcal{D}_e^{(n)}} \varphi(c^{1/p^e}) = R$$

where  $c \in R$  is some element such that  $R_c$  is regular.

20. So if  $R$  is  $n$ -DFR, then \_\_\_\_\_ for all  $\mathfrak{p}$  of height  $h$ .

**The question becomes:** Which rings are DFR?

21. Facts about Diagonal  $F$ -regularity: regular rings are DFR (exercise! Follows from Kunz's theorem), Segre products of polynomial rings are DFR [CS18] ("non-effective" USTP was known prior to this), tensor products of DFR  $k$ -algebras are DFR [CS18] (new rings with USTP!). DFR rings are strongly  $F$ -regular. DFR rings are not always Gorenstein and can have arbitrarily small  $F$ -signature.
22. **Exercise (hard):** if  $\mathfrak{p}$  is a height 1 prime and torsion element of the divisor class group, then  $\mathfrak{p}^{(n)} \neq \mathfrak{p}^n$  for  $n \gg 0$ . So DFR rings have torsion free divisor class groups [CS18].

## 4 Diagonal $F$ -regularity of Hibi rings

23. A Hibi ring is a kind of (toric) ring associated to a finite partially ordered set.
24. **Definition** Let  $P = \{v_1, \dots, v_n\}$  be a poset. The associated Hibi ring,  $k[P] \subseteq k[x_0, \dots, x_n]$  is defined as follows: we let  $\overline{P} = P \cup \{v_0\}$  where  $v_0 \leq v_i$  for all  $i$ . Then:


$$k[P] := k \left[ x_0^{a_0} \cdots x_n^{a_n} \mid \text{_____} \right]$$

25. If you know about poset ideals, then we can also write

$$k[P] = \frac{k[x_I \mid I \subseteq \overline{P} \text{ a poset ideal}]}{x_I x_J - x_{I \cup J} x_{I \cap J}}$$

26. We usually denote posets by *Hasse diagrams*: nodes represent elements of  $\overline{P}$ . Bigger elements are written above smaller elements. Draw an edge between two distinct nodes  $v_i$  and  $v_j$  if there's no node between them, i.e. if  $v_i \leq v_k \leq v_j$  implies  $v_k = v_i$  or  $v_k = v_j$ . In this case, we say  $v_j$  *covers*  $v_i$ .

27. Some examples/exercises:



28. Checking whether a Hibi ring is  $n$ -DFR boils down to solving a complicated combinatorial problem:

**Theorem 3** ([PST18]). *For each  $i$ , let  $r_i$  be the length of the longest chain going up from  $v_i$  in  $P$ . Then  $k[P]$  is  $n$ -DFR if and only if there exists some  $e$  such that the following holds: for  $0 \leq i \leq d$  and  $1 \leq m \leq n$ , let  $\alpha_{i,m}$  be integers in  $[0, p^e - 1]$  such that  $\sum_{m=1}^n \alpha_{j,m} \equiv r_j \pmod{p^e}$  for all  $j$ . Set  $N_j = \lfloor \sum_{m=1}^n \frac{\alpha_{j,m}}{p^e} \rfloor$ . For all  $i, j$ , and  $m$ , let  $\varepsilon_{j,i,m} = 1$  if  $\alpha_{j,m} > \alpha_{i,m}$  and let  $\varepsilon_{j,i,m} = 0$  otherwise. Then there exist  $\delta_{i,m} \in \mathbb{Z}$  with*


- (a)  $\delta_{i,m} \geq 0$  for all  $m$  whenever  $v_i$  is maximal in  $P$ ,
- (b)  $\delta_{j,m} \leq \varepsilon_{j,i,m} + \delta_{i,m}$  for all  $m$  whenever  $v_j$  covers  $v_i$ , and
- (c)  $\sum_{m=1}^n \delta_{j,m} = N_j$

29. **Aside for experts:** The point is that solving this combinatorial problem is the same as constructing a lifting  $(R^{\otimes n})^{1/p^e} \rightarrow R^{\otimes n}$  of a map  $R^{1/p^e} \rightarrow R$  that sends  $z = x_0^{r_0} \cdots x_n^{r_n}$  to 1. Note that  $z \in R$  and  $R_z$  is regular.

30. Using this combinatorial description, we were able to show:

**Theorem 4** ([PST18]). *If  $k[P]$  is DFR, so is  $k[P \cup \{v'\}]$ , where  $v'$  covers a single element of  $P$ .*

31. Example: Checking if  $\mathbb{F}_5[x, y, z]$  is 3-DFR:



32. Recall: polynomial rings are DFR. Using theorem 4, which posets (Hasse diagrams) do we know to correspond to DFR Hibi rings?

33. Recall: tensor products of DFR rings are DFR. Here's what the tensor product of two Hibi rings looks like:

34. **Exercise:** Convince yourself you get isomorphic rings doing the tensor product in either order!

35. **Exercise:** What are all the Hibi rings known to be DFR, using Theorem 4 and results about DFR rings in item 20?

36. **Definition:** A *top node* in a poset is a node that covers more than one element. They look like hats in the Hasse diagram.

**Theorem 5** ([PST18]). *The Hibi ring  $k[P]$  is DFR whenever the set of top nodes of  $P$  is*

---

37. The converse to this theorem is not known! Here's the first poset with incomparable top nodes:

38. We know it's 2-DFR (in fact, all Hibi rings are 2-DFR). Dylan Johnson has shown it's 3-DFR.

## 5 Questions I would like to know the answer to

39. Is the diagonal  $F$ -regularity of a toric ring independent of characteristic?
40. Is  $\mathcal{D}^{(n)}(R)$  a good metric for the singularities of  $R$ ? For instance, if  $\mathcal{D}_e^{(2)}(R) = \text{Hom}_R(R^{1/p^e}, R)$  for all  $e$ , does that imply  $R$  is regular? This is true for toric  $\mathbb{Q}$ -Gorenstein  $R$ .
41. Do we always have  $\mathcal{D}_e^{(n)}(R) \supseteq \mathcal{D}_e^{(n+1)}(R)$ ? This is true for toric  $R$ .
42. Are rings with large  $F$ -signature (say,  $> 1/2$ ) always DFR? Note that such rings have torsion-free divisor class groups by Carvajal-Rojas.
43. What kind of USTP statements can we get if the  $F$ -signature of  $\mathcal{D}^{(n)}$  is large but  $< 1$ ?

## References

- [BS15] Bhargav Bhatt and Anurag K. Singh. The  $F$ -pure threshold of a Calabi-Yau hypersurface. *Math. Ann.*, 362(1-2):551–567, 2015.
- [CS18] J. Carvajal-Rojas and D. Smolkin. The Uniform Symbolic Topology Property for Diagonally  $F$ -regular Algebras. *ArXiv:1807.03928*, July 2018.
- [ELS01] Lawrence Ein, Robert Lazarsfeld, and Karen E. Smith. Uniform bounds and symbolic powers on smooth varieties. *Invent. Math.*, 144(2):241–252, 2001.
- [HH99] Melvin Hochster and Craig Huneke. Tight closure in equal characteristic zero. 1999. Preprint. URL: [www.math.lsa.umich.edu/~hochster/tcz.ps](http://www.math.lsa.umich.edu/~hochster/tcz.ps).
- [Kun69] Ernst Kunz. Characterizations of regular local rings for characteristic  $p$ . *Amer. J. Math.*, 91:772–784, 1969.
- [PST18] J. Page, D. Smolkin, and K. Tucker. Symbolic and Ordinary Powers of Ideals in Hibi Rings. *ArXiv:1810.00149*, 2018.
- [Smo18] D. Smolkin. A New Subadditivity Formula for Test Ideals. *ArXiv:1805.08739*, 2018.