

Petersson norms of Borcherds theta lifts to $O(1, 8n + 1)$ with applications to sup-norm bounds

Simon Marshall, Hiro-aki Narita and Ameya Pitale

September 3, 2025

Abstract

We give an explicit formula for the Petersson norms of theta lifts from Maass cusp forms of level one to cusp forms on orthogonal groups $O(1, 8n + 1)$. Our formula explicitly determines archimedean local factors of the norms. As an application, we obtain bounds on the sup-norm of the lifted cusp forms on these orthogonal groups in terms of their Laplace eigenvalues.

Contents

1	Introduction	1
2	Classical automorphic forms	6
2.1	Algebraic groups	6
2.2	Automorphic forms and lifting theorem	8
3	Adelization of the Borcherds lift	9
4	Petersson norm of the theta lifting	11
4.1	The archimedean computation	13
5	Sup-norm bounds for Maass cusp forms on $O(1, N + 1)$	20
5.1	Upper and lower bounds using Fourier expansions	20
5.2	Upper bounds using the pre-trace formula	27
5.3	Proof of Theorem 1.2	30

1 Introduction

In this paper, we consider several questions regarding lifts from Maass cusp forms f of level 1 to cusp forms F_f on the orthogonal group $O(1, 8n + 1)$ with respect to the arithmetic subgroup Γ attached to even unimodular lattices L of rank $N = 8n$. The group Γ , denoted by Γ_S in (7), is the subgroup of $O(1, 8n + 1)$ stabilizing $L \oplus \mathbb{Z}^2$ with $\mathbb{Z}^2$ viewed as a lattice of the hyperbolic plane. These lifts were constructed in [37] by giving an explicit formula for the Fourier coefficients of F_f and using the explicit theta lift construction due to Borcherds [4] to prove the automorphy (see Section 2.2). In [37], it was shown that the map $f \rightarrow F_f$ preserves cuspidality and is Hecke equivariant. When restricted to Hecke eigenforms, it was shown that the lifts provide counterexamples to the generalized Ramanujan conjecture (see Theorem 2.2 below for details).

Petersson norm of the lift

One of the main results of this paper is to obtain an explicit formula for the Petersson norm of the lift F_f . For this we use the well-known methods of Rallis [45] of doubling integrals and the Siegel Weil formula. It should be remarked that Kudla-Rallis [34] suggested the regularized Siegel Weil formula, which enables us to calculate the Petersson norm without the limitation of the original Siegel-Weil formula (cf. [57]), and Gan-Qiu-Takeda [20] established the formula in full generality for dual pairs of unitary groups or symplectic-orthogonal groups. We consider Borchers's theta lift construction instead of a general theta lift since we are interested in explicit formulas for the Fourier coefficients of the lifts, an important ingredient in the sup norm application. Hence the theta lifts considered here are only from the group $SL(2)$ to $O(1, N+1)$. We assume that f is a Hecke eigenform. To use the method of [45], we first obtain an adelization of the theta lift construction of Borchers for our case. For this, we follow the work of Kudla [33] where the adelization of the Borchers construction is obtained for orthogonal groups of signature $(p, 2)$. The L^2 -norm of the adelization Φ of F_f can then be rewritten as

$$\|\Phi\|^2 = \int_{\mathrm{SL}_2(\mathbb{Q}) \backslash \mathrm{SL}_2(\mathbb{A})} \left(\int_{\mathrm{SL}_2(\mathbb{Q}) \backslash \mathrm{SL}_2(\mathbb{A})} f_0(g_1) E((g_1^*, g_2), s_0; \Xi_0) dg_1 \right) \overline{f_0(g_2)} dg_2.$$

Here, f_0 is the adelization of f , Ξ_0 is a section in an induced representation of $\mathrm{Sp}_4(\mathbb{A})$ obtained from the Weil representation corresponding to the theta lifting, and E is an Eisenstein series on Sp_4 obtained from the Siegel Weil formula. As in [44], the inner integral is Eulerian and can be written as a product of local integrals (see also [20, Section 11]). Since we are restricted to Maass forms f of level 1, in the non-archimedean case, all the data is unramified and the integral is obtained in [44]. The main contribution here is the archimedean integral computation, which is never an immediate consequence from general formulas cited above. The key ingredient of the archimedean computation is to realize (see Proposition 4.2) that the archimedean section Ξ_∞ is in the trivial K -type of the $\mathrm{Sp}_4(\mathbb{R})$ representation, i.e. it is invariant under the maximal compact subgroup of $\mathrm{Sp}_4(\mathbb{R})$.

In Section 4.1, we compute the local archimedean integral to obtain the following result (see Theorem 4.4).

Theorem. 1.1. *Let L be an even unimodular lattice of dimension N . Let $f \in S(\mathrm{SL}_2(\mathbb{Z}); -\frac{r^2+1}{4})$ be a Hecke Maass eigenform with respect to $\mathrm{SL}_2(\mathbb{Z})$ and let π be the irreducible cuspidal automorphic representation of $\mathrm{GL}_2(\mathbb{A})$ corresponding to f . Let F_f be the lift of f to a cusp form on $O(1, N+1)$ with respect to the arithmetic subgroup Γ attached to L . Then the Petersson norm of F_f is given by the formula*

$$\|F_f\|^2 = \frac{L(\frac{N}{2}, \pi, \mathrm{Ad})}{\zeta(\frac{N}{2}+1)\zeta(N)} \left(2^{1-\frac{N}{2}} \pi^2 \frac{\Gamma(\frac{N}{4} + \frac{\sqrt{-1}r}{2}) \Gamma(\frac{N}{4} - \frac{\sqrt{-1}r}{2})}{\Gamma(\frac{N}{4} + \frac{1}{2})^2} \right) \|f\|^2.$$

Here, $L(s, \pi, \mathrm{Ad})$ is the finite part of the degree 3 adjoint L -function of π .

Let us remark here that, in the case of signature $(p, 2)$, the Petersson inner product of the Borchers lift (or the Kudla-Millson lift) has been computed in Theorem 4.9 of [12]. In that case, the authors start with a holomorphic modular form of full level and the archimedean integral then corresponds to a computation involving the holomorphic discrete series of GL_2 . On the other hand, Theorem 1.1 is the first result we are aware of in the literature which gives an explicit formula for the Petersson norm in the non-holomorphic case.

In [42] and [43], we constructed the Borchers lifting from Maass forms with square free level m to cusp forms on $O(1, 5)$ with respect to arithmetic subgroups corresponding to maximal orders in definite quaternion algebras of discriminant m . The adelization of the lift and the Rallis inner product method can be applied to this case as well. The local archimedean integral can be computed similar to the computation in this paper. The main difference is the computation of the local ramified integral in the case $p|m$ which involves vector valued modular forms. We will be working on this case in the future.

Injectivity of the lifting map

As is the case with most theta lift constructions, proving the automorphy of the lift is straightforward, whereas showing non-vanishing is often highly non-trivial.

For general even unimodular lattices, an immediate corollary to Theorem 1.1 is the injectivity of the map $f \rightarrow F_f$, when restricted to Hecke eigenforms. We can extend this injectivity to non-Hecke eigenforms by using the Linear Algebra trick used in [43] (see Theorem 7.1 of [43]). Note that the latter requires information on Hecke eigenvalues of F_f when f is a Hecke eigenform, and these were computed in [37]. See Corollary 4.5 for details. Note that one could possibly extend the injectivity of the Kudla-Millson lift obtained in [12] to non-Hecke eigenforms if one had further information on the lift in the case of Hecke eigenforms.

Let us remark here that it is well known that all even unimodular lattices of dimension N form a single genus, and their equivalence classes are in one-to-one correspondence with the Γ -cusps (see Section 5.1). If we considered the Fourier expansion of F_f at a Γ -cusp corresponding to an even unimodular lattice whose image of the squared norm map exhausts all positive integers, then it is possible to follow the method in [37] to get injectivity of the map $f \rightarrow F_f$ in general. In fact, in [37], we proved the injectivity by the Fourier expansion at a cusp corresponding to a sum of copies of the E_8 -lattice.

Bounds on sup-norm of the lift F_f

Another main result of this paper is to prove bounds on the sup-norm of F_f . The problem of estimating the sup-norm of Laplace eigenfunctions on a compact Riemannian manifold is a fundamental one in harmonic analysis and mathematical physics. The most basic result on this problem is due to Avacumović [1] and Levitan [36]. If X is a compact Riemannian manifold without boundary and ϕ is an eigenfunction of the Laplacian on X with eigenvalue $-\Lambda \leq 0$, they show that

$$\frac{\|\phi\|_\infty}{\|\phi\|_2} \ll \Lambda^{\frac{\dim X - 1}{4}}. \quad (1)$$

In general, this is the best possible estimate and is indeed achieved when X is a sphere. On the other hand, one expects stronger bounds to hold on a generic manifold. For instance, when X is negatively curved, Bérard [3] showed that the bound (1) can be improved by a factor of $\sqrt{\log \Lambda}$. Moreover, it is often possible to improve (1) by a power of Λ under arithmetic assumptions on X and ϕ . The first such improvement is due to Iwaniec and Sarnak [31]. They considered X a congruence arithmetic hyperbolic surface arising from a quaternion algebra over $\mathbb{Q}$ (possibly split), and ϕ a Hecke–Maass cusp form. In this case, they improved on (1) to obtain

$$\frac{\|\phi\|_\infty}{\|\phi\|_2} \ll_\epsilon \Lambda^{\frac{5}{24} + \epsilon}. \quad (2)$$

They also obtained a lower bound $\sqrt{\log \log \Lambda}$ for an infinite sequence of ϕ , which was later improved to $\exp((1 + o(1))\sqrt{\log \Lambda / \log \log \Lambda})$ by Milićević [41]. Let us also mention several other works on the sup-norm problem [9], [10], [13], [15], [50], [51]. Among other things we also cite [46], which includes an excellent review of the sup-norm problems in terms of elliptic differential operators.

We note that in these arithmetic settings, where X is taken to be a locally symmetric space, one generally assumes that ϕ is an eigenfunction not just of the Laplacian, but of the full ring of invariant differential operators. Under this assumption on ϕ , it was shown by Sarnak [52] that the bound (1) can be strengthened to

$$\frac{\|\phi\|_\infty}{\|\phi\|_2} \ll \Lambda^{\frac{\dim X - \text{Rank } X}{4}}. \quad (3)$$

This is sharp on spaces of compact type, and is the natural analog of (1) for these eigenfunctions.

It is also interesting to investigate the sup-norm of eigenfunctions in the case when X is not compact. In this case, we now assume that ϕ is square-integrable. In the noncompact setting, the bound (1) (and (3), under the appropriate assumptions) continues to hold on fixed compact sets, and it is natural to ask whether it in fact holds globally (i.e. over the entire manifold). For this to happen one certainly needs $\|\phi\|_\infty$ to be finite, and this is not generally true, even when X is a finite volume locally symmetric space. However, it is true if one assumes that X is a finite volume locally symmetric space and ϕ is cuspidal (cf. [23]), and so we make these assumptions from now on.

In some cases, it is known that the bounds (1) and (3) hold globally. See for instance [8] for the case of Hecke–Maass forms on GL_2 over a number field, and [7] for Hecke–Maass forms on the space $\mathrm{PGL}_3(\mathbb{Z}) \backslash \mathrm{PGL}_3(\mathbb{R}) / \mathrm{PO}(3)$. (Note that [7] establishes a bound stronger than (1), but not as strong as (3).) On the other hand, it was shown in [15] that (1) fails on $\mathrm{PGL}_n(\mathbb{Z}) \backslash \mathrm{PGL}_n(\mathbb{R}) / \mathrm{PO}(n)$ for $n \geq 6$. The reason for this failure is the large peaks of the GL_n Whittaker function in higher rank, which lead to large values of cusp forms via the Fourier expansion. Moreover, it is generally expected that the large values produced in this way occur high in the cusp, at height roughly $\sqrt{\Lambda}$; [15] establishes the weaker result that the suprema of these cusp forms occur at points tending to infinity. See [6] for an upper bound that complements the lower bound of [15].

This phenomenon of cusp forms having a peak high in the cusp is a general one, which is already present for GL_2 where it is caused by the transition range of the Bessel function $K_{\sqrt{-1}r}(y)$ at $y \sim r$ [52]. However, in this case the peak produced is only of size $\Lambda^{1/12}$, which is smaller than (1). These results lead one to ask the general question of whether a sequence of cusp forms on X realize their suprema in a fixed compact set, or at a sequence of points tending to infinity.

We remark that if, instead of Maass forms, one considers holomorphic modular forms of growing weight for the group $\mathrm{SL}_2(\mathbb{Z})$, then the growth rate of the sup norm was determined by Xia [55] using the Fourier expansion. Similar results in the case of Siegel modular forms were obtained by Blomer [5].

In this paper, we investigate these questions for the forms F_f – namely, whether the bound (1) holds globally, and whether the F_f exhibit large peaks in the cusp. Our main result is the following.

Theorem. 1.2. *Let $f \in S(\mathrm{SL}_2(\mathbb{Z}); -\frac{r^2+1}{4})$ be a non-zero Hecke Maass cusp form with Fourier coefficients $\{c(m) : m \in \mathbb{Z}\}$ satisfying $c(m) = \pm c(-m)$ for all $m \in \mathbb{Z}$. Suppose L is an even unimodular lattice of dimension N . Let F_f be the lift of f to a cusp form on $\mathrm{O}(1, N+1)$ with respect to the arithmetic subgroup Γ attached to L . Let $-\Lambda$ be the Laplace eigenvalue of F_f . Then for any $\epsilon > 0$ and any r , we have*

$$\frac{\|F_f\|_\infty}{\|F_f\|_2} \ll_{N, \epsilon, \Gamma} \Lambda^{\frac{N}{4} + \frac{N(1+2\theta)}{8(N+1+2\theta)} + \epsilon} \leq \Lambda^{\frac{N}{4} + \frac{\theta}{4} + \frac{1}{8} + \epsilon},$$

where $\theta = 7/64$ is the current best estimate towards the Ramanujan conjecture for Maass forms. We also have the lower bound

$$\Lambda^{\frac{N}{8} + \frac{1}{12} - \epsilon} \ll_{N, \epsilon, \Gamma} \frac{\|F_f\|_\infty}{\|F_f\|_2}. \quad (4)$$

See Theorems 5.4 and 5.5 for more details and more general results. We note that the bound (1) in this case reads $\|F_f\|_\infty / \|F_f\|_2 \ll \Lambda^{N/4}$, so that we are not quite able to obtain this globally.

Remark 1. We are free to assume that $r \gg 1$ when proving Theorem 1.2. On the other hand, this condition holds automatically, as it is known that any cuspidal Laplace eigenfunction on the modular surface has eigenvalue $-\Lambda < -1/4$, see for instance [11, 48].

Remark 2. As alluded to above, the lower bound we prove for F_f is obtained high in the cusp. In contrast to this, there are several papers that establish power growth of eigenfunctions on hyperbolic manifolds, and more generally on locally symmetric spaces of noncompact type, in *fixed* compact sets [13, 14, 17, 35, 40, 49]. We mention in particular [14, 17], which apply to the higher-dimensional hyperbolic setting considered here. (The results of [13] also include this, but the growth exponents produced are ineffective.) In [17], Donnelly constructs compact hyperbolic $(N+1)$ -manifolds for $N \geq 4$, and sequences of Laplace

eigenfunctions $\{\phi_i\}_i$ that satisfy $\|\phi_i\|_\infty/\|\phi_i\|_2 \gg \Lambda_i^{(N-3)/4}$. In [14], this lower bound was improved to $\Lambda_i^{(N-1)/4-\epsilon}$ for N even, and it is expected that this parity condition can be removed. Moreover, similarly to F_f , the forms constructed in these papers are theta lifts from SL_2 . These results lead one to hope that F_f might satisfy the same lower bound, which should be realized in a fixed compact subset of the manifold. As this is larger than the lower bound of $\Lambda^{\frac{N}{8}+\frac{1}{12}-\epsilon}$ obtained from the peak of the Whittaker function, one might therefore expect F_f to realize their sup norms in the bulk, rather than the cusp.

We now give an outline of the proof of Theorem 1.2. The proof works by combining two ingredients. The first is a standard upper bound coming from the pre-trace formula, which is valid for any square-integrable Laplace eigenfunction on any finite-volume hyperbolic orbifold. This is Theorem 5.5, which gives

$$\frac{|F_f(n(x)a_y)|}{\|F_f\|_2} \ll \Lambda^{\frac{N}{4}} + y^{\frac{N}{2}} \Lambda^{\frac{N}{8}}. \quad (5)$$

Here, $n(x)a_y$ are Iwasawa coordinates (cf. (8)) adapted to one of the cusps.

The second ingredient is upper and lower bounds proved using the Fourier expansion of F_f . These bounds rely on the fact that these forms are theta lifts in two ways; first, by providing an explicit formula for the Fourier coefficients of F_f in terms of those of f , and secondly through the formula for $\|F_f\|_2$ from Theorem 1.1. The lower bound comes from the first nonzero Fourier coefficients of F_f , combined with the transition behavior of the Bessel function; see the end of Section 5.1.

The upper bound is stated in Theorem 5.4, and gives

$$\frac{1}{\|F_f\|_2} |F_f(n(x)a_y)| \ll_{\epsilon, N, L} \begin{cases} y^{-N/2-1-2\theta} r^{3N/4+1+2\theta+\epsilon} & 1 \ll y \leq r^{11/12}; \\ y^{-N/2+1-2\theta} r^{3N/4-5/6+2\theta+\epsilon} & r^{11/12} < y \leq r/2\pi; \\ e^{-Cy} & r/2\pi < y. \end{cases} \quad (6)$$

Here, $\theta = 7/64$ is the current best estimate towards the Ramanujan conjecture for Maass forms, and we note that $\Lambda \sim r^2$. We combine (5) and (6) by noting that the first bound is strong low in the cusp, while the second is strong high in the cusp. In fact, the first bound gives $\frac{\|F_f\|_\infty}{\|F_f\|_2} \ll \Lambda^{N/4}$ when $y < \Lambda^{1/4}$, while for $y > \Lambda^{1/4}$ the second gives $\frac{\|F_f\|_\infty}{\|F_f\|_2} \ll \Lambda^{N/4+\frac{1}{4}+\frac{\theta}{2}}$. Finding the point at which they are equally strong gives the upper bound of Theorem 1.2.

Bounds on Fourier coefficients of the lift F_f

As discussed above, Theorems 1.2 and 5.4 on sup-norm bounds rely on an estimate for the Fourier coefficients of F_f . By using standard techniques, one can obtain the Hecke bound for the Fourier coefficients $\{A(\lambda) : \lambda \in L \setminus \{0\}\}$ of F_f given by $|A(\lambda)| \ll |\lambda|^{\frac{N}{2}}$. Here, the implied constant depends on the sup-norm of F_f , the quantity we wish to bound. Hence, the Hecke bound is not adequate for obtaining the sup-norm bounds on F_f . Using the explicit formula for $A(\lambda)$ in terms of the Fourier coefficients of the Maass form f , we are able to obtain in Proposition 5.1 the improved bounds of the form $|A(\lambda)| \ll |\lambda|^{\frac{N}{2}-1+\epsilon}$. In the special case of primitive λ , i.e., satisfying $\frac{1}{n}\lambda \notin L$ for all integers $n > 1$, we get an even better bound $|A(\lambda)| \ll |\lambda|^{2\theta+1+\epsilon}$, where $\theta = 7/64$ is the current best estimate towards the Ramanujan conjecture for Maass forms. Here the implied constants depend on ϵ, r and $\|f\|_2$.

These bounds on the size of the Fourier coefficients of the lift are analogous to the results obtained by Ikeda and Katsurada [28] in the context of Siegel cusp forms. The Hecke bound for a Siegel cusp form of genus n and weight k is given by $|A(T)| \ll (\det(2T))^{\frac{k}{2}}$, where $\{A(T)\}$ are the Fourier coefficients with T running through $n \times n$ half integral, positive definite, symmetric matrices. In [28], for the subset of Ikeda lifts, the authors prove better bounds given by $|A(T)| \ll (\det(2T))^{\frac{k}{2}-\frac{1}{2}}$. Furthermore, they show that if T is primitive, then we have $|A(T)| \ll (\det(2T))^{\frac{k}{2}-\frac{1}{2}-\frac{n}{4}}$.

Outline of the paper

The paper is outlined as follows. In Section 2, after reviewing basic notions on orthogonal groups, their algebraic subgroups and real hyperbolic spaces, we introduce automorphic forms that are the focus of this paper. We recall the construction of the theta lifting map as worked out in [37] and state the main theorem from that work regarding these lifts. In Section 3, we carry out the adelization of the theta lifts. In Section 4, we set up the global integrals for the Petersson norm of the lifting using the method of Rallis. The bulk of the computation is done in Section 4.1, where the archimedean integral is calculated. We end Section 4.1 with the explicit formula for the Petersson norm of the lift and the injectivity of the lifting map. In Section 5.1, we compute upper and lower bounds on the sup-norm of the lift using its Fourier expansion. In Section 5.2, we apply the pre-trace formula method to obtain upper bounds for any L^2 eigenfunction of the Laplacian on an $(N + 1)$ -dimensional hyperbolic orbifold of finite volume. We put the results from these methods together to obtain the proof of Theorem 1.2 in Section 5.3.

Notation

If $A \in \mathbb{C}$ and $B \in \mathbb{R}$, we use the notation $A \ll B$ to mean that there is a constant $C > 0$ such that $|A| \leq CB$, and $A \sim B$ means that $A \ll B \ll A$.

Acknowledgement

The authors would like to thank Jens Funke for discussions regarding the local computations of the Petersson norm. We would like to thank Abhishek Saha for giving expert advice on the analytic number theoretic aspect of the paper. We would also like to thank Shingo Sugiyama for his fruitful comments related to this paper, which has led us to cite [6] and [48], and Peter Humphries for sharing the reference [11]. The first author would like to thank Victoria University of Wellington for their hospitality while this paper was being completed. The first author was supported by NSF grant DMS-1902173. The second author was supported by Waseda University Grant for Special Research Projects (Project number: 2020C-611). This work was supported by the Research Institute for Mathematical Sciences, an International Joint Usage/Research Center located in Kyoto University.

2 Classical automorphic forms

2.1 Algebraic groups

For $N \in \mathbb{N}$, let $S \in M_N(\mathbb{Q})$ be a positive definite symmetric matrix and put $Q := \begin{bmatrix} & & 1 \\ & -S & \\ 1 & & \end{bmatrix}$. We then define a $\mathbb{Q}$ -algebraic group $\mathcal{G}$ by the group

$$\mathcal{G}(\mathbb{Q}) := \{g \in M_{N+2}(\mathbb{Q}) \mid {}^t g Q g = Q\}$$

of $\mathbb{Q}$ -rational points. We introduce another $\mathbb{Q}$ -algebraic group $\mathcal{H}$ by the group

$$\mathcal{H}(\mathbb{Q}) := \{h \in M_N(\mathbb{Q}) \mid {}^t h S h = S\}$$

of $\mathbb{Q}$ -rational points. Let q_S , resp. q_Q , denote the quadratic form on $\mathbb{Q}^N$, resp. $\mathbb{Q}^{N+2}$, associated to S , resp. Q , i.e.

$$q_S(v) = \frac{1}{2} {}^t v S v, \quad q_Q(w) = \frac{1}{2} {}^t w Q w$$

for $v \in \mathbb{Q}^N$ and $w \in \mathbb{Q}^{N+2}$. Then $\mathcal{H}$, resp. $\mathcal{G}$, is the orthogonal group associated to this quadratic form q_S , resp. q_Q . For every place $v \leq \infty$ of $\mathbb{Q}$ we put $G_v := \mathcal{G}(\mathbb{Q}_v)$ and $H_v := \mathcal{H}(\mathbb{Q}_v)$.

In addition, we introduce the standard proper $\mathbb{Q}$ -parabolic subgroup $\mathcal{P}$ of $\mathcal{G}$ with the Levi decomposition $\mathcal{P} = \mathcal{N}\mathcal{L}$, where the $\mathbb{Q}$ -subgroups $\mathcal{N}$ and $\mathcal{L}$ are defined by

$$\begin{aligned}\mathcal{N}(\mathbb{Q}) &:= \left\{ n(x) = \begin{bmatrix} 1 & {}^t x S & \frac{1}{2} {}^t x S x \\ & 1_N & x \\ & & 1 \end{bmatrix} \mid x \in \mathbb{Q}^N \right\}, \\ \mathcal{L}(\mathbb{Q}) &:= \left\{ \begin{bmatrix} \alpha & & \\ & \delta & \\ & & \alpha^{-1} \end{bmatrix} \mid \alpha \in \mathbb{Q}^\times, \delta \in \mathcal{H}(\mathbb{Q}) \right\}.\end{aligned}$$

Let us regard $J = \mathbb{Z}^2$ as a lattice of the real hyperbolic plane, and let L be a maximal lattice with respect to S . We then put

$$L_0 := \left\{ \begin{bmatrix} x \\ y \\ z \end{bmatrix} \in \mathbb{Q}^{N+2} \mid x, z \in \mathbb{Z}, y \in L \right\} = L \oplus J,$$

which is a maximal lattice with respect to Q . Here, see [54, Chapter II, Section 6.1] for the definition of maximal lattices. Through the bilinear form induced by the quadratic form q_S , the dual lattice $L^\sharp := \text{Hom}_{\mathbb{Z}}(L, \mathbb{Z})$ is identified with a sublattice of $\mathbb{Q}^N$ containing L .

Assumption: We will assume that L is an unimodular even lattice. This implies that $8|N$ and $L^\sharp = L$.

For each finite prime $p < \infty$ we introduce $L_{0,p} := L_0 \otimes_{\mathbb{Z}} \mathbb{Z}_p$ and put

$$K_p := \{g \in G_p \mid gL_{0,p} = L_{0,p}\},$$

which forms a maximal open compact subgroup of G_p . On the other hand, let $R := \begin{bmatrix} 1 & & \\ & S & \\ & & 1 \end{bmatrix}$ and put

$$K_\infty := \{g \in G_\infty \mid {}^t g R g = R\},$$

which is a maximal compact subgroup of G_∞ . Let $K_f := \prod_{p < \infty} K_p$ and $K := K_f \times K_\infty$. The groups K_f and K form maximal compact subgroups of $\mathcal{G}(\mathbb{A}_f)$ and $\mathcal{G}(\mathbb{A})$ respectively. We furthermore put $U := U_f \times H_\infty$ with $U_f := \prod_{p < \infty} U_p$, where

$$U_p := \{h \in H_p \mid hL_p = L_p\}$$

with $L_p := L \otimes_{\mathbb{Z}} \mathbb{Z}_p$. We now set

$$\Gamma_S := \mathcal{G}(\mathbb{Q}) \cap K_f G_\infty = \{\gamma \in \mathcal{G}(\mathbb{Q}) \mid \gamma L_0 = L_0\}. \quad (7)$$

We have the following result [37, Lemma 2.1],

Lemma. 2.1. *i) (Strong approximation theorem for $\mathcal{G}$) The class number of $\mathcal{G} = \text{O}(Q)$ with respect to $G_\infty K_f$ is one. Namely $\mathcal{G}(\mathbb{A}) = \mathcal{G}(\mathbb{Q}) G_\infty K_f$.*

ii) The class number of $\mathcal{H} = \text{O}(S)$ with respect to U coincides with the number of Γ_S -cusps.

The real Lie group G_∞ admits an Iwasawa decomposition

$$G_\infty = N_\infty A_\infty K_\infty,$$

where

$$N_\infty := \{n(x) \mid x \in \mathbb{R}^N\}, \quad A_\infty := \left\{ a_y = \begin{bmatrix} y & & \\ & 1_N & \\ & & y^{-1} \end{bmatrix} \mid y \in \mathbb{R}_+^\times \right\}. \quad (8)$$

From the Iwasawa decomposition we can identify the homogeneous space G_∞/K_∞ with the $(N+1)$ -dimensional real hyperbolic space $H_N := \{(x, y) \mid x \in \mathbb{R}^N, y \in \mathbb{R}_{>0}\}$ by the natural map

$$n(x)a_y \mapsto (x, y).$$

The cusp forms we are going to study are regarded as cusp forms on the real hyperbolic space H_N .

2.2 Automorphic forms and lifting theorem

For $\lambda \in \mathbb{C}$ and a congruence subgroup $\Gamma \subset \mathrm{SL}_2(\mathbb{R})$ we denote by $S(\Gamma, \lambda)$ the space of Maass cusp forms of weight 0 on the complex upper half plane $\mathfrak{h} := \{u + \sqrt{-1}v \in \mathbb{C} \mid v > 0\}$ whose eigenvalue with respect to the hyperbolic Laplacian is λ .

For $r \in \mathbb{C}$ we denote by $\mathcal{M}(\Gamma_S, r)$ the space of smooth functions F on G_∞ satisfying the following conditions:

- i) $\Omega \cdot F = \frac{1}{2N} \left(r^2 - \frac{N^2}{4} \right) F$, where Ω is the Casimir operator defined in [37, (2.3)],
- ii) for any $(\gamma, g, k) \in \Gamma_S \times G_\infty \times K_\infty$, we have $F(\gamma g k) = F(g)$,
- iii) F is of moderate growth.

Let $r \in \mathbb{R}$. We say that $F \in \mathcal{M}(\Gamma_S, \sqrt{-1}r)$ is a cusp form if it vanishes at the cusps of Γ_S . By [37, Section 2.3], any cusp form $F \in \mathcal{M}(\Gamma_S, \sqrt{-1}r)$ has a Fourier expansion of the form

$$F(n(x)a_y) = \sum_{\lambda \in L \setminus \{0\}} A(\lambda) y^{\frac{N}{2}} K_{\sqrt{-1}r}(4\pi|\lambda|_S y) \exp(2\pi\sqrt{-1}^t \lambda S x). \quad (9)$$

Here, $|\lambda|_S := \sqrt{q_S(\lambda)}$ and $K_{\sqrt{-1}r}$ is the K -Bessel function.

Let

$$f(\tau) = \sum_{n \neq 0} c(n) W_{0, \frac{\sqrt{-1}r}{2}}(4\pi|n|v) \exp(2\pi\sqrt{-1}nu) \in S(\mathrm{SL}_2(\mathbb{Z}); -\frac{r^2+1}{4}) \quad (10)$$

be a Maass cusp form on $\mathfrak{h}$, where we use the Whittaker function $W_{0, \frac{\sqrt{-1}r}{2}}$ to describe the Fourier expansion of f . Recall that we have supposed that L is an even unimodular lattice of rank N divisible by 8. For $0 \neq \lambda \in L$, define

$$A(\lambda) := |\lambda|_S \sum_{d \mid d_\lambda} c\left(-\frac{|\lambda|_S^2}{d^2}\right) d^{\frac{N}{2}-2}, \quad (11)$$

where d_λ denotes the greatest integer such that $\frac{1}{d_\lambda} \lambda \in L$. By Theorems 3.1, 3.3, 4.11 and 5.6 from [37] we have

Theorem. 2.2. *Let L be an even unimodular lattice. Let $f \in S(\mathrm{SL}_2(\mathbb{Z}); -\frac{r^2+1}{4})$ with Fourier expansion (10). Let $F_f : H_N \rightarrow \mathbb{C}$ be given by the Fourier expansion (9) with Fourier coefficients $A(\lambda)$ defined in (11). Then*

- i) *The map $f \rightarrow F_f$ is a map from $S(\mathrm{SL}_2(\mathbb{Z}); -\frac{r^2+1}{4})$ to $\mathcal{M}(\Gamma_S, \sqrt{-1}r)$ preserving cuspidality.*
- ii) *If f is a Hecke eigenform, then so is F_f .*

iii) Suppose f is a Hecke eigenform. Let π_{F_f} be the cuspidal automorphic representation of $\mathcal{G}(\mathbb{A})$ generated by F_f . Then π_{F_f} is irreducible, and thus has the decomposition into the restricted tensor product $\otimes'_{v \leq \infty} \pi_{F_f, v}$ of irreducible admissible representations $\pi_{F_f, v}$ of G_v . For $v = p < \infty$, the representation $\pi_{F_f, p}$ is the spherical constituent of an unramified principal series representation of G_p .

iv) The finite part of the degree $N + 2$ standard L -function of π_{F_f} is given by

$$L(s, \pi_{F_f}) = L(s, \pi_f, \text{Sym}^2) \prod_{i=-(\frac{N}{2}-1)}^{\frac{N}{2}-1} \zeta(s-i),$$

where π_f is the cuspidal, automorphic representation of $\text{GL}_2(\mathbb{A})$ generated by f and $L(s, \pi_f, \text{Sym}^2)$ is the degree 3 symmetric square L -function of π_f .

v) For every finite prime $p < \infty$, $\pi_{F_f, p}$ is non-tempered while $\pi_{F_f, \infty}$ is tempered.

Let us remark here that, in [37], the injectivity of the map $f \rightarrow F_f$ was proven for general even unimodular lattices by using the Fourier expansion at a cusp corresponding to a sum of copies of the E_8 -lattice, which has the property that it contains vectors of squared length M for all $M \in \mathbb{Z}_{>0}$. Corollary 4.5 below provides another proof for the injectivity of the map $f \rightarrow F_f$, which also works in full generality.

3 Adelization of the Borchers lift

In [37], the automorphy of the lift F_f is proved by showing that it is a Borchers theta lift. In [37, Section 3] the classical construction of the theta functions and Borchers lifts has been explained in details. The first main result of this paper is the Petersson norm for F_f . We wish to use the Siegel-Weil formula and the Rallis inner product formula for this. For this we first need to adelize the Borchers lift. The main reference for this is [33], where the adelic Borchers lift has been worked out for signature $(p, 2)$. We will do this for signature $(1, N+1)$. Recall that we have assumed that L is an even unimodular lattice. From now on, let V_N be the quadratic space of dimension $N+2$ defined over $\mathbb{Q}$ equipped with the quadratic form q_Q defined in Section 2.1. Let B_Q be the bilinear form corresponding to the quadratic form q_Q . By $V_N(\mathbb{R})$ and $V_N(\mathbb{A})$ we denote the real quadratic space and the adelic quadratic space attached to V_N respectively. The former is often denoted simply by $\mathbb{R}^{1, N+1}$. By $V_N(\mathbb{A}_f)$ we denote the space formed by the finite adeles in $V_N(\mathbb{A})$.

Let $\mathcal{D}$ be the Grassmanian of positive oriented lines in the real quadratic space $V_N(\mathbb{R})$. For $(x, y) \in H_N$, set $\nu(x, y) := \frac{1}{\sqrt{2}} {}^t(y + y^{-1}q_S(x), -y^{-1}x, y^{-1}) \in V_N(\mathbb{R})$ satisfying $B_Q(\nu(x, y), \nu(x, y)) = 1$. Then, we can identify H_N with one of the two connected components $\mathcal{D}^+$ of $\mathcal{D}$ via

$$H_N \ni (x, y) \rightarrow \mathbb{R} \cdot \nu(x, y) \in \mathcal{D}^+. \quad (12)$$

Denote by $X = \mathcal{G}(\mathbb{Q}) \backslash (\mathcal{D} \times \mathcal{G}(\mathbb{A}_f)) / K_f$. By Lemma 2.1, we have $\mathcal{G}(\mathbb{A}) = \mathcal{G}(\mathbb{Q})\mathcal{G}(\mathbb{R})K_f$. We have $X \simeq \Gamma_S \backslash \mathcal{D}^+$, where $\Gamma_S = \mathcal{G}(\mathbb{Q}) \cap \mathcal{G}(\mathbb{R})K_f$.

Let $S(V_N(\mathbb{A}))$, $S(V_N(\mathbb{A}_f))$ and $S(V_N(\mathbb{R}))$ be the space of Schwartz Bruhat functions of $V_N(\mathbb{A})$, $V_N(\mathbb{A}_f)$ and $V_N(\mathbb{R})$ respectively. For $\nu \in \mathcal{D}$, we have the map

$$\begin{aligned} \iota_\nu : V_N(\mathbb{R}) &\rightarrow \mathbb{R} \cdot \nu \oplus (\nu^\perp, q_S|_{\nu^\perp}) \simeq \mathbb{R}^{1, N+1} \\ \lambda &\mapsto (\iota_\nu^+(\lambda), \iota_\nu^-(\lambda)). \end{aligned}$$

Here, ι_ν^+ and ι_ν^- are the projections from $V_N(\mathbb{R})$ to $\mathbb{R}\nu$ and $\nu^\perp, q_S|_{\nu^\perp}$ respectively. For $\lambda \in V_N(\mathbb{R})$, set $R(\lambda, \nu) := -2q_Q(\iota_\nu^+(\lambda))$ and $(\lambda, \lambda)_\nu := 2q_Q(\lambda) + 2R(\lambda, \nu)$. We can see that

$$(\lambda, \lambda)_\nu = 2q_Q(\lambda) + 2R(\lambda, \nu) = 2q_Q(\iota_\nu^+(\lambda)) + 2q_Q(\iota_\nu^-(\lambda)) - 4q_Q(\iota_\nu^+(\lambda))$$

$$= 2q_Q(\iota_\nu^-(\lambda)) - 2q_Q(\iota_\nu^+(\lambda)) = 2q_Q(\lambda_{\nu-}) - 2q_Q(\lambda_{\nu+}).$$

Here we are denoting $\iota_\nu^+(\lambda) = \lambda_{\nu+}$ and $\iota_\nu^-(\lambda) = \lambda_{\nu-}$. Let $A^\circ(\mathcal{D})$ be the space of smooth functions on $\mathcal{D}$. We then introduce the Gaussian

$$\tilde{\phi}_\infty(\lambda, \nu) := \exp(\pi(\lambda, \lambda)_\nu) = \exp(2\pi q_Q(\lambda_{\nu-}) - 2\pi q_Q(\lambda_{\nu+})). \quad (13)$$

We can regard this as a $A^\circ(\mathcal{D})$ -valued Schwartz function on $V_N(\mathbb{R})$ by

$$V_N(\mathbb{R}) \ni \lambda \rightarrow (\mathcal{D} \ni \nu \mapsto \tilde{\phi}_\infty(\lambda, \nu)).$$

Note that, in [33], there is a minus sign in the exponent instead of plus sign. The reason for the difference is that we are considering signature $(1, N+1)$ instead of $(p, 2)$. Consider the polynomial $P(\lambda, \nu)$ obtained by applying the operator $\exp(-\Delta/(8\pi))$ to the polynomial $2^{-\frac{N}{4}-3}x_1^{\frac{N}{2}}$ defined in [37, Section 3.2], where Δ denotes the Laplacian for the $(N+2)$ -dimensional Euclidean space with the coordinate $(x_1, \dots, x_{N+2})$. For $\lambda \in V_N(\mathbb{R})$ and $\nu \in \mathcal{D}$, define

$$\phi_\infty(\lambda, \nu) := P(\lambda, \nu)\tilde{\phi}_\infty(\lambda, \nu). \quad (14)$$

Lemma. 3.1. *For all $h \in G_\infty$, we have*

$$\phi_\infty(h\lambda, h\nu) = \phi_\infty(\lambda, \nu).$$

Proof. The lemma follows from the observation that both P and $\tilde{\phi}_\infty$ depend only on $q_Q(\lambda)$ and $B_Q(\lambda, \nu)$, and by definition G_∞ preserves this. $\square$

The group $\mathrm{SL}_2(\mathbb{A})$ acts on $S(V_N(\mathbb{A}))$ via the Weil representation ω determined by the standard additive character ψ on $\mathbb{A}/\mathbb{Q}$ such that $\psi_\infty(x) = \exp(2\pi\sqrt{-1}x)$. If $\Phi \in S(V_N(\mathbb{A}))$, then the Weil representation is given by

$$\omega\left(\begin{bmatrix} a & \\ & a^{-1} \end{bmatrix}\right)\Phi(x) = |a|^{\frac{N}{2}+1}\Phi(ax), \quad \omega\left(\begin{bmatrix} 1 & t \\ & 1 \end{bmatrix}\right)\Phi(x) = \psi(tq_Q(x))\Phi(x), \quad \omega\left(\begin{bmatrix} & 1 \\ -1 & \end{bmatrix}\right)\Phi(x) = \hat{\Phi}(x), \quad (15)$$

where $\hat{\Phi}$ denotes the Fourier transform of Φ with respect to a self-dual measure of $V_N(\mathbb{A})$ with respect to $V_N(\mathbb{A})^2 \ni (x, y) \mapsto \psi(B_Q(x, y)) \in \mathbb{C}^{(1)}$. In the first equation, we need to add a factor $\chi_{q_Q}(a) := \langle (-1)^{\frac{N}{2}+1} \det(Q), a \rangle$ on the right hand side. But, note that $\det(Q) = -1$ and $\frac{N}{2} + 1$ is odd. Hence, $\chi_{q_Q}(a) = 1$ for all a .

By ω_v we denote the v -component of ω at a place v . Let $\hat{\Phi}$ denote the Fourier transform of $\Phi \in S(V_N(\mathbb{Q}_v))$ with respect to a self-dual measure of $V_N(\mathbb{Q}_v)$ with respect to $V_N(\mathbb{Q}_v)^2 \ni (x, y) \mapsto \psi_v(B_Q(x, y)) \in \mathbb{C}^{(1)}$. Over the Schwartz Bruhat space $S(V_N(\mathbb{Q}_v))$ at v we also provide a description of ω_v as follows:

$$\omega_v\left(\begin{bmatrix} a & \\ & a^{-1} \end{bmatrix}\right)\Phi(x) = |a|_v^{\frac{N}{2}+1}\Phi(ax), \quad \omega_v\left(\begin{bmatrix} 1 & t \\ & 1 \end{bmatrix}\right)\Phi(x) = \psi_v(tq_Q(x))\Phi(x), \quad \omega_v\left(\begin{bmatrix} & 1 \\ -1 & \end{bmatrix}\right)\Phi(x) = \hat{\Phi}(x). \quad (16)$$

In general, we have to add a factor $\gamma_{q_Q, v}^{-1}(1)$, which denotes the local constant called the Weil constant, to the right hand side of the last equation above. But in our case, since L_p is self-dual for all $p < \infty$ and dimension of V_N is $N+2$, we see that $\gamma_{q_Q, v}(1) = 1$ for all places v (See [56, Chapitre II] for details).

Suppose $\tau = u + \sqrt{-1}v$ lies in the complex upper half plane, and let $g_\tau = \begin{bmatrix} 1 & u \\ & 1 \end{bmatrix} \begin{bmatrix} v^{1/2} & \\ & v^{-1/2} \end{bmatrix} \in \mathrm{SL}_2(\mathbb{R})$. Then

$$\begin{aligned} (\omega(g_\tau)\tilde{\phi}_\infty)(\lambda, \nu) &= v^{\frac{N}{4}+\frac{1}{2}} \exp(2\pi\sqrt{-1}uq_Q(\lambda))\tilde{\phi}_\infty(\sqrt{v}\lambda, \nu) \\ &= v^{\frac{N}{4}+\frac{1}{2}} \exp(2\pi\sqrt{-1}u(q_Q(\lambda_{\nu+}) + q_Q(\lambda_{\nu-}))) \exp(2\pi vq_Q(\lambda_{\nu-}) - 2\pi vq_Q(\lambda_{\nu+})) \\ &= v^{\frac{N}{4}+\frac{1}{2}} \exp(2\pi\sqrt{-1}(\tau q_Q(\lambda_{\nu+}) + \bar{\tau} q_Q(\lambda_{\nu-}))). \end{aligned} \quad (17)$$

The $\mathrm{SL}_2(\mathbb{A})$ action commutes with the action of $\mathcal{G}(\mathbb{A})$, which we denote by $\omega(h)\Phi(x) := \Phi(h^{-1}x)$. For $\nu \in \mathcal{D}$, $h \in \mathcal{G}(\mathbb{A}_f)$ and $g \in \mathrm{SL}_2(\mathbb{A})$, let $\theta(g, \nu, h)$ be the linear functional on $S(V_N(\mathbb{A}_f))$ defined by

$$S(V_N(\mathbb{A}_f)) \ni \phi \mapsto \theta(g, \nu, h; \phi) := \sum_{\lambda \in V_N(\mathbb{Q})} \omega(g) \left(\phi_\infty(\cdot, \nu) \otimes \omega(h)\phi \right) (\lambda). \quad (18)$$

Lemma. 3.2. *Let $h_0 \in \mathcal{G}(\mathbb{Q})$ and $g_0 \in \mathrm{SL}_2(\mathbb{Q})$. We have*

$$\theta(g, h_0\nu, h_0h; \phi) = \theta(g, \nu, h; \phi), \quad \theta(g_0g, \nu, h; \phi) = \theta(g, \nu, h; \phi). \quad (19)$$

Proof. The statement for h_0 follows from Lemma 3.1, and a change of variable. For the statement of g_0 , we can look at the three cases. For $g_0 = \begin{bmatrix} 1 & t \\ & 1 \end{bmatrix}$, the result follows from $\psi|_{\mathbb{Q}} \equiv 1$. For $g_0 = \begin{bmatrix} a & \\ & a^{-1} \end{bmatrix}$, we get the result from a change of variable $\lambda \mapsto a\lambda$. For $g_0 = \begin{bmatrix} & 1 \\ -1 & \end{bmatrix}$, the result is obtained by Poisson summation. $\square$

If $g_1 \in \mathrm{SL}_2(\mathbb{A}_f)$ and $h_1 \in \mathcal{G}(\mathbb{A}_f)$, then we have

$$\theta(gg_1, \nu, hh_1; \phi) = \theta(g, \nu, h; \omega(g_1, h_1)\phi), \quad (20)$$

with $\omega(g_1, h_1)$ meaning $\omega(g_1)\omega(h_1)$. Hence, if $\phi \in S(V_N(\mathbb{A}_f))^{K_f}$, then the map

$$(\nu, h) \mapsto \theta(g, \nu, h; \phi)$$

on $\mathcal{D} \times \mathcal{G}(\mathbb{A}_f)$ descends to a function on $X = \Gamma_S \backslash \mathcal{D}^+$. We may view it as a linear functional on $S(V_N(\mathbb{A}_f))^{K_f}$, and obtain

$$\theta : \mathrm{SL}_2(\mathbb{Q}) \backslash \mathrm{SL}_2(\mathbb{A}) \times X \rightarrow \left(S(V_N(\mathbb{A}_f))^{K_f} \right)^\vee \quad (g, \nu, h) \mapsto \theta(g, \nu, h; \cdot).$$

Since we have assumed that L is even unimodular, it is self dual, and in this case $S(V_N(\mathbb{A}_f))^{K_f}$ is a one dimensional space spanned by the characteristic function ϕ_0 of $\otimes_{p < \infty} L_{0,p}$.

Let $f \in S(\mathrm{SL}_2(\mathbb{Z}); -\frac{r^2+1}{4})$ be as in Theorem 2.2. Write $g \in \mathrm{SL}_2(\mathbb{A})$ as $g = \gamma g_\infty k$ with $\gamma \in \mathrm{SL}_2(\mathbb{Q})$, $g_\infty \in \mathrm{SL}_2(\mathbb{R})$ and $k \in \prod_\ell \mathrm{SL}_2(\mathbb{Z}_\ell)$. Define $f_0 : \mathrm{SL}_2(\mathbb{A}) \rightarrow \mathbb{C}$ be $f_0(g) := f(g_\infty \langle \sqrt{-1} \rangle)$.

As a function of $g \in \mathrm{SL}_2(\mathbb{A})$, the function $g \mapsto f_0(g)\theta(g, \nu, h; \phi_0)$ is left- $\mathrm{SL}_2(\mathbb{Q})$ invariant and right- $\mathrm{SO}(2, \mathbb{R}) \prod_\ell \mathrm{SL}_2(\mathbb{Z}_\ell)$ invariant using (19), (20) and $\hat{\phi}_0 = \phi_0$ since L is self-dual. This allows us to define the adelic Borchers lift on X by

$$\Phi(\nu, h, f_0) := \int_{\mathrm{SL}_2(\mathbb{Q}) \backslash \mathrm{SL}_2(\mathbb{A})} f_0(g) \bar{\theta}(g, \nu, h; \phi_0) dg. \quad (21)$$

Proposition. 3.3. *For $(x, y) \in H_N$, let $\nu(x, y) \in \mathcal{D}^+$ by (12). Then we have*

$$\Phi(\nu(x, y), 1, f_0) = F_f(n(x)a_y).$$

Proof. This follows from Section 3 of [37]. $\square$

4 Petersson norm of the theta lifting

We study the Petersson norm

$$\|\Phi(*, *, f_0)\|^2 := \int_{\mathcal{G}(\mathbb{Q}) \backslash \mathcal{G}(\mathbb{A})} \Phi(\nu, h, f_0) \overline{\Phi(\nu, h, f_0)} d\nu dh$$

of $\Phi(\nu, h, f_0)$ by the well known approach originally due to S. Rallis (cf. [45]). In view of Lemma 2.1 and Proposition 3.3 this coincides with

$$\|F_f\|_2^2 := \int_{\Gamma_S \backslash G_\infty} F_f(h_\infty) \overline{F_f(h_\infty)} dh_\infty,$$

which is nothing but the Petersson norm of our lift in the classical setting. This is convergent since $\Phi(\nu, h, f_0)$ is cuspidal.

We have that

$$\begin{aligned} \|\Phi(*, *, f_0)\|^2 &= \int_{\mathcal{G}(\mathbb{Q}) \backslash \mathcal{G}(\mathbb{A})} \int_{(\mathrm{SL}_2(\mathbb{Q}) \backslash \mathrm{SL}_2(\mathbb{A}))^2} f_0(g_1) \bar{\theta}(g_1, \nu, h; \phi_0) \overline{f_0(g_2) \bar{\theta}(g_2, \nu, h; \phi_0)} dg_1 dg_2 d\nu dh \\ &= \int_{(\mathrm{SL}_2(\mathbb{Q}) \backslash \mathrm{SL}_2(\mathbb{A}))^2} f_0(g_1) \overline{f_0(g_2)} I(g_1, g_2, \phi_0) dg_1 dg_2, \end{aligned} \quad (22)$$

where

$$I(g_1, g_2, \phi_0) := \int_{\mathcal{G}(\mathbb{Q}) \backslash \mathcal{G}(\mathbb{A})} \bar{\theta}(g_1, \nu, h; \phi_0) \theta(g_2, \nu, h; \phi_0) d\nu dh.$$

Since we are in the convergent range, the change in order of integration is justified. In view of the doubling variables of the Weil representation (cf. [26, Section 11]) we have

$$\bar{\theta}(g_1, \nu, h, \phi_0) \theta(g_2, \nu, h, \phi_0) = \theta((g_1, g_2), \delta(\nu), \delta(h), \phi_0 \otimes \phi_0). \quad (23)$$

Here, for $g_1, g_2 \in \mathrm{SL}_2$ we regard $(g_1, g_2) \in \mathrm{SL}_2 \times \mathrm{SL}_2$ as its image of the diagonal embedding $\mathrm{SL}_2 \times \mathrm{SL}_2 \hookrightarrow \mathrm{Sp}_4$ given by

$$\mathrm{SL}_2 \times \mathrm{SL}_2 \ni \left(\begin{bmatrix} a & b \\ c & d \end{bmatrix}, \begin{bmatrix} a' & b' \\ c' & d' \end{bmatrix} \right) \mapsto \begin{bmatrix} a & & -b & \\ & a' & & b' \\ -c & & d & \\ & c' & & d' \end{bmatrix} \in \mathrm{Sp}_4,$$

and the map δ denotes the canonical diagonal embedding of $\mathcal{G}$ into the orthogonal group defined by the quadratic space $(\mathbb{Q}^{N+2} \oplus \mathbb{Q}^{N+2}, q_Q \oplus q_Q)$, for which note that $\mathrm{Sp}_4 \times \delta(\mathcal{G})$ forms a dual pair.

On the right hand side of (23), the theta series is a linear functional on $S(V_N(\mathbb{A})) \oplus S(V_N(\mathbb{A}))$ with the choice of Schwartz function being $\Phi_0 := (\phi_\infty \oplus \phi_\infty) \otimes (\phi_0 \oplus \phi_0)$.

By the convergent Siegel-Weil formulas [57, Théorème 5], the integral $I(g_1, g_2, \phi_0)$ can be written as a special value of a certain Siegel Eisenstein series. Let us describe this next. Let $P(\mathbb{A}) = N(\mathbb{A})M(\mathbb{A})$ be the adelized Siegel parabolic subgroup of $\mathrm{Sp}_4(\mathbb{A})$ with the Levi part $M(\mathbb{A}) \simeq \mathrm{GL}_2(\mathbb{A})$ and the unipotent radical $N(\mathbb{A}) \simeq \mathrm{Sym}_2(\mathbb{A})$, and let $\tilde{K}$ be the standard maximal compact subgroup $\prod_{p < \infty} \mathrm{Sp}_4(\mathbb{Z}_p) \times (\mathrm{O}(4)(\mathbb{R}) \cap \mathrm{Sp}_4(\mathbb{R}))$. The Weil representation ω^D of $\mathrm{Sp}_4(\mathbb{A})$ on $S(V_N(\mathbb{A})^2)$, obtained by the doubling of ω , is realized as follows:

$$\begin{aligned} \omega^D \left(\begin{bmatrix} A & 0_2 \\ 0_2 & {}^t A^{-1} \end{bmatrix} \right) \Phi(X) &= |\det(A)|^{\frac{N}{2}+1} \Phi(XA) \quad (A \in \mathrm{GL}_2(\mathbb{A})), \\ \omega^D \left(\begin{bmatrix} 1_2 & Y \\ 0_2 & 1_2 \end{bmatrix} \right) \Phi(X) &= \psi \left(\frac{1}{2} \mathrm{tr}((X, X) \cdot Y) \right) \Phi(X) \quad (Y \in \mathrm{Sym}_2(\mathbb{A})), \\ \omega^D \left(\begin{bmatrix} 0_2 & 1_2 \\ -1_2 & 0_2 \end{bmatrix} \right) \Phi(X) &= \hat{\Phi}(X), \end{aligned} \quad (24)$$

where Φ denotes a Schwartz Bruhat function on $V_N(\mathbb{A}) \oplus V_N(\mathbb{A})$, and $(X, X) := (\frac{1}{2} {}^t x_i Q x_j) \in \mathrm{Sym}_2(\mathbb{A})$ for $X = (x_1, x_2) \in V_N(\mathbb{A})^2$. For the first formula we note that there is a factor $\chi_{q_Q}(\det(A))$, which is proved to be trivial.

We now also provide local representations ω_v^D of ω^D at each $v \leq \infty$. With the same notation for ω_v

$$\omega_v^D \left(\begin{bmatrix} A & 0_2 \\ 0_2 & {}^t A^{-1} \end{bmatrix} \right) \Phi(X) = |\det(A)|^{\frac{N}{2}+1} \Phi(XA) \quad (A \in \mathrm{GL}_2(\mathbb{Q}_v)),$$

$$\begin{aligned}\omega_v^D\left(\begin{bmatrix} 1_2 & Y \\ 0_2 & 1_2 \end{bmatrix}\right)\Phi(X) &= \psi\left(\frac{1}{2}\mathrm{tr}((X, X) \cdot Y)\right)\Phi(X) \quad (Y \in \mathrm{Sym}_2(\mathbb{Q}_v)), \\ \omega_v^D\left(\begin{bmatrix} 0_2 & 1_2 \\ -1_2 & 0_2 \end{bmatrix}\right)\Phi(X) &= \hat{\Phi}(X),\end{aligned}\tag{25}$$

where Φ denotes a Schwartz Bruhat function on $V_N(\mathbb{Q}_v) \oplus V_N(\mathbb{Q}_v)$, and $(X, X) := (\frac{1}{2}x_i Q x_j) \in \mathrm{Sym}_2(\mathbb{Q}_v)$ for $X = (x_1, x_2) \in V_N(\mathbb{Q}_v)^2$. For the formula above we remark that there is the factor $\chi_{q_{Q,v}}(\det(A))$ (resp. $\gamma_{q_{Q,v}}(1)^{-2}$) in the first formula (resp. third formula), which turn out to be trivial.

We have the Iwasawa decomposition $\mathrm{Sp}_4(\mathbb{A}) = P(\mathbb{A})\tilde{K}$. Write any $g \in \mathrm{Sp}_4(\mathbb{A})$ as $g = nm(a)k$ with some $n = \begin{pmatrix} 1_2 & X \\ 0_2 & 1_2 \end{pmatrix}$ ($X \in \mathrm{Sym}_2(\mathbb{A})$), $m(a) = \begin{pmatrix} a & 0_2 \\ 0_2 & {}^t a^{-1} \end{pmatrix}$ ($a \in \mathrm{GL}_2(\mathbb{A})$) and $k \in \tilde{K}$. We then set $|a(g)| := |\det(a)|_{\mathbb{A}}$. This is a well-defined function on $\mathrm{Sp}_4(\mathbb{A})$ that is left $N(\mathbb{A})M(\mathbb{Q})$ -invariant and right $\tilde{K}$ -invariant. For $s \in \mathbb{C}$ and $\Phi \in S(V_N(\mathbb{A})^2)$, define $\Xi(g, s) := \omega^D(g)\Phi(0) \cdot |a(g)|^{s-s_0}$, where $s_0 = (N-1)/2$. From (24), we see that $\omega^D(nm(a)g)\Phi(0) = |\det(a)|^{\frac{N}{2}+1}\omega^D(g)\Phi(0)$. Recall that the modular character on the Siegel parabolic is $\delta_P(p) = |a(p)|^3$. Hence, we get

$$\Xi(pg, s) = |a(p)|^s \delta_P(p)^{\frac{1}{2}} \Xi(g, s) \text{ for } p \in P(\mathbb{A}), g \in \mathrm{Sp}_4(\mathbb{A}).$$

Hence we can conclude that $\Xi \in \mathrm{Ind}_{P(\mathbb{A})}^{\mathrm{Sp}_4(\mathbb{A})}(\delta_P^{\frac{s}{3}})$, which is the normalized parabolic induction. For any $F \in \mathrm{Ind}_{P(\mathbb{A})}^{\mathrm{Sp}_4(\mathbb{A})}(\delta_P^{\frac{s}{3}})$, the Siegel Eisenstein series is defined by

$$E(g, s; F) := \sum_{\gamma \in P(\mathbb{Q}) \backslash \mathrm{Sp}_4(\mathbb{Q})} F(\gamma g, s),$$

which converges absolutely for $\mathrm{Re}(s) > 3/2$ (cf. [57, Théorème 1]).

We now state the Siegel Weil formula for our case as follows (cf. [57, Théorème 5]):

Proposition. 4.1. *Let $\Phi_0 \in S(V_N(\mathbb{A})) \oplus S(V_N(\mathbb{A}))$ be defined by $(\phi_\infty \oplus \phi_\infty) \otimes (\phi_0 \oplus \phi_0)$ as above. For $s \in \mathbb{C}$, $g \in \mathrm{Sp}_4(\mathbb{A})$, let $\Xi_0(g, s) := \omega^D(g)\Phi_0(0)|a(g)|^{s-s_0}$, where $s_0 = (N-1)/2$. Then, for $g_1, g_2 \in \mathrm{SL}_2(\mathbb{A})$, we have*

$$I(g_1, g_2, \phi_0) = E((g_1, g_2), s_0; \Xi_0).\tag{26}$$

Returning to the Petersson norm calculation (22), we now have

$$\|\Phi(*, *, f_0)\|^2 = \int_{\mathrm{SL}_2(\mathbb{Q}) \backslash \mathrm{SL}_2(\mathbb{A})} \left(\int_{\mathrm{SL}_2(\mathbb{Q}) \backslash \mathrm{SL}_2(\mathbb{A})} f_0(g_1) E((g_1, g_2), s_0; \Xi_0) dg_1 \right) \overline{f_0(g_2)} dg_2.\tag{27}$$

Call the inner integral $Z(s_0, f_0, \Xi_0)(g_2)$. By Theorem 3.6 of [44], we have

$$Z(s_0, f_0, \Xi_0)(g_2) = \prod_{v \leq \infty} Z_v(s_0, f_v, \Xi_v)(g_v),$$

where

$$Z_v(s_0, f_v, \Xi_v)(g_v) = \int_{\mathrm{SL}_2(\mathbb{Q}_v)} \Xi_v(Q_2(h, 1), s_0) f_v(g_v h) dh\tag{28}$$

with local components $\{f_v\}$ of f_0 . Here

$$Q_2 := \begin{bmatrix} 1 & & & \\ & 1 & & \\ & & 1 & \\ 1 & -1 & & \end{bmatrix}.$$

For $v < \infty$, all the data is unramified, and hence, by Proposition 4.1 of [44], we get

$$Z_v(s_0, f_v, \Xi_v)(g_v) = \frac{L_v(s_0 + \frac{1}{2}, \pi_v, \text{Ad})}{\zeta_v(s_0 + \frac{3}{2})\zeta_v(2s_0 + 1)} f_v(g_v). \quad (29)$$

Recall that $\pi_f = \otimes'_v \pi_v$ is the automorphic cuspidal representation of $\text{GL}_2(\mathbb{A})$ generated by f_0 . Here, $L(s, \pi_v, \text{Ad})$ is the degree 3 adjoint L -function of π_v .

4.1 The archimedean computation

In this section we will compute the archimedean integral $Z_\infty(s_0, f_\infty, \Xi_\infty)$. We begin with the following proposition.

Proposition. 4.2. *Let $\tilde{K}_\infty$ be the maximal compact subgroup of $\text{Sp}_4(\mathbb{R})$ given by $\tilde{K}_\infty := \text{Sp}_4(\mathbb{R}) \cap \text{O}_4(\mathbb{R})$, which is isomorphic to the unitary group $U(2)$ of degree two. For the Schwartz functions ϕ_∞ in (14), set $\Phi_\infty := \phi_\infty \oplus \phi_\infty$. Then $\Xi_\infty(g, s_0) := \omega^D(g)\Phi_\infty(0)$ is right $\tilde{K}_\infty$ -invariant (with respect to the Weil representation).*

Proof. For the proof we need the following lemma.

Lemma. 4.3. *i) The Schwartz function ϕ_∞ in (14) is right $\text{SO}(2)$ -invariant (with respect to the Weil representation).*

ii) Let τ_Λ be an irreducible representation of $\tilde{K}_\infty \simeq U(2)$ with dominant weight $\Lambda = (\lambda_1, \lambda_2) \in \mathbb{Z}^2$ with $\lambda_1 \geq \lambda_2$. The infinitesimal action of τ_Λ , also denoted by τ_Λ , is described by the following explicit formula:

$$\begin{aligned} \tau_\Lambda(H_1)v_k &= (\lambda_2 + k)v_k, \quad \tau_\Lambda(H_2)v_k = (\lambda_1 - k)v_k, \\ \tau_\Lambda(X)v_k &= (k+1)v_{k+1}, \quad \tau_\Lambda(\bar{X})v_k = (d - (k-1))v_{k-1}, \end{aligned}$$

where

- $\{v_k\}_{0 \leq k \leq d := \lambda_1 - \lambda_2}$ denotes a set of weight vectors which forms a basis of the representation space for τ_Λ .

•

$$\begin{aligned} H_1 &:= \begin{bmatrix} 0 & 0 & -\sqrt{-1} & 0 \\ 0 & 0 & 0 & 0 \\ \sqrt{-1} & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{bmatrix}, \quad H_2 := \begin{bmatrix} 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & -\sqrt{-1} \\ 0 & 0 & 0 & 0 \\ 0 & \sqrt{-1} & 0 & 0 \end{bmatrix}, \\ X &:= \frac{1}{2} \begin{bmatrix} 0 & 1 & 0 & -\sqrt{-1} \\ -1 & 0 & -\sqrt{-1} & 0 \\ 0 & \sqrt{-1} & 0 & 1 \\ \sqrt{-1} & 0 & -1 & 0 \end{bmatrix}, \end{aligned}$$

and $\bar{X}$ denotes the complex conjugate of X .

Proof. (i) Putting $\nu_0 = \nu(0, 1)$ we see

$$\phi_\infty(\lambda, \nu_0) = P(\lambda, \nu_0) \exp(-2\pi q_R(\lambda)),$$

where q_R denotes the quadratic form defined by $R := \begin{pmatrix} 1 & & \\ & S & \\ & & 1 \end{pmatrix}$ (cf. Section 2.1). It suffices to calculate

the action of $\begin{bmatrix} \cos \theta & \sin \theta \\ -\sin \theta & \cos \theta \end{bmatrix}$ on $\phi_\infty(\lambda, \nu_0)$ in (14) via the Weil representation. In fact, we see that the

calculation does not depend on the choice of $\nu \in \mathcal{D}$. This is due to the well known commutativity of the actions of the dual pair by the Weil representation, which is also ensured by the argument of Section 3. The main difficulty is the calculation of the action of $\begin{bmatrix} 0 & 1 \\ -1 & 0 \end{bmatrix}$ on ϕ_∞ . We remark that such a calculation is well known for the Gaussian multiplied by a harmonic polynomial. For instance we can find it in the proof of the automorphy for holomorphic theta series with harmonic polynomials.

When the matrix $\begin{bmatrix} \cos \theta & \sin \theta \\ -\sin \theta & \cos \theta \end{bmatrix}$ is diagonal i.e. $\sin \theta = 0$ the calculation is settled in a straight forward manner. Let us thus assume $\sin \theta \neq 0$. We then have that

$$\begin{bmatrix} \cos \theta & \sin \theta \\ -\sin \theta & \cos \theta \end{bmatrix} = \begin{bmatrix} 1 & -\frac{1}{\tan \theta} \\ 0 & 1 \end{bmatrix} \begin{bmatrix} 0 & 1 \\ -1 & 0 \end{bmatrix} \begin{bmatrix} \sin \theta & 0 \\ 0 & \frac{1}{\sin \theta} \end{bmatrix} \begin{bmatrix} 1 & -\frac{1}{\tan \theta} \\ 0 & 1 \end{bmatrix}.$$

The action by $\begin{bmatrix} \sin \theta & 0 \\ 0 & \frac{1}{\sin \theta} \end{bmatrix} \begin{bmatrix} 1 & -\frac{1}{\tan \theta} \\ 0 & 1 \end{bmatrix}$ is calculated to be

$$|\sin \theta|^{\frac{N}{2}+1} \exp(2\pi\sqrt{-1}(-\frac{1}{\tan \theta}q_Q(\sin \theta \cdot \lambda))(\exp(-\frac{\Delta}{8\pi})(-2^{\frac{N}{2}-4}(\sin \theta \cdot x_1)^{\frac{N}{2}})) \exp(-2\pi q_R(\sin \theta \cdot \lambda)),$$

where x_1 denotes the first entry of $\lambda \in \mathbb{R}^{N+2}$, more precisely the coordinate of $\mathbb{R} \cdot \nu_0 = \{\frac{1}{\sqrt{2}}(x_1, 0, x_1) \mid x_1 \in \mathbb{R}\}$. This is deduced immediately from the definition of the Weil representation (cf. (16)).

As the next step we consider the action by $\begin{bmatrix} 0 & 1 \\ -1 & 0 \end{bmatrix}$, which is given by the Fourier transform of the above expression multiplied by the Weil constant. For this purpose we introduce

$$\phi_{\tau, \infty}(x) := (\exp(-\frac{\Delta}{8\pi \operatorname{Im}(\tau)})(2^{\frac{N}{2}-3}x_1^{\frac{N}{2}})) \exp(2\pi\sqrt{-1}(\operatorname{Re}(\tau)q_Q(\lambda) + \operatorname{Im}(\tau)\sqrt{-1}q_R(\lambda))) \quad (\tau \in \mathfrak{h}).$$

The result of the action by $\begin{bmatrix} \sin \theta & 0 \\ 0 & \frac{1}{\sin \theta} \end{bmatrix} \begin{bmatrix} 1 & -\frac{1}{\tan \theta} \\ 0 & 1 \end{bmatrix}$ is rewritten as

$$|\sin \theta|^{N+1} \phi_{-\cos \theta \sin \theta + \sqrt{-1} \sin^2 \theta, \infty}(x).$$

From [4, Corollary 3.5] the Fourier transform of $\phi_{\tau, \infty}$ is as follows:

$$\widehat{\phi_{\tau, \infty}}(x) = (\tau/\sqrt{-1})^{-\frac{1}{2}}(\sqrt{-1}\bar{\tau})^{-\frac{N+1}{2}}(-\tau)^{-\frac{N}{2}}\phi_{-1/\tau, \infty}(x).$$

Putting $\tau = -\cos \theta \sin \theta + \sqrt{-1} \sin^2 \theta$ the result of the action by $\begin{bmatrix} 0 & 1 \\ -1 & 0 \end{bmatrix}$ is calculated to be

$$\phi_{-1/\tau, \infty}(x),$$

taking into account the Weil constant for q_Q and the constant factor of the self-dual measure, both of which are trivial for an even unimodular lattice. As the last step, the action of $\begin{bmatrix} 1 & -\frac{1}{\tan \theta} \\ 0 & 1 \end{bmatrix}$ is given by

$$\phi_{-1/\tau - \frac{1}{\tan \theta}, \infty}(x) = \phi_{\sqrt{-1}, \infty} = \phi_\infty.$$

(ii) Let $\mathfrak{k}$ and $\mathfrak{u}(2)$ be the Lie algebra of K and $U(2)$ respectively, and note the following isomorphism

$$\mathfrak{k} \ni \begin{bmatrix} A & B \\ -B & A \end{bmatrix} \mapsto A + \sqrt{-1}B \in \mathfrak{u}(2)$$

as Lie algebras. This isomorphism is naturally extended to the isomorphism between their complexifications: $\mathfrak{k}_{\mathbb{C}} \simeq \mathfrak{u}(2)_{\mathbb{C}}$, the latter of which coincides with the complex matrix algebra $M_2(\mathbb{C})$ of degree two.

Via this isomorphism, H_1 , H_2 , X , and $\bar{X}$ correspond to $\begin{bmatrix} 1 & 0 \\ 0 & 0 \end{bmatrix}$, $\begin{bmatrix} 0 & 0 \\ 0 & 1 \end{bmatrix}$, $\begin{bmatrix} 0 & 1 \\ 0 & 0 \end{bmatrix}$, and $\begin{bmatrix} 0 & 0 \\ 1 & 0 \end{bmatrix}$ respectively. Recall that the irreducible representation of $U(2)$ with highest weight $\Lambda = (\lambda_1, \lambda_2)$ is realized by $\det^{\lambda_2} \text{sym}^{\lambda_1 - \lambda_2}(\mathbb{R}^2)$, where $\text{sym}^{\lambda_1 - \lambda_2}(\mathbb{R}^2)$ denotes the $(\lambda_1 - \lambda_2)$ -th symmetric tensor representation of the standard representation of $U(2)$. The formula is obtained by considering the pullback of the infinitesimal action of the irreducible representation of $U(2)$ with highest weight Λ . $\square$

For the proof of Proposition 4.2, we note that $\Xi_\infty(g, s_0)$ is right $\tilde{K}_\infty$ -invariant if and only if $\Xi_\infty(g, s_0)$ belongs to the one-dimensional $\tilde{K}_\infty$ -type τ_Λ with dominant weight $\Lambda = (0, 0)$. By definition of the Lie algebra action, we have, for $i = 1, 2$

$$\tau_\Lambda(\sqrt{-1}H_i)\Xi_\infty(g, s_0) := \left. \frac{d}{d\theta} \right|_{\theta=0} \Xi_\infty(g \exp(\theta\sqrt{-1}H_i), s_0).$$

Since

$$\exp(\theta\sqrt{-1}H_1) = \begin{bmatrix} \cos(\theta) & 0 & \sin(\theta) & 0 \\ 0 & 1 & 0 & 0 \\ -\sin(\theta) & 0 & \cos(\theta) & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix}, \quad \exp(\theta\sqrt{-1}H_2) = \begin{bmatrix} 1 & 0 & 0 & 0 \\ 0 & \cos(\theta) & 0 & \sin(\theta) \\ 0 & 0 & 1 & 0 \\ 0 & -\sin(\theta) & 0 & \cos(\theta) \end{bmatrix},$$

part (i) of Lemma 4.3 implies that $\tau_\Lambda(\sqrt{-1}H_i)\Xi_\infty(g, s_0) = 0$ for $i = 1, 2$. Hence, by part (ii) of Lemma 4.3, we see that $\Xi_\infty(g, s_0)$ is the weight $(0, 0)$ vector in $\tau_{(\lambda, -\lambda)}$ for $\lambda \geq 0$. In the terminology of part (ii) of Lemma 4.3, $\Xi_\infty(g, s_0)$ is the vector v_λ in the basis of $\tau_{(\lambda, -\lambda)}$.

For $\theta \in \mathbb{R}$, set $r(\theta) := \begin{bmatrix} \cos(\theta) & \sin(\theta) \\ -\sin(\theta) & \cos(\theta) \end{bmatrix}$ and $R(\theta) = \begin{bmatrix} r(\theta) & \\ & \iota_{r(\theta)^{-1}} \end{bmatrix}$. By definition of the local Weil representation (25), we see that $(\omega^D(R(\theta))\Phi_\infty)(Y) = \Phi_\infty(Yr(\theta))$. Note that, if $Y = (y_1, y_2) \in V_N(\mathbb{R})^2$, then $Yr(\theta) = (\cos(\theta)y_1 - \sin(\theta)y_2, \sin(\theta)y_1 + \cos(\theta)y_2)$. Using the definition of ϕ_∞ and the fact that

$$q_Q(\cos(\theta)y_1 - \sin(\theta)y_2) + q_Q(\sin(\theta)y_1 + \cos(\theta)y_2) = q_Q(y_1) + q_Q(y_2),$$

we see that $\Phi_\infty(Yr(\theta)) = \Phi_\infty(Y)$. Hence we see that $\Xi_\infty(gR(\theta), s_0) = \Xi_\infty(g, s_0)$ for all g and θ . Since $\exp(\sqrt{-1}\theta(X + \bar{X})) = R(\theta)$, this implies that $\tau_\Lambda(\sqrt{-1}(X + \bar{X}))\Xi_\infty = 0$. By part (ii) of Lemma 4.3, we can conclude that $\lambda = 0$, as required. This completes the proof of the proposition. $\square$

We will now compute the archimedean integral (28)

$$Z_\infty(s_0, f_\infty, \Xi_\infty)(g_2) = \int_{\text{SL}_2(\mathbb{R})} \Xi_\infty(Q_2(h, 1), s_0) f_\infty(g_2 h) dh.$$

Note that π_∞ is the irreducible principle series of $\text{GL}_2^+(\mathbb{R})$ with all even $\text{SO}(2)$ -types, and the Maass cusp form f is viewed as the weight 0 vector f_∞ in π_∞ . Hence, in the induced model for π_∞ , we have

$$f_\infty\left(\begin{bmatrix} a & * \\ & a^{-1} \end{bmatrix} k\right) = a^{\sqrt{-1}r+1}, \quad a \in \mathbb{R}^+, k \in \text{SO}(2, \mathbb{R}). \quad (30)$$

Here r depends on the Laplace eigenvalue of the Maass form f . By Proposition 4.2, the local section Ξ_∞ is right $\tilde{K}_\infty$ -invariant. The local integral Z_∞ is an element of π_∞ and a simple change of variable shows that it is also right $\text{SO}(2)$ -invariant. Since, f_∞ is the unique (up to scalars) element of π_∞ which is right $\text{SO}(2)$ -invariant, we see that

$$Z_\infty(s_0, f_\infty, \Xi_\infty)(g_2) = B(s_0) f_\infty(g_2).$$

The final task is to compute $B(s_0)$ which can be achieved by

$$B(s_0) = Z_\infty(s_0, f_\infty, \Xi_\infty)(1) = \int_{\text{SL}_2(\mathbb{R})} \Xi_\infty(Q_2(h, 1), s_0) f_\infty(h) dh.$$

Using the Iwasawa decomposition, we see that, if u is a right $\mathrm{SO}(2)$ -invariant function on $\mathrm{SL}_2(\mathbb{R})$, then

$$\int_{\mathrm{SL}_2(\mathbb{R})} u(g) dg = 2\pi \int_0^\infty \int_{-\infty}^\infty u\left(\begin{bmatrix} a & \\ & a^{-1} \end{bmatrix} \begin{bmatrix} 1 & x \\ & 1 \end{bmatrix}\right) a^{-1} dx da.$$

Hence, we get

$$\begin{aligned} B(s_0) &= 2\pi \int_0^\infty \int_{-\infty}^\infty \Xi_\infty(Q_2(\begin{bmatrix} a & \\ & a^{-1} \end{bmatrix} \begin{bmatrix} 1 & x \\ & 1 \end{bmatrix}, 1), s_0) f_\infty(\begin{bmatrix} a & \\ & a^{-1} \end{bmatrix} \begin{bmatrix} 1 & x \\ & 1 \end{bmatrix}) a^{-1} dx da \\ &= 2\pi \int_0^\infty \int_{-\infty}^\infty \Xi_\infty(Q_2(\begin{bmatrix} a & \\ & a^{-1} \end{bmatrix} \begin{bmatrix} 1 & x \\ & 1 \end{bmatrix}, 1), s_0) a^{\sqrt{-1}r} dx da. \end{aligned}$$

We need to first simplify the integrand. For this, note that

$$Q_2 = \begin{bmatrix} 1 & & & \\ & 1 & & \\ & 1 & 1 & \\ 1 & & & 1 \end{bmatrix} \begin{bmatrix} 1 & & & \\ & 1 & & \\ & & 1 & \\ & -1 & & 1 \end{bmatrix}.$$

Hence, we have

$$\begin{aligned} &\Xi_\infty(Q_2(\begin{bmatrix} a & \\ & a^{-1} \end{bmatrix} \begin{bmatrix} 1 & x \\ & 1 \end{bmatrix}, 1), s_0) \\ &= \Xi_\infty\left(\begin{bmatrix} 1 & & & \\ & 1 & & \\ & 1 & 1 & \\ 1 & & & 1 \end{bmatrix} \begin{bmatrix} 1 & & & \\ & 1 & & \\ & & 1 & \\ & -1 & & 1 \end{bmatrix} \begin{bmatrix} a & & -ax & \\ & 1 & & \\ & & a^{-1} & \\ & & & 1 \end{bmatrix}, s_0\right) \\ &= \Xi_\infty\left(\begin{bmatrix} a & & & \\ & 1 & & \\ & & a^{-1} & \\ & & & 1 \end{bmatrix} \begin{bmatrix} 1 & & & \\ & 1 & & \\ & a & 1 & \\ a & & & 1 \end{bmatrix} \begin{bmatrix} 1 & & & \\ & 1 & & \\ & & 1 & \\ & -1 & & 1 \end{bmatrix} \begin{bmatrix} 1 & & -x & \\ & 1 & & \\ & & 1 & \\ & & & 1 \end{bmatrix}, s_0\right) \\ &= a^{s_0+3/2} \Xi_\infty\left(\begin{bmatrix} 1 & & & \\ & 1 & & \\ & a & 1 & \\ a & & & 1 \end{bmatrix} \begin{bmatrix} 1 & & & \\ & 1 & & \\ & & 1 & \\ & -1 & & 1 \end{bmatrix} \begin{bmatrix} 1 & & -x & \\ & 1 & & \\ & & 1 & \\ & & & 1 \end{bmatrix}, s_0\right) \\ &= a^{s_0+3/2} \Xi_\infty\left(\begin{bmatrix} 1 & & & \\ & 1 & & \\ & a & 1 & \\ a & & & 1 \end{bmatrix} \begin{bmatrix} 1 & & -x & \\ & 1 & & \\ & & 1 & \\ & & & 1 \end{bmatrix} \begin{bmatrix} 1 & & & \\ & 1 & & \\ & & 1 & \\ & -1 & & 1 \end{bmatrix}, s_0\right) \\ &= a^{s_0+3/2} \Xi_\infty\left(\begin{bmatrix} 1 & & & \\ & 1 & & \\ & a & 1 & \\ a & & & 1 \end{bmatrix} \begin{bmatrix} 1 & & -x & \\ & 1 & & \\ & & 1 & \\ & & & 1 \end{bmatrix}, s_0\right). \end{aligned}$$

Here, we have used that Ξ_∞ lies in the induced representation and is right $\tilde{K}_\infty$ -invariant. Now, write

$$\begin{bmatrix} 1 & \\ a & 1 \end{bmatrix} = \begin{bmatrix} 1 & x' \\ & 1 \end{bmatrix} \begin{bmatrix} y^{1/2} & \\ & y^{-1/2} \end{bmatrix} r(\theta) \text{ with } x' = \frac{a}{1+a^2}, y = \frac{1}{1+a^2}, e^{\sqrt{-1}\theta} = \frac{1-\sqrt{-1}a}{\sqrt{1+a^2}}, r(\theta) = \begin{bmatrix} c(\theta) & s(\theta) \\ -s(\theta) & c(\theta) \end{bmatrix}.$$

Here, $c(\theta) = \cos(\theta)$ and $s(\theta) = \sin(\theta)$. Hence,

$$\begin{aligned}
& \Xi_\infty(Q_2([{}^a_{a^{-1}}][{}^1_x], 1), s_0) \\
&= a^{s_0+3/2} \Xi_\infty\left(\begin{bmatrix} 1 & & x' \\ & 1 & x' \\ & & 1 \end{bmatrix} \begin{bmatrix} y^{1/2} & & & \\ & y^{1/2} & & \\ & & y^{-1/2} & \\ & & & y^{-1/2} \end{bmatrix} \begin{bmatrix} c(\theta) & & s(\theta) \\ & c(\theta) & s(\theta) \\ & -s(\theta) & c(\theta) \\ -s(\theta) & & c(\theta) \end{bmatrix} \right. \\
&\quad \left. \begin{bmatrix} 1 & & -x \\ & 1 & \\ & & 1 \\ & & & 1 \end{bmatrix}, s_0\right) \\
&= \frac{a^{s_0+3/2}}{(1+a^2)^{s_0+3/2}} \Xi_\infty\left(\begin{bmatrix} c(\theta) & & s(\theta) \\ & c(\theta) & s(\theta) \\ & -s(\theta) & c(\theta) \\ -s(\theta) & & c(\theta) \end{bmatrix} \begin{bmatrix} 1 & & -x \\ & 1 & \\ & & 1 \\ & & & 1 \end{bmatrix}, s\right) \\
&= \left(\frac{a}{1+a^2}\right)^{s_0+3/2} \Xi_\infty\left(\begin{bmatrix} 1 & -s(\theta)c(\theta)x & -c(\theta)^2x \\ & 1 & \\ & s(\theta)^2x & s(\theta)c(\theta)x \\ & & 1 \end{bmatrix} \begin{bmatrix} c(\theta) & & s(\theta) \\ & c(\theta) & s(\theta) \\ & -s(\theta) & c(\theta) \\ -s(\theta) & & c(\theta) \end{bmatrix}, s\right) \\
&= \left(\frac{a}{1+a^2}\right)^{s_0+3/2} \Xi_\infty\left(\begin{bmatrix} 1 & -s(\theta)c(\theta)x & -c(\theta)^2x \\ & 1 & \\ & s(\theta)^2x & s(\theta)c(\theta)x \\ & & 1 \end{bmatrix}, s\right) \\
&= \left(\frac{a}{1+a^2}\right)^{s_0+3/2} \Xi_\infty\left(\begin{bmatrix} 1 & -s(\theta)c(\theta)x & -c(\theta)^2x \\ & 1 & \\ & & 1 \\ & & s(\theta)c(\theta)x \end{bmatrix} \begin{bmatrix} 1 & & & \\ & 1 & & \\ & & 1 & \\ & s(\theta)^2x & & 1 \end{bmatrix}, s\right) \\
&= \left(\frac{a}{1+a^2}\right)^{s_0+3/2} \Xi_\infty\left(\begin{bmatrix} 1 & & & \\ & 1 & & \\ & & 1 & \\ & s(\theta)^2x & & 1 \end{bmatrix}, s\right).
\end{aligned}$$

Note that $s(\theta)^2 = a^2/(1+a^2)$. Substituting this in the integral for $B(s_0)$, we get

$$B(s_0) = 2\pi \int_0^\infty \int_{-\infty}^\infty \left(\frac{a}{1+a^2}\right)^{s_0+3/2} \Xi_\infty\left(\begin{bmatrix} 1 & & & \\ & 1 & & \\ & & 1 & \\ & s(\theta)^2x & & 1 \end{bmatrix}, s_0\right) a^{\sqrt{-1}r} dx da.$$

A change of variable $x \rightarrow s(\theta)^{-2}x$ gives us

$$\begin{aligned}
B(s_0) &= 2\pi \int_0^\infty \int_{-\infty}^\infty \left(\frac{a}{1+a^2}\right)^{s_0+3/2} \frac{1+a^2}{a^2} \Xi_\infty\left(\begin{bmatrix} 1 & & & \\ & 1 & & \\ & & 1 & \\ & x & & 1 \end{bmatrix}, s_0\right) a^{\sqrt{-1}r} dx da \\
&= 2\pi \int_0^\infty \frac{a^{s_0-1/2+\sqrt{-1}r}}{(1+a^2)^{s_0+1/2}} da \int_{-\infty}^\infty \Xi_\infty\left(\begin{bmatrix} 1 & & & \\ & 1 & & \\ & & 1 & \\ & x & & 1 \end{bmatrix}, s_0\right) dx.
\end{aligned}$$

Let us consider the integral in the a variable first. Change of variable $u = a^2$ gives us

$$\int_0^\infty \frac{a^{s_0-1/2+\sqrt{-1}r}}{(1+a^2)^{s_0+1/2}} da = \frac{1}{2} \int_0^\infty \frac{u^{\frac{s_0}{2}-\frac{3}{4}+\frac{\sqrt{-1}r}{2}}}{(1+u)^{s_0+1/2}} du.$$

Recall the beta function

$$B(v, w) = \frac{\Gamma(v)\Gamma(w)}{\Gamma(v+w)}.$$

One integral representation for the beta function (see pg 7 of [38]) is

$$B(v, w) = \int_0^\infty \frac{u^{v-1}}{(1+u)^{v+w}} du \text{ for } \operatorname{Re}(v) > 0, \operatorname{Re}(w) > 0.$$

We have $v = s_0/2 + 1/4 + \sqrt{-1}r/2$ and $w = s_0/2 + 1/4 - \sqrt{-1}r/2$. Hence, we get

$$\int_0^\infty \frac{a^{s_0-1/2+\sqrt{-1}r}}{(1+a^2)^{s_0+1/2}} da = \frac{1}{2} \frac{\Gamma(s_0/2 + 1/4 + \sqrt{-1}r/2) \Gamma(s_0/2 + 1/4 - \sqrt{-1}r/2)}{\Gamma(s_0 + 1/2)}.$$

Now, let us compute the integral in the x variable. We will again use the Iwasawa decomposition

$$\begin{bmatrix} 1 & \\ x & 1 \end{bmatrix} = \begin{bmatrix} 1 & x' \\ & y^{-1/2} \end{bmatrix} r(\theta) \text{ with } x' = \frac{x}{1+x^2}, y = \frac{1}{1+x^2}, e^{\sqrt{-1}\theta} = \frac{1 - \sqrt{-1}x}{\sqrt{1+x^2}}.$$

Hence

$$\begin{aligned} \int_{-\infty}^\infty \Xi_\infty \left(\begin{bmatrix} 1 & & \\ & 1 & \\ & x & 1 \end{bmatrix}, s_0 \right) dx &= \int_{-\infty}^\infty \Xi_\infty \left(\begin{bmatrix} 1 & & \\ & 1 & x' \\ & & 1 \end{bmatrix} \begin{bmatrix} 1 & & \\ & y^{1/2} & \\ & & 1 \end{bmatrix}, s_0 \right) dx \\ &= \int_{-\infty}^\infty \frac{1}{(1+x^2)^{\frac{s_0}{2}+\frac{3}{4}}} dx = 2 \int_0^\infty \frac{1}{(1+x^2)^{\frac{s_0}{2}+\frac{3}{4}}} dx. \end{aligned}$$

The change of variable $x = \tan(\theta)$ gives us

$$\int_{-\infty}^\infty \Xi_\infty \left(\begin{bmatrix} 1 & & \\ & 1 & \\ & x & 1 \end{bmatrix}, s_0 \right) dx = 2 \int_0^{\pi/2} \frac{1}{(\sec^2(\theta))^{\frac{s_0}{2}+\frac{3}{4}}} \sec^2(\theta) d\theta = 2 \int_0^{\pi/2} \cos(\theta)^{s_0-\frac{1}{2}} d\theta$$

On pg 8 of [38], we have the formula

$$\int_0^{\pi/2} \sin(\theta)^{2x-1} \cos(\theta)^{2y-1} d\theta = \frac{1}{2} B(x, y) \text{ for } \operatorname{Re}(x) > 0, \operatorname{Re}(y) > 0.$$

We have $x = 1/2$ and $y = s_0/2 + 1/4$. Hence, we have

$$\int_{-\infty}^\infty \Xi_\infty \left(\begin{bmatrix} 1 & & \\ & 1 & \\ & x & 1 \end{bmatrix}, s_0 \right) dx = B(1/2, s_0/2 + 1/4) = \frac{\Gamma(1/2)\Gamma(s_0/2 + 1/4)}{\Gamma(s_0/2 + 3/4)}.$$

Putting all this together, and using properties of the gamma function, we get

$$\begin{aligned} B(s_0) &= \pi \frac{\Gamma(s_0/2 + 1/4 + \sqrt{-1}r/2)\Gamma(s_0/2 + 1/4 - \sqrt{-1}r/2)}{\Gamma(s_0 + 1/2)} \frac{\Gamma(1/2)\Gamma(s_0/2 + 1/4)}{\Gamma(s_0/2 + 3/4)} \\ &= 2^{1-\frac{N}{2}} \pi^2 \frac{\Gamma(\frac{N}{4} + \frac{\sqrt{-1}r}{2})\Gamma(\frac{N}{4} - \frac{\sqrt{-1}r}{2})}{\Gamma(\frac{N}{4} + \frac{1}{2})^2}. \end{aligned}$$

Putting together (27), (29) and the formula of $B(s_0)$ above, we get the following theorem.

Theorem. 4.4. *The Petersson norm of the Borchers theta lift is given by*

$$\|\Phi(*, *, f_0)\|^2 = \frac{L(\frac{N}{2}, \pi_f, \text{Ad})}{\zeta(\frac{N}{2} + 1)\zeta(N)} \left(2^{1-\frac{N}{2}} \pi^2 \frac{\Gamma(\frac{N}{4} + \frac{\sqrt{-1}r}{2})\Gamma(\frac{N}{4} - \frac{\sqrt{-1}r}{2})}{\Gamma(\frac{N}{4} + \frac{1}{2})^2} \right) \|f_0\|^2. \quad (31)$$

The above theorem together with Proposition 3.3 gives us Theorem 1.1 from the introduction. We can now obtain the important corollary regarding injectivity of the Borchers theta lift.

Corollary. 4.5. *Let $f \in S(\text{SL}_2(\mathbb{Z}); -\frac{r^2+1}{4})$ and F_f be its Borchers lift defined in Section 2.2. Then the map $f \rightarrow F_f$ is injective.*

Proof. If f is a Hecke eigenform, then by Theorem 2.2, so is F_f . By Proposition 3.3, the Petersson norm of F_f is given by Theorem 4.4 above. Hence, $f \rightarrow F_f$ is an injective map when restricted to Hecke eigenforms. To conclude the same about non-Hecke eigenforms, we follow the exact argument as in the proof of Theorem 7.1 of [43]. Note that, for this we need formulas for Hecke eigenvalues of F_f in terms of those of f , and these are obtained in Theorem 4.11 of [37]. $\square$

5 Sup-norm bounds for Maass cusp forms on $O(1, N+1)$

In this section, we will obtain lower and upper bounds for the sup-norm of F_f . For the lower bounds we will use the Fourier expansion of F_f . For the upper bound, we will use a combination of Fourier expansion of F_f and the pre-trace formula method.

5.1 Upper and lower bounds using Fourier expansions

Recall from Section 2.2 that we have $f \in S(\text{SL}_2(\mathbb{Z}); -\frac{r^2+1}{4})$ given by the Fourier expansion

$$\begin{aligned} f(u + iv) &= \sum_{n \neq 0} c(n) W_{0, \frac{\sqrt{-1}r}{2}}(4\pi|n|v) \exp(2\pi\sqrt{-1}nu) \\ &= \sum_{n \neq 0} 2c(n) |n|^{1/2} v^{1/2} K_{\frac{\sqrt{-1}r}{2}}(2\pi|n|v) \exp(2\pi\sqrt{-1}nu). \end{aligned}$$

We have the Borchers theta lift F_f obtained in Section 2.2 given by the Fourier expansion

$$F_f(n(x)a_y) = \sum_{\lambda \in L \setminus \{0\}} A(\lambda) y^{\frac{N}{2}} K_{\sqrt{-1}r}(4\pi|\lambda|_S y) \exp(2\pi\sqrt{-1}^t \lambda S x) \quad (32)$$

with

$$A(\lambda) = |\lambda|_S \sum_{d|\lambda} c \left(-\frac{|\lambda|_S^2}{d^2} \right) d^{\frac{N}{2}-2}.$$

Let us make the assumption that f is a Hecke eigenform with $c(m) = \pm c(-m)$ for all $m \in \mathbb{Z}$. As in Remark 1, we also note that r is bounded from below, i.e. $r \gg 1$.

Fourier expansion of F_f at the cusps of Γ_S

In addition to the Fourier expansion (32) of F_f at the cusp at ∞ (a neighborhood of $y = \infty$ in H_N), we will also need the same at other cusps of Γ_S . Let us briefly recall this from Section 3.3 of [37]. Let the adelization of F_f (denoted by the same symbol) be as defined in equations (3.2)-(3.5) of [37].

There is a bijection between the cusps of Γ_S and the set $\mathcal{P}(\mathbb{Q}) \backslash \mathcal{G}(\mathbb{Q}) / \Gamma_S$, which sends $c \in \mathcal{G}(\mathbb{Q})$ to the cusp $c_\infty^{-1}\infty$. Moreover, we know from Lemma 2.1 that there is a bijection between $\mathcal{P}(\mathbb{Q}) \backslash \mathcal{G}(\mathbb{Q}) / \Gamma_S$ and the set of double cosets $\mathcal{H}(\mathbb{Q}) \backslash \mathcal{H}(\mathbb{A}) / U_f U_\infty$, which have representatives in $\mathcal{H}(\mathbb{A}_f)$. This bijection is defined so that if $c \in \mathcal{G}(\mathbb{Q})$ corresponds to $h \in \mathcal{H}(\mathbb{A}_f)$, then there exist $\gamma \in \mathcal{P}(\mathbb{Q})$ and $g \in G_\infty K_f$ such that

$$c = \gamma \begin{bmatrix} 1 & & \\ & h & \\ & & 1 \end{bmatrix} g. \quad (33)$$

Note that we will simply write the matrix $\begin{bmatrix} 1 & & \\ & h & \\ & & 1 \end{bmatrix}$ as h if the meaning is clear from the context.

Let $c \in \mathcal{G}(\mathbb{Q})$, and let γ , h , and g be as in (33). We write c_f (respectively c_∞) for the finite part (respectively the infinite part) of c , and likewise for γ . We then see $\gamma_f^{-1} c_f = h k_f$ with some $k_f \in K_f$. Using the left invariance of F_f by $\mathcal{G}(\mathbb{Q})$ and the right invariance by $K_f K_\infty$, we get for $x \in \mathbb{R}^N$, $y \in \mathbb{R}_{>0}$,

$$\begin{aligned} F_f(n(x) a_y h) &= F_f(n(x) a_y \gamma_f^{-1} c_f) = F_f(\gamma_f^{-1} c_f n(x) a_y) \\ &= F_f((\gamma_\infty^{-1} c_\infty)^{-1} n(x) a_y) = F_f(c_\infty^{-1} n(x') a_{y'}), \end{aligned} \quad (34)$$

where we have the Iwasawa decomposition $\gamma_\infty n(x) a_y = n(x') a_{y'} k'_\infty$ with some $(x', y') \in \mathbb{R}^N \times \mathbb{R}_{>0}$ and some $k'_\infty \in K_\infty$, the latter of which can be omitted in the above equations by the right K_∞ -invariance of F_f . We therefore see that $F_f(n(x) a_y h)$ is indeed an expression for F_f in Iwasawa coordinates with respect to the cusp $c_\infty^{-1}\infty$ corresponding to c .

The definition of the adelic $F_f(n(x) a_y h)$ given in (3.2), (3.3) of [37] implies

$$F_f(n(x) a_y h) = \sum_{\lambda \in L_h \setminus \{0\}} A_h(\lambda) y^{\frac{N}{2}} K_{\sqrt{-1}r} (4\pi |\lambda|_S y) \exp(2\pi \sqrt{-1}^t \lambda S x). \quad (35)$$

Here, L_h is the lattice in the genus of L associated to h , given by $L_h = \mathbb{Q}^N \cap (\mathbb{R}^N \times \prod_{p < \infty} h_p L_p)$, and

$$A_h(\lambda) = |\lambda|_S \sum_{d|d_\lambda} c \left(-\frac{|\lambda|_S^2}{d^2} \right) d^{\frac{N}{2}-2},$$

with d_λ being the smallest positive integer such that $\frac{1}{d_\lambda} \lambda \in L_h$. Note that L_h is also an even unimodular lattice. Moreover, by (34), equation (35) gives the Fourier expansion of F_f in the cusp $c_\infty^{-1}\infty$ corresponding to c .

Bounds on $\|F_f\|_2$, $|A_h(\lambda)|$ and $K_{\sqrt{-1}r}(y)$

Note that $c(1) \neq 0$ and we have the following estimate on the first Fourier coefficient $c(1)$ given by [29, Theorem 2] and [25, Corollary 0.3]. For any $\epsilon > 0$

$$r^{-\epsilon} \cosh(\pi r/2) \ll_\epsilon \frac{|c(1)|^2}{\|f\|_2^2} \ll_\epsilon r^\epsilon \cosh(\pi r/2). \quad (36)$$

Here $\|f\|_2$ is the Petersson norm of f , and we are not assuming it to be equal to 1.

Proposition. 5.1. *For any $\epsilon > 0$ we have an estimate as follows:*

i)

$$\sqrt{\sinh(\pi r/2)} r^{-\frac{N}{4} + \frac{1}{2}} \ll_N \frac{\|f\|_2}{\|F_f\|_2} \ll_N \sqrt{\sinh(\pi r/2)} r^{-\frac{N}{4} + \frac{1}{2}}.$$

ii) For all $\lambda \in L_h \setminus \{0\}$ and $\epsilon > 0$, we have

$$\begin{aligned} \frac{|A_h(\lambda)|}{\|f\|_2} &\ll_\epsilon |\lambda|_S^{2\theta+1+\epsilon} d_\lambda^{\frac{N}{2}-2-2\theta} r^\epsilon \sqrt{\cosh(\pi r/2)} \\ &\ll_\epsilon |\lambda|_S^{\frac{N}{2}-1+\epsilon} r^\epsilon \sqrt{\cosh(\pi r/2)}, \end{aligned}$$

where $\theta = 7/64$ is the current best estimate towards the Ramanujan conjecture for Maass forms.*Proof.* By Theorem 4.4, we have

$$\|F_f\|_2^2 = \frac{L(\frac{N}{2}, \pi_f, \text{Ad})}{\zeta(\frac{N}{2}+1)\zeta(N)} \left(2^{1-\frac{N}{2}} \pi^2 \frac{\Gamma(\frac{N}{4} + \frac{\sqrt{-1}r}{2}) \Gamma(\frac{N}{4} - \frac{\sqrt{-1}r}{2})}{\Gamma(\frac{N}{4} + \frac{1}{2})^2} \right) \|f\|_2^2.$$

We have

$$1 \ll_N \frac{2^{1-\frac{N}{2}} \pi^2 L(\frac{N}{2}, \pi_f, \text{Ad})}{\zeta(\frac{N}{2}+1)\zeta(N)\Gamma(\frac{N}{4} + \frac{1}{2})^2} \ll_N 1.$$

Since $N/2 > 1 + 2\theta$, the L -function above is given by a convergent Dirichlet series, and hence can be bounded by estimates independent of f .

For the terms involving the Gamma function we use the following standard properties

$$\Gamma\left(\frac{N}{4} + \frac{\sqrt{-1}r}{2}\right) \Gamma\left(\frac{N}{4} - \frac{\sqrt{-1}r}{2}\right) = |\Gamma\left(\frac{N}{4} + \frac{\sqrt{-1}r}{2}\right)|^2 = \frac{\pi r/2}{\sinh(\pi r/2)} \prod_{k=1}^{\frac{N}{4}-1} (k^2 + (r/2)^2).$$

Hence, we have

$$\frac{\pi r/2}{\sinh(\pi r/2)} r^{\frac{N}{2}-2} \ll_N \Gamma\left(\frac{N}{4} + \frac{\sqrt{-1}r}{2}\right) \Gamma\left(\frac{N}{4} - \frac{\sqrt{-1}r}{2}\right) \leq \frac{\pi r/2}{\sinh(\pi r/2)} (N^2 + r^2)^{\frac{N}{4}-1} \ll_N \frac{\pi r/2}{\sinh(\pi r/2)} r^{\frac{N}{2}-2}.$$

Here, we have used $r \gg 1$. Hence we obtain the estimate for $\|f\|_2/\|F_f\|_2$ in the statement of the proposition.To obtain the estimate for $|A_h(\lambda)|$, we have

$$|A_h(\lambda)| \leq |\lambda|_S \sum_{d|\lambda} |c\left(-\frac{|\lambda|_S^2}{d^2}\right)| d^{\frac{N}{2}-2} = |\lambda|_S \sum_{d|\lambda} |c(1)| \mu\left(\frac{|\lambda|_S^2}{d^2}\right) d^{\frac{N}{2}-2},$$

where $\mu(m)$ is the Hecke eigenvalue for f for the Hecke operator $T(m)$. We have, by [32, Appendix 2, Proposition 2] (see also [30, Section 8.5]), $|\mu(m)| \leq m^\theta \tau(m)$ where $\tau(m)$ is the number of divisors of m and $\theta = 7/64$ is the current best estimate towards the Ramanujan conjecture for Maass forms. Using $\tau(m) \ll_\epsilon m^\epsilon$ for every $m > 0$, the right hand inequality of (36) and $\frac{N}{2} - 2 - 2\theta - \epsilon > 0$, we have

$$\begin{aligned} |A_h(\lambda)| &\leq |\lambda|_S \sum_{d|\lambda} |c(1)| \tau\left(\frac{|\lambda|_S^2}{d^2}\right) \left(\frac{|\lambda|_S^2}{d^2}\right)^\theta d^{\frac{N}{2}-2} \\ &\ll_\epsilon |\lambda|_S^{2\theta+1+\epsilon} |c(1)| \sum_{d|\lambda} d^{\frac{N}{2}-2-2\theta-\epsilon} \\ &\ll_\epsilon |\lambda|_S^{2\theta+1+\epsilon} |c(1)| d_\lambda^{\frac{N}{2}-2-2\theta} \end{aligned}$$

$$\begin{aligned} &\ll_{\epsilon} |\lambda|_S^{2\theta+1+\epsilon} d_{\lambda}^{\frac{N}{2}-2-2\theta} \|f\|_2 r^{\epsilon} \sqrt{\cosh(\pi r/2)} \\ &\ll_{\epsilon} |\lambda|_S^{\frac{N}{2}-1+\epsilon} \|f\|_2 r^{\epsilon} \sqrt{\cosh(\pi r/2)}, \end{aligned}$$

which completes the proof of the proposition. $\square$

We will require the estimate for the K -Bessel function.

Lemma. 5.2. *Let $r \gg 1$ be a real number.*

i) *If $1 \ll y < r$, we have*

$$e^{\pi r/2} K_{\sqrt{-1}r}(y) \ll r^{-1/4} (r-y)^{-1/4}.$$

ii) *If $y = r + O(r^{1/3})$ and $y \gg 1$, we have*

$$e^{\pi r/2} K_{\sqrt{-1}r}(y) = \pi(2/y)^{1/3} \text{Ai}(\xi e^{-2\pi i/3}) + O(y^{-2/3}) \ll r^{-1/3},$$

where $\xi = i(y-r)(-iy/2)^{-1/3}$.

iii) *If $2r > y > r$, we have*

$$e^{\pi r/2} K_{\sqrt{-1}r}(y) \ll r^{-1/4} (y-r)^{-1/4} \exp(-Cr^{-1/2}(y-r)^{3/2})$$

for some $C > 0$.

iv) *If $y \geq 2r$, we have*

$$e^{\pi r/2} K_{\sqrt{-1}r}(y) \ll \exp(-Cy)$$

for some $C > 0$.

Proof. The bound (i) follows from the asymptotic of Erdélyi [19, 7.13.2, (19)] (after noting that $r+y \sim r$ in the range under consideration), and formula (ii) is due to Balogh [2, (8)]. We shall derive (iii) and (iv) from the formula [19, 7.13.2, (18)] of Erdélyi, which gives

$$e^{\pi r/2} K_{\sqrt{-1}r}(y) \ll (y^2 - r^2)^{-1/4} \exp(\pi r/2 - (y^2 - r^2)^{1/2} - r \sin^{-1}(r/y)).$$

To derive (iii) from this, we must show that the argument of the exponential satisfies

$$-\pi r/2 + (y^2 - r^2)^{1/2} + r \sin^{-1}(r/y) \gg r^{-1/2} (y-r)^{3/2} \quad (37)$$

when $2r > y > r$. To do this, define $\rho = y/r$. If we define $f(\rho) = -\pi/2 + (\rho^2 - 1)^{1/2} + \sin^{-1}(\rho^{-1})$, then we have

$$-\pi r/2 + (y^2 - r^2)^{1/2} + r \sin^{-1}(r/y) = r f(\rho).$$

We have $f'(\rho) = \sqrt{\rho^2 - 1}/\rho \sim \sqrt{\rho - 1}$ when $2 > \rho > 1$, and $f(1) = 0$. This implies that $f(\rho) \sim (\rho - 1)^{3/2}$ in this range, which gives (37).

To establish (iv), we note that when $\rho \geq 2$ we have $f'(\rho) = \sqrt{1 - \rho^{-2}} \geq 1/2$, which implies that $f(\rho) \geq f(2) + (\rho - 2)/2$ in this range. Because $f(2) > 0$, this implies that $f(\rho) \gg \rho$, and hence that

$$-\pi r/2 + (y^2 - r^2)^{1/2} + r \sin^{-1}(r/y) = r f(\rho) \gg y.$$

Hence, we get

$$e^{\pi r/2} K_{ir}(y) \ll (y^2 - r^2)^{-1/4} e^{-Cy} \ll y^{-1/2} e^{-Cy} \ll e^{-Cy},$$

since $y \geq 2r$ and $r \gg 1$. $\square$

Finally, we need a lemma to estimate the number of vectors in the lattice L_h with prescribed norms.

Lemma. 5.3. *Let L_h be the even unimodular lattices as above with dimension N . Set $k = N/2 - 1$. Then, for $m \in \mathbb{Z}_{>0}$, $k' \in \mathbb{R}_{>0}$ and $\epsilon > 0$, we have*

$$\sum_{\substack{\lambda \in L_h \\ |\lambda|_S^2 = m}} d_\lambda^{k-k'} \ll_{L, \epsilon} m^{k+\epsilon}, \quad (38)$$

where we note that the implied constant is independent of h .

Proof. We have

$$\begin{aligned} \sum_{\substack{\lambda \in L_h \\ |\lambda|_S^2 = m}} d_\lambda^{k-k'} &= \sum_{d^2|m} \sum_{\substack{|\lambda|_S^2 = m \\ d_\lambda = d}} d^{k-k'} = \sum_{d^2|m} d^{k-k'} \sum_{\substack{|\lambda|_S^2 = m \\ d_\lambda = d}} 1 \\ &= \sum_{d^2|m} d^{k-k'} \sum_{\substack{|\lambda|_S^2 = m/d^2 \\ d_\lambda = 1}} 1 \leq \sum_{d^2|m} d^{k-k'} \sum_{|\lambda|_S^2 = m/d^2} 1. \end{aligned}$$

Since L_h is assumed to be an even unimodular lattice of dimension N , [53, p.109, Corollary 2] and [16, Proposition 1.3.5] implies that, for any $M > 0$, we have $\#\{\lambda \in L_h \mid |\lambda|_S^2 = M\} \ll_L \tau(M) M^{\frac{N}{2}-1}$. Note that we have $\tau(M) \ll_\epsilon M^\epsilon$. Using this we get

$$\begin{aligned} \sum_{\substack{\lambda \in L_h \\ |\lambda|_S^2 = m}} d_\lambda^{k-k'} &\ll_{L, \epsilon} \sum_{d^2|m} d^{k-k'} \left(\frac{m}{d^2}\right)^{k+\epsilon} = \sum_{d^2|m} \frac{m^{k+\epsilon}}{d^{k+k'+2\epsilon}} \\ &\ll_{L, \epsilon} \sum_{d^2|m} m^{k+\epsilon} \leq m^{k+\epsilon} \tau(m) \ll_{L, \epsilon} m^{k+\epsilon}, \end{aligned}$$

as required. $\square$

Upper bound for sup-norm using Fourier expansion of F_f

In this section we will obtain an upper bound for $|F_f(n(x)a_y h)|/||F_f||_2$ using the Fourier expansion (35) of F_f and the bounds on $||F_f||_2$, $|A_h(\lambda)|$, $K_{\sqrt{-1}r}(y)$ and bounds on lattice points obtained in Proposition 5.1 and Lemmas 5.2 and 5.3.

Theorem. 5.4. *Let $f \in S(\mathrm{SL}_2(\mathbb{Z}); -\frac{r^2+1}{4})$ be a non-zero Hecke eigenform with Fourier coefficients $c(-m) = \pm c(m)$ for all $m \in \mathbb{Z}$, and recall that $r \gg 1$. Let $F_f \in \mathcal{M}(\Gamma_S, \sqrt{-1}r)$ be the Borchers theta lift of f . For any $\epsilon > 0$, $x \in \mathbb{R}^N$, and any $h \in \mathcal{H}(\mathbb{A}_f)$ corresponding to a cusp of Γ_S , the following holds:*

$$\frac{1}{||F_f||_2} |F_f(n(x)a_y h)| \ll_{\epsilon, N, L} \begin{cases} y^{-N/2-1-2\theta} r^{3N/4+1+2\theta+\epsilon} & 1 \ll y \leq r^{11/12}; \\ y^{-N/2+1-2\theta} r^{3N/4-5/6+2\theta+\epsilon} & r^{11/12} < y \leq r/2\pi; \\ e^{-Cy} & r/2\pi < y. \end{cases}$$

Proof. Using the Fourier expansion of F_f given by (35), we see that it suffices to bound

$$\frac{1}{||F_f||_2} \sum_{\lambda \in L_h} |A_h(\lambda)| y^{N/2} |K_{\sqrt{-1}r}(4\pi|\lambda|_S y)|.$$

Using Proposition 5.1 and Lemma 5.3, this is bounded by

$$S_y := r^{-N/4+1/2+\epsilon} y^{N/2} \sum_{m \geq 1} m^{N/2-1/2+\theta+\epsilon} e^{\pi r/2} |K_{\sqrt{-1}r}(4\pi\sqrt{m}y)|. \quad (39)$$

We break the sum in (39) into four ranges and denote $S_y := S_y^{(1)} + S_y^{(2)} + S_y^{(3)} + S_y^{(4)}$, where $S_y^{(1)}$ is obtained by summing over $4\pi\sqrt{my} \leq r/2$, i.e. $m \leq (r/8\pi y)^2$, $S_y^{(2)}$ is obtained by summing over $r/2 < 4\pi\sqrt{my} \leq r$, i.e. $(r/8\pi y)^2 < m \leq (r/4\pi y)^2$, $S_y^{(3)}$ is obtained by summing over $r < 4\pi\sqrt{my} \leq 2r$ and $S_y^{(4)}$ is obtained by summing over $2r < 4\pi\sqrt{my}$.

Computing $S_y^{(1)}$: In this range, the bound (i) from Lemma 5.2 becomes $e^{\pi r/2} K_{\sqrt{-1}r}(4\pi\sqrt{my}) \ll r^{-1/2}$, so we have

$$\begin{aligned} S_y^{(1)} &\ll r^{-N/4+1/2+\epsilon} y^{N/2} r^{-1/2} \sum_{1 \leq m \leq (r/8\pi y)^2} m^{N/2-1/2+\theta+\epsilon} \\ &\ll r^{-N/4+\epsilon} y^{N/2} (r/y)^{N+1+2\theta+\epsilon} \ll \frac{r^{\frac{3N}{4}+1+2\theta+\epsilon}}{y^{\frac{N}{2}+1+2\theta}}. \end{aligned}$$

Computing $S_y^{(2)}$: In this range we have $m \sim (r/y)^2$, so that

$$S_y^{(2)} \ll r^{-N/4+1/2+\epsilon} y^{N/2} (r/y)^{N-1+2\theta+\epsilon} \sum_{(r/8\pi y)^2 < m \leq (r/4\pi y)^2} e^{\pi r/2} |K_{\sqrt{-1}r}(4\pi\sqrt{my})|. \quad (40)$$

If the sum on the right hand side of (40) is nonempty, its highest term is $m_* = \lfloor (r/4\pi y)^2 \rfloor$. Lemma 5.2 implies that $e^{\pi r/2} K_{\sqrt{-1}r}(4\pi\sqrt{m_*y}) \ll r^{-1/3}$ for all y , and so applying this to bound the contribution of m_* to the sum gives

$$\sum_{(r/8\pi y)^2 < m \leq (r/4\pi y)^2} e^{\pi r/2} |K_{\sqrt{-1}r}(4\pi\sqrt{my})| \ll r^{-1/3} + \sum_{(r/8\pi y)^2 < m \leq m_*-1} e^{\pi r/2} |K_{\sqrt{-1}r}(4\pi\sqrt{my})|.$$

For the remaining terms in the sum, we apply Lemma 5.2 (i), which gives

$$\sum_{(r/8\pi y)^2 < m \leq m_*-1} e^{\pi r/2} |K_{\sqrt{-1}r}(4\pi\sqrt{my})| \ll r^{-1/4} \sum_{(r/8\pi y)^2 < m \leq m_*-1} (r - 4\pi\sqrt{my})^{-1/4}.$$

Because the function $x \rightarrow (r - 4\pi\sqrt{xy})^{-1/4}$ is increasing on the interval $((r/8\pi y)^2, m_*)$, we may bound the sum by an integral as follows:

$$\begin{aligned} r^{-1/4} \sum_{(r/8\pi y)^2 < m \leq m_*-1} (r - 4\pi\sqrt{my})^{-1/4} &< r^{-1/4} \int_{(r/8\pi y)^2}^{m_*} (r - 4\pi\sqrt{xy})^{-1/4} dx \\ &\leq r^{-1/4} \int_{(r/8\pi y)^2}^{(r/4\pi y)^2} (r - 4\pi\sqrt{xy})^{-1/4} dx. \end{aligned}$$

We then have

$$\begin{aligned} r^{-1/4} \int_{(r/8\pi y)^2}^{(r/4\pi y)^2} (r - 4\pi\sqrt{xy})^{-1/4} dx &= r^{-1/4} \int_{r/8\pi y}^{r/4\pi y} (r - 4\pi uy)^{-1/4} 2u du \\ &\leq r^{3/4} / 2\pi y \int_{r/8\pi y}^{r/4\pi y} (r - 4\pi uy)^{-1/4} du \\ &= r^{3/4} / 8\pi^2 y^2 \int_{r/2}^r (r - u)^{-1/4} du \\ &\ll r^{3/2} y^{-2}. \end{aligned}$$

Combining these gives

$$S_y^{(2)} \ll \frac{r^{\frac{3N}{4}-\frac{1}{2}+2\theta+\epsilon}}{y^{\frac{N}{2}-1+2\theta}} (r^{-1/3} + r^{3/2} y^{-2}).$$

We note that this term dominates the contribution from $S_y^{(1)}$.

Computing $S_y^{(3)}$: Following the computation of $S_y^{(2)}$, we can see that we get the same bound for $S_y^{(3)}$ as we got for $S_y^{(2)}$ above. To see this, note that we again have $m \sim (r/y)^2$, so it suffices to bound the sum

$$\sum_{(r/4\pi y)^2 < m \leq (r/2\pi y)^2} e^{\pi r/2} |K_{\sqrt{-1}r}(4\pi\sqrt{my})|.$$

As in the case of $S_y^{(2)}$, we may bound the extremal term in this sum by $r^{-1/3}$. For the other terms, we may use the bound $e^{\pi r/2} K_{\sqrt{-1}r}(4\pi\sqrt{my}) \ll r^{-1/4}(y-r)^{-1/4}$ coming from Lemma 5.2 (iii), which gives a contribution of $\ll r^{3/2}y^{-2}$ as in the case of $S_y^{(2)}$.

Computing $S_y^{(4)}$: Here, we use part (iv) of Lemma 5.2, which gives

$$S_y^{(4)} \ll r^{-N/4+1/2+\epsilon} y^{N/2} \sum_{(r/2\pi y)^2 < m} m^{N/2-1/2+\theta+\epsilon} \exp(-C\sqrt{my}).$$

Because $y \gg 1$ we have $\sqrt{my} \gg \sqrt{m} + y$, so

$$\begin{aligned} S_y^{(4)} &\ll r^{-N/4+1/2+\epsilon} y^{N/2} e^{-Cy} \sum_{(r/2\pi y)^2 < m} m^{N/2-1/2+\theta+\epsilon} \exp(-C\sqrt{m}) \\ &\ll r^{-N/4+1/2+\epsilon} y^{N/2} e^{-Cy} \sum_{m \geq 1} m^{N/2-1/2+\theta+\epsilon} \exp(-C\sqrt{m}) \\ &\ll r^{-N/4+1/2+\epsilon} e^{-Cy} \ll e^{-Cy}, \end{aligned}$$

where the final inequality follows because $y \gg r$.

Finally, we combine these estimates to obtain the proposition. First note that, when $y > r/2\pi$, the sums $S_y^{(1)}$, $S_y^{(2)}$ and $S_y^{(3)}$ are empty, and so we get

$$\frac{|F_f(n(x)a_y h)|}{\|F_f\|_2} \ll S_y = S_y^{(4)} \ll e^{-Cy},$$

as required. When $y \leq r/2\pi$, we can check that the contribution from $S_y^{(2)}$ dominates the one from $S_y^{(4)}$, and hence we get

$$\frac{|F_f(n(x)a_y h)|}{\|F_f\|_2} \ll S_y \ll \frac{r^{\frac{3N}{4}-\frac{1}{2}+2\theta+\epsilon}}{y^{\frac{N}{2}-1+2\theta}} (r^{-1/3} + r^{3/2}y^{-2}).$$

The term $y^{-2}r^{3/2}$ dominates when $y \leq r^{11/12}$, while $r^{-1/3}$ dominates when $y > r^{11/12}$, which gives the first two bounds of the theorem. $\square$

Lower bound for the sup-norm

We now use the Fourier expansion to obtain a lower bound for the sup norm of F_f . When doing this, it will be convenient to work in a cusp $c_\infty^{-1}\infty$ corresponding to $c \in \mathcal{G}(\mathbb{Q})$ such that the corresponding unimodular lattice L_h has a vector of length one. As all even unimodular lattices form a single genus, we may do this by taking L_h to be a sum of copies of the E_8 lattice.

Recall again that we have the Fourier expansion

$$F_f(n(x)a_y h) = \sum_{\lambda \in L_h \setminus \{0\}} A_h(\lambda) y^{\frac{N}{2}} K_{\sqrt{-1}r}(4\pi|\lambda|_S y) \exp(2\pi\sqrt{-1}^t \lambda S x).$$

Fix $\lambda_0 \in L_h$ to be an element of norm 1. We then have $A_h(\lambda_0) = c(-1) = \pm c(1) \neq 0$. Using the above Fourier expansion of F_f , the Cauchy-Schwartz inequality, and the fact that L is a unimodular lattice, we

get

$$\begin{aligned}
\|F_f\|_\infty &= \text{vol}(\mathbb{R}^N/L)^{-1} \int_{\mathbb{R}^N/L} \|F_f\|_\infty dx \\
&\geq \int_{\mathbb{R}^N/L} |F_f(n(x)a_y h)| \exp(-2\pi\sqrt{-1}^t \lambda_0 Sx) dx \\
&\geq \left| \int_{\mathbb{R}^N/L} F_f(n(x)a_y h) \exp(-2\pi\sqrt{-1}^t \lambda_0 Sx) dx \right| \\
&= |A_h(\lambda_0) y^{N/2} K_{\sqrt{-1}r}(4\pi y)|.
\end{aligned}$$

By part (ii) of Lemma 5.2, there is a value of y such that $4\pi y = r + O(r^{1/3})$, and $|K_{\sqrt{-1}r}(4\pi y)| \gg e^{-\pi r/2} r^{-1/3}$. Choosing this value of y and using $|A_h(\lambda_0)| = |c(1)|$ gives

$$\|F_f\|_\infty \gg |c(1)| r^{N/2-1/3} e^{-\pi r/2}. \quad (41)$$

The lower bound of Theorem 1.2 now follows from (41) using the bound (36), and part i) of Proposition 5.1.

5.2 Upper bounds using the pre-trace formula

In this section, we use a pre-trace inequality to obtain upper bounds on the lifted form F_f . The bound we prove holds for any square-integrable Laplace eigenfunction on any hyperbolic orbifold X of finite volume with the Laplacian Δ . To state it, we will need the notion of the height of a point $x \in X$ in the cusp, denoted by $\text{ht}(x)$, which we recall in this section. We note that only for this section, $\sqrt{-1}r$ will denote the spectral parameter of an eigenfunction on X , instead of the parameter of the Maass form f . We note that these two spectral parameters are the same size, so this should not lead to any confusion.

Theorem. 5.5. *Let X be a finite-volume hyperbolic orbifold of dimension $N+1$. Let $\psi \in L^2(X)$ be an L^2 -normalized Laplace eigenfunction with spectral parameter $\sqrt{-1}r$, so that $(\Delta + r^2 + N^2/4)\psi = 0$. We have*

$$\psi(x) \ll (1 + |r|)^{N/2} + \text{ht}(x)^{N/2} (1 + |r|)^{N/4}, \quad x \in X.$$

We shall assume in the proof that $r > 1$, as the other case is similar.

Background on hyperbolic geometry

We let G^0 be the group of isometries of H_N , which can be identified with the index two subgroup of $O(1, N)$ preserving the upper sheet of the two-sheeted hyperboloid. Let d be the standard distance function on H_N , and let $\partial H_N \simeq S^N$ denote the boundary sphere.

Let $X = \Gamma \backslash H_N$, where $\Gamma < G^0$ is a lattice. Let $O(\Gamma) \subset \partial H_N$ denote the set of fixed points of parabolic elements of Γ . We fix a set of representatives Ξ for the Γ -orbits in $O(\Gamma)$, which can be identified with the set of cusps of X . We let Γ_ξ be the stabilizer of $\xi \in \Xi$ in Γ . For each $\xi \in \Xi$ we choose a horoball B_ξ tangent to the boundary at ξ , and denote $V_\xi = \Gamma_\xi \backslash B_\xi$. By [47, Thm 12.7.4], we may choose each B_ξ so that the following hold:

- i) V_ξ is mapped isometrically to its image in X .
- ii) We have $X = X_0 \coprod V_\xi$, where X_0 is compact.
- iii) If $\gamma \in \Gamma - \Gamma_\xi$, then $B_\xi \cap \gamma B_\xi = \emptyset$.

We note that [47] states this theorem for manifolds, rather than orbifolds, but one may easily obtain the result for an orbifold by using Selberg's lemma to pass to a finite-index torsion-free subgroup of Γ . We now define the height function $\text{ht} : X \rightarrow \mathbb{R}_{\geq 1}$. If $x \in X_0$, we set $\text{ht}(x) = 1$. Next, suppose that $x \in V_\xi$ for some ξ . Let B_∞ denote the standard horoball $\{(x, y) : x \in \mathbb{R}^N, y > 1\}$, and choose $g_\xi \in G^0$ such that $g_\xi B_\xi = B_\infty$. We define $\text{ht}(x) = y(g_\xi x)$ (the y -coordinate of $g_\xi x$), which is independent of the choice of g_ξ . This has the key property that for any $C > 0$, the set of points $x \in X$ with $\text{ht}(x) \leq C$ is compact.

Test functions

In this section, we construct a test function for use in the pre-trace inequality. We shall do this using the Harish-Chandra transform, which we now recall. Define φ_s to be the standard spherical function on H_N or G^0 with spectral parameter s . We continue to normalize s so that $\sqrt{-1}\mathbb{R}$ is the tempered axis. Let $K \simeq O(N)$ be the standard maximal compact subgroup of G^0 . For a K -biinvariant function $k \in C_c^\infty(G^0)$, we define its Harish-Chandra transform by

$$\widehat{k}(s) = \int_{G^0} k(g) \varphi_s(g) dg.$$

This is inverted by

$$k(g) = C_N \int_{\sqrt{-1}\mathbb{R}} \widehat{k}(s) \varphi_s(g) |c(s)|^{-2} ds$$

for a constant C_N , where $c(s)$ is Harish-Chandra's c -function. We may now define the test function we shall use.

Lemma. 5.6. *There exists a K -biinvariant function $k_r \in C_c^\infty(G^0)$ with the following properties:*

- (i) k_r is supported in a fixed compact set that is independent of r .
- (ii) $\widehat{k}_r(s) \geq 0$ for $s \in \sqrt{-1}\mathbb{R} \cup (0, N/2]$.
- (iii) $\widehat{k}_r(\sqrt{-1}r) = 1$.
- (iv) $k_r(g) \ll r^N (1 + rd(g, e))^{-N/2}$. In particular, $\|k_r\|_\infty \ll r^N$.

Proof. Let $h \in C^\infty(\mathbb{C})$ be a function of Paley-Wiener type (i.e. the Fourier transform of a function in $C_c^\infty(\mathbb{R})$) that is real and non-negative on $\mathbb{R}$, and satisfies $h(0) = 1$. Define h_r by $h_r(s) = h(r-s) + h(r+s)$. h_r is real valued on $\mathbb{R}$, and satisfies $h_r(r) \geq 1$. We also have $h_r(\bar{s}) = \overline{h_r(s)}$ and $h_r(s) = h_r(-s)$ so h_r is real-valued on $\sqrt{-1}\mathbb{R}$.

Define k_r to be the K -biinvariant function on G^0 satisfying $\widehat{k}_r(\sqrt{-1}s) = h_r^2(s)$. Because h_r is of Paley-Wiener type, (i) follows from the Paley-Wiener theorem of Gangolli-Varadarajan (cf. [21, Theorem 6.6.8]). For $s \in \sqrt{-1}\mathbb{R} \cup (0, N/2]$, we have that $h_r(s)$ is real, and hence $\widehat{k}_r(s) = h_r^2(s) \geq 0$. We also have $\widehat{k}_r(\sqrt{-1}r) = h_r^2(r) \geq 1$, and so we may arrange that $\widehat{k}_r(\sqrt{-1}r) = 1$ by scaling h_r .

Finally, (iv) follows from inverting the Harish-Chandra transform and applying bounds for the spherical function. We have

$$k_r(g) = \int_{\sqrt{-1}\mathbb{R}} \widehat{k}_r(s) \varphi_s(g) |c(s)|^{-2} ds.$$

We have $|c(s)|^{-2} \ll (1 + |s|)^N$ for $s \in \sqrt{-1}\mathbb{R}$. By [39, Theorem 1.3], or by applying [18] together with stationary phase, we have

$$\varphi_s(g) \ll (1 + |s|d(g, e))^{-N/2}$$

for all $s \in \sqrt{-1}\mathbb{R}$ and g in the support of k_r , which implies

$$k_r(g) \ll \int_{\sqrt{-1}\mathbb{R}} |\widehat{k}_r(s)| (1 + |s|d(g, e))^{-N/2} (1 + |s|)^N ds.$$

The rapid decay of $\widehat{k}_r(s)$ away from $s = \pm\sqrt{-1}r$ effectively truncates the integral to the region where $s = \pm\sqrt{-1}(r + o(r))$, which gives (iv). $\square$

The pre-trace inequality

Let k_r be as in Lemma 5.6. The fundamental inequality we shall use to bound ψ is

$$|\psi(x)|^2 \leq \sum_{\gamma \in \Gamma} k_r(x^{-1}\gamma x). \quad (42)$$

This may be derived from the pre-trace formula, by using the positivity property (ii) of $\widehat{k}_r$ to drop all terms on the spectral side other than $|\psi(x)|^2$ (including the continuous spectrum). For this we remark that the parameters of the discrete spectrum are contained in $\sqrt{-1}\mathbb{R} \cup (0, N/2]$, which parametrize equivalence classes of irreducible unitary spherical principal series representations of $O(1, N)$ together with spherical complimentary series representations. To confirm this fact on the representation theory of $O(1, N)$ we refer to [24] and [27, pp31–32, Remarks] for instance.

It may also be proved in an elementary way by an application of Cauchy–Schwartz and unfolding, see for instance [14, Lemma 6.5], with the test function ω there taken to be the function k_r^0 satisfying $\widehat{k}_r^0 = h_r$ so that $k_r^0 * (k_r^0)^* = k_r$.

Let $R > 0$ be a constant, independent of r , such that the support of k_r is contained in the open ball of radius R about the origin $(0, 1)$ in H_N . For x in any fixed compact subset of X , (42) and (iv) give $|\psi(x)| \ll r^{N/2}$, so we may assume that $\text{ht}(x) > e^R > 1$. In particular, x is contained in a cusp neighborhood V_ξ , which we shall assume to be fixed for the rest of the proof. Moreover, until further notice we identify x with a choice of lift $x \in B_\xi$. Under this assumption, we shall show that only $\gamma \in \Gamma_\xi$ contribute to (42). Indeed, suppose that $\gamma \in \Gamma$ satisfies $k_r(x^{-1}\gamma x) \neq 0$. This implies that $d(\gamma x, x) < R$. This, together with $\text{ht}(x) > e^R$, implies that $\gamma x \in B_\xi$, so that $\gamma B_\xi \cap B_\xi \neq \emptyset$, and hence that $\gamma \in \Gamma_\xi$ as required.

We now assume that our cusp ξ is the standard point at infinity, and denote Γ_ξ and B_ξ by Γ_∞ and B_∞ , as the proof for the other cusps is similar. We therefore have

$$|\psi(x)|^2 \leq \sum_{\gamma \in \Gamma_\infty} k_r(x^{-1}\gamma x).$$

We next apply our upper bound (iv) for k_r , which gives

$$|\psi(x)|^2 \ll r^N \sum_{\substack{\gamma \in \Gamma_\infty \\ d(\gamma x, x) < R}} (1 + rd(\gamma x, x))^{-N/2}.$$

We may identify Γ_∞ with a lattice in $\text{Isom}(\mathbb{R}^N)$. By [47, Thm 7.5.2], Γ_∞ has a finite index subgroup of translations of rank N , which we identify with a lattice $L \subset \mathbb{R}^N$. The action of L on H_N will be written additively. We let $\gamma_1, \dots, \gamma_k$ be coset representatives for $L \backslash \Gamma_\infty$, and define $x_i = \gamma_i x$. We therefore have

$$|\psi(x)|^2 \ll r^N \sum_{i=1}^k \sum_{\substack{\ell \in L \\ d(\ell + x_i, x) < R}} (1 + rd(\ell + x_i, x))^{-N/2}.$$

Theorem 5.5 now follows from the following lemma.

Lemma. 5.7. *Let $v_1, v_2 \in \mathbb{R}^N$, and let $x_i = (v_i, y) \in H_N$ with $y > 1$. We have*

$$\sum_{\substack{\ell \in L \\ d(\ell + x_1, x_2) < R}} (1 + rd(\ell + x_1, x_2))^{-N/2} \ll 1 + y^N r^{-N/2}.$$

Proof. We may assume without loss of generality that v_1, v_2 lie in a fixed fundamental domain for $\mathbb{R}^N/L$. It follows from an elementary computation that there is $C_1 > 0$, depending on R , such that $d(\ell + x_1, x_2) < R$ implies $\|\ell\| < C_1 y$. It may also be seen that $d(\ell + x_1, x_2) < R$ implies

$$d(\ell + x_1, x_2) \sim_R \|\ell + v_1 - v_2\|/y = \|\ell\|/y + O(1/y).$$

In particular, there is $C_2 > 0$ such that when $d(\ell + x_1, x_2) < R$ and $\|\ell\| > C_2$ we have $d(\ell + x_1, x_2) \sim_R \|\ell\|/y$. The finitely many ℓ with $\|\ell\| \leq C_2$ contribute $O(1)$ to the sum, so it suffices to show that

$$\sum_{\substack{\ell \in L \\ C_2 < \|\ell\| < C_1 y}} (1 + rd(\ell + x_1, x_2))^{-N/2} \sim_R \sum_{\substack{\ell \in L \\ C_2 < \|\ell\| < C_1 y}} (1 + r\|\ell\|/y)^{-N/2} \ll 1 + y^N r^{-N/2}. \quad (43)$$

We break the sum

$$\sum_{\substack{\ell \in L \\ C_2 < \|\ell\| < C_1 y}} (1 + r\|\ell\|/y)^{-N/2}$$

into those ℓ with $\|\ell\| < y/r$, and the complement. If $\|\ell\| < y/r$, we have $1 + r\|\ell\|/y \sim 1$, so the contribution these terms make is asymptotically bounded by $\#\{\ell \in L : \|\ell\| < y/r\} \ll 1 + y^N r^{-N}$ which is smaller than the right hand side of (43). If $\|\ell\| \geq y/r$, we have $1 + r\|\ell\|/y \sim r\|\ell\|/y$, so the contribution from these terms is bounded by

$$\sum_{\substack{\ell \in L \\ C_2 < \|\ell\| < C_1 y}} (r\|\ell\|/y)^{-N/2} = y^{N/2} r^{-N/2} \sum_{\substack{\ell \in L \\ C_2 < \|\ell\| < C_1 y}} \|\ell\|^{-N/2} \ll y^N r^{-N/2}$$

which is again less than the right hand side of (43). This completes the proof. $\square$

5.3 Proof of Theorem 1.2

In this section we will give the proof of Theorem 1.2 from the introduction. First note that Theorem 5.5 gives

$$|F(n_x a_y h)|/\|F_f\|_2 \ll r^{N/2} + y^{N/2} r^{N/4}$$

for all $h \in \mathcal{H}(\mathbb{A}_f)$ and all $y \gg 1$, where we are using (34) to convert between the classical and adelic pictures. By Theorem 5.4, it suffices to consider the range where $1 \ll y < r/2\pi$. We consider the point $y_0 = r^{(N/2+1+2\theta)/(N+1+2\theta)}$, which is chosen so that the expressions $y^{N/2} r^{N/4}$ and $y^{-N/2-1-2\theta} r^{3N/4+1+2\theta}$ appearing in Theorem 5.5 and Theorem 5.4 are both equal to $r^{N/2+N(1+2\theta)/4(N+1+2\theta)}$ (our desired upper bound) when evaluated at y_0 . An elementary computation gives us that $y_0 < r^{11/12}$.

The expression $r^{N/2} + y^{N/2} r^{N/4}$ is increasing in y , so for $1 \ll y \leq y_0$ we have

$$\begin{aligned} |F(n_x a_y h)|/\|F_f\|_2 &\ll r^{N/2} + y^{N/2} r^{N/4} \leq r^{N/2} + y_0^{N/2} r^{N/4} \\ &\ll r^{N/2+N(1+2\theta)/4(N+1+2\theta)}. \end{aligned}$$

We next suppose that $y_0 \leq y < r/2\pi$. Because the upper bound given by Theorem 5.4 is decreasing in y (including across the transition point at $y = r^{11/12}$), we may obtain an upper bound for $|F(n_x a_y h)|/\|F_f\|_2$ by evaluating the upper bound of Theorem 5.4 at y_0 . As $y_0 < r^{11/12}$, this gives

$$|F(n_x a_y h)|/\|F_f\|_2 \ll y_0^{-N/2-1-2\theta} r^{3N/4+1+2\theta+\epsilon} = r^{N/2+N(1+2\theta)/4(N+1+2\theta)+\epsilon}.$$

Note that we have $\Lambda \sim r^2$ since $r \gg 1$. Combining the above computations completes the proof of the upper bound in Theorem 1.2.

References

- [1] AVACUMOVIĆ, G. V.: *Über die Eigenfunktionen auf geschlossenen Riemannschen Mannigfaltigkeiten*, Math. Z. **65** (1956), 327–344.
- [2] BALOGH, C.: *Asymptotic expansions of the modified Bessel function of the third kind of imaginary order*, SIAM Journal on Applied Mathematics, Vol. **15**, No. 5 (Sep., 1967), 1315–1323.
- [3] BÉRARD, P. H.: *On the wave equation on a compact Riemannian manifold without conjugate points*, Math. Z. **155** no. 3 (1977), 249–276.
- [4] BORCHERDS, R. E.: *Automorphic forms with singularities on Grassmanians*. Invent. Math., **132** (1998), 491–562.
- [5] BLOMER, V.: *On the size of Ikeda lifts*. Manuscripta math., **148** (2015), 341–349.
- [6] BLOMER, V., HARCOS, G. AND MAGA, P.: *Analytic properties of spherical cusp forms on $GL(n)$* . J. d’Analyse Mathématique **140** (2020), 483–510.
- [7] BLOMER, V., HARCOS, G. AND MAGA, P.: *On the global sup-norm of $GL(3)$ cusp forms*. Israel J. Math. **229** (2019), 357–379.
- [8] BLOMER, V., HARCOS, G., MAGA, P. AND MILIĆEVIĆ, D.: *The sup-norm problem for $GL(2)$ over number fields*. J. Eur. Math. Soc. **22** (2020), 1–53.
- [9] BLOMER, V. AND HOLOWINSKI R.: *Bounding sup-norms of cusp forms of large level*. Invent. Math., **179** (2010), 645–681.
- [10] BLOMER, V. AND POHL, A.: *The sup-norm problem on the Siegel modular space of degree two*. Amer. J. Math., **138** (2016), 999–1027.
- [11] BOOKER, A., STRÖMBERGSSON, A., AND VENKATESH, A.: *Effective computation of Maass cusp forms*. Int. Math. Res. Not., **2006** (2006), 71281, doi.org/10.1155/IMRN/2006/71281.
- [12] BRUINIER, J. H. AND FUNKE J.: *On the injectivity of the Kudla–Millson lift and surjectivity of the Borcherds lift*. Moonshine: the first quarter century and beyond, 12–39, London Math. Soc. Lecture Note Ser., 372, Cambridge Univ. Press, Cambridge, (2010).
- [13] BRUMLEY, F. AND MARSHALL, S.: *Lower bounds for Maass forms on simisimple groups*. Compositio Math., **156** (2020), 959–1003.
- [14] BRUMLEY, F., MARSHALL, S.: *Concentration properties of theta lifts on orthogonal groups*, preprint arxiv:2309.06433.
- [15] BRUMLEY, F. AND TEMPLIER, N.: *Large values of cusp forms on GL_n* . Selecta Math., **26** (2020), 26–63.
- [16] BUMP, D.: *Automorphic forms and representations*, volume 5 of Cambridge Studies in Advanced Mathematics, Cambridge University Press, Cambridge, 1997.
- [17] DONNELEY, H.: *Exceptional sequences of eigenfunctions for hyperbolic manifolds*. Proc. Amer. Math. Soc., **135** (2007), 1551–1555.
- [18] DUISTERMAAT, J.: *On the similarity between the Iwasawa projection and the diagonal part*, Mémoires de la Soc. Math. de France, 2e série, tome **15** (1984), 129–138.
- [19] ERDÉLYI, A.: *Higher transcendental functions, volume 2*, New York : McGraw-Hill, 1953–55.

- [20] GAN, W.T., QIU, Y. AND TAKEDA, S.: *The regularized Siegel Weil formula (the second term identity) and the Rallis inner product formula*. Invent. math., **198** (2014), 739–831.
- [21] GANGOLLI, R. AND VARADARAJAN, V. S.: *Harmonic analysis of spherical functions on real reductive groups*. Ergeb. Math. Grenzgeb., **101** Springer-Verlag, Berlin, 1988.
- [22] HARCOS, G. AND MICHEL, P.: *The subconvexity problem for Rankin-Selberg L -functions and equidistribution of Heegner points II*. Invent. Math., **163** (2006), 581–655.
- [23] HARISH-CHANDRA: *Automorphic forms on semisimple Lie groups*. Notes by J. G. M. Mars. Lecture Notes in Mathematics, No. 62. Springer-Verlag, Berlin, 1968.
- [24] HIRAI, T.: *On irreducible representations of the Lorentz group of n -th order*. Proc. Japan Acad. **38** (1962), 258–262.
- [25] HOFFSTEIN, J. AND LOCKHART, P.: *Coefficients of Maass forms and the Siegel zero*. Ann. Math., **140** (1994), 161–176.
- [26] HOWE, R.: *θ -series and invariant theory*. Proceedings of Symposia in Pure Math., **33** part 1 (1979), 275–285.
- [27] HOWE, R. AND TAN, E. C.: *Homogeneous functions on light cones: the infinitesimal structure of some degenerate principal series representations*, Bull. Amer. Math. Soc. (N.S.), **28** (1993), 1–74.
- [28] IKEDA, T. AND KATSURADA, H.: *Estimates for the Fourier coefficients of the Duke-Imamoglu-Ikeda lift*. Forum Math., **35**(4) (2023), 975–990.
- [29] IWANIEC, H.: *Small eigenvalues of Laplacian for $\Gamma_0(N)$* . Acta Arith., **56** (1990), 65–82.
- [30] IWANIEC, H.: *Spectral Methods of Automorphic Forms, second edition*. Grad. Stud. Math., 53 American Mathematical Society, Providence, R. I. Revista Matemática Iberoamericana, Madrid, 2002, xii+220 pp.
- [31] IWANIEC, H. AND SARNAK, P.: *L^∞ norms of eigenfunctions of arithmetic surfaces*. Ann. Math., **141** (1995), 301–320.
- [32] KIM, H.: *Functoriality for the exterior square of GL_4 and the symmetric fourth of GL_2 . With appendix 1 by Dinakar Ramakrishnan and appendix 2 by Kim and Peter Sarnak*. J. Amer. Math. Soc., **16** (2003), 139–183.
- [33] KUDLA, S.: *Integrals of Borcherds Forms*. Compositio Math., **137** (2003), 293–349.
- [34] KUDLA, S. AND RALLIS, S.: *On the Weil-Siegel formula*. J. Reine Angew. Math., **387** (1988), 1–68.
- [35] LAPID, E., AND OFFEN, O.: *Compact unitary periods*. Compos. Math., **143** no. 2 (2007), 323–338.
- [36] LEVITAN, B. M.: *On the asymptotic behavior of the spectral function of a self-adjoint differential equation of second order*, Isv. Akad. Nauk SSSR Ser. Mat. **16** (1952), 325–352.
- [37] LI, Y., NARITA, H. AND PITALE, A.: *An explicit construction of non-tempered cusp forms on $O(1, 8n + 1)$* . Ann. Math. Quebec, **44** (2020), 349–384.
- [38] MAGNUS, W., OBERHETTINGER, F. AND SONI, R. P.: *Formulas and Theorems for the Special Functions of Mathematical Physics*, 3rd ed., Die Grundlehren der Mathematischen Wissenschaften, vol. 117, Springer, New York, 1966.
- [39] MARSHALL, S.: *L^p norms of higher rank eigenfunctions and bounds for spherical functions*, J. Eur. Math. Soc. **18** issue 7 (2016), 1437–1493.

- [40] MILIĆEVIĆ, D.: *Large values of eigenfunctions on arithmetic hyperbolic 3-manifolds*. Geom. Funct. Anal., **21** no. 6 (2011), 1375–1418.
- [41] MILIĆEVIĆ, D.: *Large values of eigenfunctions on arithmetic hyperbolic surfaces*. Duke Math. J. **155** no. 2 (2010), 365–401.
- [42] MUTO, M., NARITA, H. AND PITALE, A.: *Lifting to $GL(2)$ over a division quaternion algebra and an explicit construction of CAP representations*. Nagoya Math. J., **222**, issue 01 (2016), 137–185.
- [43] NARITA, H., PITALE, A. AND WAGH, S.: *An explicit lifting construction of CAP forms on $O(1, 5)$* . Int. J. Number Theory, **19**, no. 6 (2023), 1337–1378.
- [44] PITALE, A., SAHA, A. AND SCHMIDT, R.: *On the standard L -function for $GSp_{2n} \times GL_1$ and algebraicity of symmetric fourth L -values for GL_2* . Ann. Math. Quebec., **45** (2020), 113–159.
- [45] RALLIS, S.: *Injectivity properties of lifting associated to Weil representations*. Compositio Math., **52** (1984), 139–169.
- [46] RAMACHAR, P. AND WAKATSUKI, S.: *Subconvex bounds for Hecke-Maass forms on compact arithmetic quotients of semisimple Lie groups*, Math. Z., **298** (2021), 1383–1424.
- [47] RATCLIFFE, J.: *Foundations of Hyperbolic Manifolds*, second edition. Graduate Texts in Mathematics 149, Springer, (2006).
- [48] ROELCKE, W.: *Über die Wellengleichung bei Grenzkreisgruppen erster Art*. Sitzungsberichte der Heidelberger Akademie der Wissenschaften, Springer-Verlag Berlin Heidelberg, 1956.
- [49] RUDNICK, Z. AND SARNAK, P.: *The behavior of eigenstates of arithmetic hyperbolic manifolds*, Communications in Math. Physics, **161** (1994), 195–213.
- [50] SAHA, A.: *On sup-norms of cusp forms of powerful level*. J. Eur. Math. Soc. (JEMS). **19** no. 11 (2017), 3549–3573.
- [51] SAHA, A.: *Hybrid sup-norm bounds for Maass newforms of powerful level*. Algebra Number Theory. **11** no. 5 (2017), 1009–1045.
- [52] SARNAK, P.: *Letter to Morawetz*. <http://www.math.princeton.edu/sarnak>
- [53] SERRE, J. P.: *A course in arithmetic*. Graduate text in math., vol. 7, Springer-Verlag, New York, (1973).
- [54] SHIMURA, G.: *Arithmetic and analytic theories of quadratic forms and Clifford groups*, Mathematical surveys and monographs, vol. 109, American Mathematical Society (2004).
- [55] XIA, H.: *On L^∞ -norms of holomorphic cusp forms*, J. Number Th., **124** (2007), 325–327.
- [56] WEIL, A.: *Sur certains groupes d’opérateurs unitaires*, Acta Math., **111** (1964), 143–211.
- [57] WEIL, A.: *Sur la formule de Siegel dans la théorie des groupes classiques*, Acta Math., **113** (1965), 143–211.