

---

**I.** Let  $D$  be the unit disc consisting of all  $(x, y)$  for which  $x^2 + y^2 \leq 1$ . Let  $f(x, y) = x^2 - y^2$ . Use the methods (8) of section 12.7 (i. e. critical points and investigation of the values of  $f$  to the boundary of  $D$ ) to determine the maximum value of  $f$  on  $D$ , and the point or points at which it occurs.

**II.** Let  $C$  be the unit circle where  $x^2 + y^2 = 1$ . Let  $f(x, y) = x^2 - y^2$ . Write the (three) equations that would need to be solved if the method of Lagrange multipliers is to be used to find the maximum value of  $f$  on  $C$ , but do *not* try to solve the equations or proceed further with finding the maximum. (4)



---

**IV.** State the Extreme Value Theorem for functions of two variables.

(3)

**V.**

(10)

1. Draw a (reasonably large-sized) picture showing a portion of the graph  $z = f(x, y)$  for a typical function  $f(x, y)$ , where  $f(x, y) > 0$ , which has domain  $R$  consisting of all  $(x, y)$  with  $0 \leq x \leq 1$  and  $0 \leq y \leq 1$ .

2. Using the picture for part (a), draw a typical cross-section of the region under the graph and above the  $xy$ -plane for a fixed value  $y_0$  of  $y$ . Write an expression for its area.

3. Using the picture for part (a), draw a vector  $\vec{v}$  tangent to the graph  $z = f(x, y)$  at a typical point  $(x_0, y_0, f(x_0, y_0))$ , so that the  $\vec{i}$  component of  $\vec{v}$  is  $0 \cdot \vec{i}$  and the  $\vec{j}$  component of  $\vec{v}$  is  $1 \cdot \vec{j}$ . Write an expression for the  $\vec{k}$  component of  $\vec{v}$ .

**VI.** Evaluate the following integral by reversing the order of integration:  $\int_0^1 \int_{3y}^3 e^{x^2} dx dy$ .  
(7)

**VII.** Write an integral in spherical coordinates whose value is the volume of the upper hemisphere of radius  $R$  where  $x^2 + y^2 + z^2 \leq R^2$ ,  $z \geq 0$ . Carry out the integration to determine that the volume is  $2\pi R^3/3$ .  
(7)