

# Sharp Sobolev Trace Inequalities on Riemannian Manifolds with Boundaries

YANYAN LI

*Rutgers University*

AND

MEIJUN ZHU

*Rutgers University and University of British Columbia*

## Abstract

In this paper, we establish some sharp Sobolev trace inequalities on  $n$ -dimensional, compact Riemannian manifolds with smooth boundaries. More specifically, let

$$q = 2(n-1)/(n-2), \quad \frac{1}{S} = \inf \left\{ \int_{\mathbb{R}_+^n} |\nabla u|^2 : \nabla u \in L^2(\mathbb{R}_+^n), \int_{\partial \mathbb{R}_+^n} |u|^q = 1 \right\}.$$

We establish for any Riemannian manifold with a smooth boundary, denoted as  $(M, g)$ , that there exists some constant  $A = A(M, g) > 0$ ,  $(\int_{\partial M} |u|^q ds_g)^{2/q} \leq S \int_M |\nabla_g u|^2 dv_g + A \int_{\partial M} u^2 ds_g$ , for all  $u \in H^1(M)$ . The inequality is sharp in the sense that the inequality is false when  $S$  is replaced by any smaller number. © 1997 John Wiley & Sons, Inc.

## 0 Introduction

It is well-known that sharp Sobolev-type inequalities are important in the study of partial differential equations, especially those that arise in geometry and physics. There has been much work on such inequalities and their applications (see, for example, Trudinger [35], Moser [30], Aubin [3], Talenti [34], Lieb [27, 28], Brezis-Nirenberg [9], Cherrier [13], Brezis-Lieb [8], Carleson-Chang [11], Escobar [14, 16], Beckner [6], Adimurthi and Yadava [1], Hebey and Vaugon [21, 20], Hebey [19], and the references therein).

For  $n \geq 3$ , it was shown by Aubin [3] and Talenti [34] that, for  $p = 2n/(n-2)$ ,

$$\frac{1}{S_1} = \inf \left\{ \frac{\int_{\mathbb{R}^n} |\nabla u|^2}{(\int_{\mathbb{R}^n} |u|^p)^{2/p}} \mid u \in L^p(\mathbb{R}^n) \setminus \{0\}, \nabla u \in L^2(\mathbb{R}^n) \right\}$$

is achieved and the extremal functions are found. In particular,

$$(A) \quad \frac{1}{S_1} = \pi n(n-2) \left( \frac{\Gamma(n/2)}{\Gamma(n)} \right)^{2/n}.$$

It was shown by P. L. Lions [29] that, for  $q = 2(n-1)/(n-2)$ ,

$$(0.1) \quad \frac{1}{S} = \inf \left\{ \frac{\int_{\mathbb{R}_+^n} |\nabla u|^2}{(\int_{\partial\mathbb{R}_+^n} |u|^q)^{2/q}} \mid \nabla u \in L^2(\mathbb{R}_+^n), u \in L^q(\partial\mathbb{R}_+^n) \setminus \{0\} \right\}$$

is achieved. The extremal functions were found independently by Escobar [16] and Beckner [6]. In particular,

$$(B) \quad \frac{1}{S} = \frac{n-2}{2} \sigma_n^{1/(n-1)},$$

where  $\sigma_n$  denotes the volume of the unit sphere in  $\mathbb{R}^n$ .

In this paper we study some Sobolev-type trace inequalities on Riemannian manifolds with boundaries. Throughout this paper we denote  $p = 2n/(n-2)$ ,  $q = 2(n-1)/(n-2)$ , and  $S_1$  and  $S$  as in (A) and (B), respectively.

**THEOREM 0.1 (Main Theorem)** *For  $n \geq 3$ , let  $(M, g)$  be some smooth  $n$ -dimensional, compact, Riemannian manifold with a smooth boundary. Then there exists some constant  $A = A(M, g) > 0$  such that, for all  $u \in H^1(M)$ ,*

$$(0.2) \quad \left( \int_{\partial M} |u|^q ds_g \right)^{2/q} \leq S \int_M |\nabla_g u|^2 dv_g + A \int_{\partial M} u^2 ds_g,$$

where  $dv_g$  denotes the volume form of  $(M, g)$  and  $ds_g$  denotes the induced volume form on  $\partial M$ .

**REMARK 0.2** The constant  $S$  in front of  $\int_M |\nabla_g u|^2 dv_g$  is sharp. It cannot be replaced by any smaller number.

**REMARK 0.3** In general,  $\int_{\partial M} u^2 ds_g$  cannot be replaced by  $\int_{\partial M} u^r ds_g$  for  $r < 2$ . For instance, this is the case for any bounded domain in  $\mathbb{R}^n$  with the flat metric.

**REMARK 0.4** The above theorem in the special case  $n \geq 5$  and  $(M, g)$  a bounded, smooth domain in  $\mathbb{R}^n$  with the Euclidean metric was obtained by Adimurthi and Yadava in [1]. Our method in proving Theorem 0.1 is different from theirs.

**REMARK 0.5** Clearly we only need to consider the case when  $M$  is connected. Throughout the paper, we assume this.

The present work is stimulated by some recent work of Hebey and Vaugon [21], where they proved a conjecture of Aubin [4]: For  $n \geq 3$  and  $(M, g)$  any

smooth  $n$ -dimensional, compact manifold without boundary, there exists some constant  $C > 0$  such that, for all  $u \in H^1(M)$ ,

$$(0.3) \quad \left( \int_M |u|^p dv_g \right)^{2/p} \leq S_1 \int_M |\nabla_g u|^2 dv_g + C \int_M u^2 dv_g.$$

One of the main ingredients in their proof of (0.3) is, through the Moser iteration technique, to obtain an appropriate upper bound for blowup minimum-type solutions to certain critical exponent equations with the zero Dirichlet boundary condition. Such asymptotic analysis was obtained by Han in [18], using the Moser iteration technique, for blowup minimum-type solutions to certain critical exponent equations with the zero Dirichlet boundary conditions in general domains in  $\mathbb{R}^n$ , which extend results of Atkinson and Peletier [2] and Brezis and Peletier [10] on balls in  $\mathbb{R}^n$ . Such extension was conjectured by Brezis and Peletier and was proven by Rey [31] using a different method.

During the past few years, energy-independent asymptotic analysis for blowup solutions to certain critical exponent equations has been obtained. See Schoen [32], Zhang [36], Chang, Gursky, and Yang [12], Li [24, 25, 23], Schoen and Zhang [33], and the references therein.

As in [21], one of the main ingredients in our proof of Theorem 0.1 is some asymptotic analysis for blowup minimum-type solutions. However, we need to overcome new difficulties since what we encounter here are certain nonlinear Neumann boundary conditions rather than zero Dirichlet boundary conditions as in [18] and [21]. Moreover, Theorem 0.1 for  $n = 3$  is subtler: In addition to the upper bound of solutions obtained by the Moser iteration technique, we also need to obtain an appropriate lower bound.

Another main ingredient is local balance checking via the Pohozaev identity. Using similar methods, we have established some other Sobolev-type inequalities. In particular, we have extended theorem 1 in [1] from dimension  $n \geq 5$  to  $n \geq 3$ . This will be addressed in a forthcoming paper.

## 1 Preliminary Estimates

We first present two weaker inequalities from which one can deduce that minimum-type solutions can blow up at only one point. Although this step is well-known, we include a proof here for the reader's convenience.

**PROPOSITION 1.1** *For all  $\varepsilon > 0$ , there exists some constant  $B(\varepsilon)$  depending only on  $\varepsilon$ ,  $M$ , and  $g$  such that*

$$\begin{aligned} & \left( \int_{\partial M} |u|^q ds_g \right)^{2/q} \\ & \leq (S + \varepsilon) \int_M |\nabla_g u|^2 dv_g + B(\varepsilon) \int_M u^2 dv_g, \quad \forall u \in H^1(M). \end{aligned}$$

PROOF: By partition of unity, it follows easily from (0.1). We omit the details.  $\blacksquare$

PROPOSITION 1.2 *For all  $\varepsilon > 0$ , there exists some constant  $A(\varepsilon)$  depending only on  $\varepsilon$ ,  $M$ , and  $g$  such that for all  $u \in H^1(M)$ ,*

$$(1.1) \quad \left( \int_{\partial M} |u|^q ds_g \right)^{2/q} \leq (S + \varepsilon) \int_M |\nabla_g u|^2 dv_g + A(\varepsilon) \int_M u^2 dv_g.$$

PROOF: We prove this proposition using an argument by contradiction. Suppose the contrary of (1.1), namely, that there exists some constant  $\delta > 0$  such that for all  $\alpha > 1$ ,

$$(1.2) \quad \xi_\alpha := \inf_{H^1(M) \setminus \{0\}} \frac{\int_M |\nabla_g u|^2 dv_g + \alpha \int_{\partial M} u^2 ds_g}{(\int_{\partial M} |u|^q ds_g)^{2/q}} \leq \frac{1}{S} - \delta.$$

*Claim.* There exists some nonnegative function  $u_\alpha \in H^1(M)$  satisfying

$$(1.3) \quad \xi_\alpha = \int_M |\nabla_g u_\alpha|^2 dv_g + \alpha \int_{\partial M} u_\alpha^2 ds_g, \quad \int_{\partial M} u_\alpha^q ds_g = 1.$$

PROOF OF CLAIM: We sketch this well-known proof for the reader's convenience. Let  $\{u^{(m)}\}$  be a minimizing sequence with

$$\|u^{(m)}\|_{q,\partial M} = \left( \int_{\partial M} |u^{(m)}|^q ds_g \right)^{1/q} = 1$$

and  $u^{(m)} \geq 0$ . Clearly,  $\|u^{(m)}\|_{H^1(M)} \leq C$ . After passing to a subsequence,  $u^{(m)}$  converges weakly to some  $u \in H^1(M)$ ,  $u \geq 0$ . It is not difficult to see that

$$\int_{\partial M} (|u^{(m)}|^q - |u^{(m)} - u|^q) ds_g = \int_{\partial M} u^q ds_g + o(1),$$

and consequently

$$\int_{\partial M} |u^{(m)} - u|^q ds_g \leq 1 + o(1), \quad \int_{\partial M} u^q ds_g \leq 1,$$

where  $o(1)$  denotes some quantity tending to zero as  $m$  tends to  $\infty$ .

Therefore, by the Sobolev embedding theorems and Proposition 1.1, we have, for  $\varepsilon_0 > 0$ ,

$$\begin{aligned}
\xi_\alpha &= \int_M |\nabla u^{(m)}|^2 + \alpha \int_{\partial M} |u^{(m)}|^2 + o(1) \\
&= \int_M |\nabla(u^{(m)} - u)|^2 + \int_M |\nabla u|^2 + \alpha \|u\|_{2,\partial M}^2 + o(1) \\
&= \int_M |\nabla(u^{(m)} - u)|^2 + \frac{B(\varepsilon_0)}{S + \varepsilon_0} \int_M |u^{(m)} - u|^2 + \int_M |\nabla u|^2 \\
&\quad + \alpha \|u\|_{2,\partial M}^2 + o(1) \\
&\geq \frac{1}{S + \varepsilon_0} \left( \int_{\partial M} |u^{(m)} - u|^q \right)^{2/q} + \xi_\alpha \left( \int_{\partial M} u^q \right)^{2/q} + o(1) \\
&\geq \frac{1}{S + \varepsilon_0} \int_{\partial M} |u^{(m)} - u|^q + \xi_\alpha \int_{\partial M} u^q + o(1) \\
&= \left( \frac{1}{S + \varepsilon_0} - \xi_\alpha \right) \int_{\partial M} |u^{(m)} - u|^q + \xi_\alpha + o(1).
\end{aligned}$$

Choose  $\varepsilon_0 > 0$  small so that  $\frac{1}{S + \varepsilon_0} - \xi_\alpha \geq \delta/2$ ; we have from the above that  $\int_{\partial M} |u^{(m)} - u|^q = o(1)$ . It follows easily that  $u$  is a minimum of (1.2). ■

Now let  $u_\alpha$  be some nonnegative function in  $H^1(M)$  satisfying (1.3). It is easy to see from (1.3) that  $\|u_\alpha\|_{H^1(M)}$  is bounded by some constant independent of  $\alpha$ . It follows that, after passing to some subsequence,  $u_\alpha$  weakly converges to some  $\bar{u} \in H^1(M)$ . This leads to

$$(1.4) \quad \int_M |u_\alpha - \bar{u}|^2 dv_g + \int_{\partial M} |u_\alpha - \bar{u}|^2 ds_g = o(1),$$

and therefore, in view of (1.3),

$$(1.5) \quad \bar{u} = 0 \quad \text{on } \partial M.$$

Here and in the following,  $o(1)$  denotes some quantity tending to zero as  $\alpha$  tends to  $\infty$ .

Therefore, by Proposition 1.1, (1.4), and (1.5), we have, for  $\varepsilon_0 > 0$ ,

$$\begin{aligned}
\xi_\alpha &= \int_M |\nabla u_\alpha|^2 + \alpha \int_{\partial M} |u_\alpha|^2 \\
&= \int_M |\nabla(u_\alpha - \bar{u})|^2 + \int_M |\nabla \bar{u}|^2 + \alpha \|u_\alpha\|_{2,\partial M}^2 + o(1)
\end{aligned}$$

$$\begin{aligned}
&\geq \int_M |\nabla(u_\alpha - \bar{u})|^2 + o(1) \\
&\geq \frac{1}{S + \varepsilon_0} \left( \int_{\partial M} |u_\alpha - \bar{u}|^q \right)^{2/q} + o(1) \\
&= \frac{1}{S + \varepsilon_0} + o(1).
\end{aligned}$$

Sending  $\alpha$  to  $\infty$ , we obtain from the above and (1.2) that

$$\frac{1}{S} - \delta \geq \frac{1}{S + \varepsilon_0}.$$

Sending  $\varepsilon_0$  to zero, we reach a contradiction.  $\blacksquare$

## 2 Asymptotic Analysis

From now on, we begin to prove Theorem 0.1 through an argument by contradiction. Suppose the contrary of Theorem 0.1 is true; then we have, for all  $\alpha \geq 1$ ,

$$(2.1) \quad \xi_\alpha < \frac{1}{S},$$

where  $\xi_\alpha$  is defined in (1.2). As in Section 1, there exists some nonnegative function  $u_\alpha \in H^1(M)$  satisfying (1.3). It follows that  $u_\alpha$  satisfies

$$(2.2) \quad \begin{cases} -\Delta_g u_\alpha = 0 & \text{in } M \\ \frac{\partial_g u_\alpha}{\partial \nu} = \xi_\alpha u_\alpha^{q-1} - \alpha u_\alpha & \text{on } \partial M. \end{cases}$$

In this section we establish, by using the Moser iteration technique, an appropriate upper bound for  $u_\alpha$ .

For all  $\varepsilon > 0$ , it follows from (1.3), (2.1), and Proposition 1.2 that there exists some  $A(\varepsilon)$  such that

$$\begin{aligned}
1 + \frac{\varepsilon}{S} &> (S + \varepsilon) \xi_\alpha \\
&= (S + \varepsilon) \|\nabla_g u_\alpha\|_{2,M}^2 + \alpha(S + \varepsilon) \|u_\alpha\|_{2,\partial M}^2 \\
&\geq \left( \int_{\partial M} |u_\alpha|^q \right)^{2/q} + [\alpha(S + \varepsilon) - A(\varepsilon)] \|u_\alpha\|_{2,\partial M}^2 \\
&= 1 + [\alpha(S + \varepsilon) - A(\varepsilon)] \|u_\alpha\|_{2,\partial M}^2.
\end{aligned}$$

Sending  $\alpha$  to  $\infty$ , we have

$$(S + \varepsilon) \liminf_{\alpha \rightarrow \infty} \xi_\alpha \geq 1 \quad \text{and} \quad 1 + \frac{\varepsilon}{S} \geq 1 + (S + \varepsilon) \limsup_{\alpha \rightarrow \infty} \alpha \|u_\alpha\|_{2,\partial M}^2.$$

Sending  $\varepsilon$  to 0, we have, by using (2.1), that

$$(2.3) \quad \lim_{\alpha \rightarrow \infty} \xi_\alpha = \frac{1}{S}$$

and

$$(2.4) \quad \lim_{\alpha \rightarrow \infty} \alpha \|u_\alpha\|_{2,\partial M}^2 = 0.$$

PROPOSITION 2.1 *There exists  $\bar{x}_\alpha \in \partial M$  such that for all  $\delta > 0$ ,*

$$\lim_{\alpha \rightarrow \infty} \int_{B_\delta(\bar{x}_\alpha) \cap \partial M} u_\alpha^q = 1.$$

Before proving the previous proposition, we present a well-known lemma (see, e.g., [5] for results of this type).

LEMMA 2.2 *Suppose  $\{y_\alpha\} \in \partial M$  for a sequence of  $\alpha \rightarrow \infty$  satisfies, for some  $0 < \beta < 1$ ,  $\delta > 0$ ,*

$$(2.5) \quad \int_{B_\delta(y_\alpha) \cap \partial M} u_\alpha^q \leq \beta.$$

Then

$$(2.6) \quad \lim_{\alpha \rightarrow \infty} \int_{B_{\delta/2}(y_\alpha) \cap \partial M} u_\alpha^q = 0.$$

PROOF: Let  $\eta = \eta_\alpha \in C^\infty(\overline{M})$  be some cutoff function satisfying

$$\eta = \begin{cases} 1 & \text{in } B_{\delta/2}(y_\alpha) \cap \overline{M} \\ 0 & \text{in } \overline{M} \setminus B_\delta(y_\alpha) \end{cases}$$

and

$$|\nabla_g \eta| + |\nabla_g^2 \eta| \leq C(\delta, M, g).$$

For  $1 < r \leq q - 1$ , multiplying the first equation in (2.2) by  $\eta^2 u_\alpha^r$  and integrating by parts, we obtain, by using the boundary condition of  $u_\alpha$  in (2.2), that

$$\int_M \nabla_g u_\alpha \cdot \nabla_g (\eta^2 u_\alpha^r) \, dv_g = \xi_\alpha \int_{\partial M} \eta^2 u_\alpha^{q-1+r} \, ds_g - \alpha \int_{\partial M} \eta^2 u_\alpha^{r+1} \, ds_g.$$

Direct calculation yields

$$\begin{aligned}
 & \int_M \nabla_g u_\alpha \cdot \nabla_g (\eta^2 u_\alpha^r) dv_g \\
 (2.7) = & \frac{4r}{(r+1)^2} \int_M |\nabla_g (u_\alpha^{(r+1)/2} \eta)|^2 dv_g + \frac{r-1}{(r+1)^2} \int_M u_\alpha^{r+1} \Delta_g (\eta^2) dv_g \\
 & - \frac{4r}{(r+1)^2} \int_M u_\alpha^{r+1} |\nabla_g \eta|^2 dv_g - \frac{r-1}{(r+1)^2} \int_{\partial M} u_\alpha^{r+1} \frac{\partial_g (\eta^2)}{\partial \nu} ds_g.
 \end{aligned}$$

It follows that

$$\begin{aligned}
 & \int_M |\nabla_g (u_\alpha^{(r+1)/2} \eta)|^2 \\
 & = -\frac{r-1}{4r} \int_M u_\alpha^{r+1} \Delta_g (\eta^2) + \int_M u_\alpha^{r+1} |\nabla_g \eta|^2 + \frac{r-1}{4r} \int_{\partial M} u_\alpha^{r+1} \frac{\partial_g (\eta^2)}{\partial \nu} \\
 & + \frac{\xi_\alpha (r+1)^2}{4r} \int_{\partial M} u_\alpha^{q-1+r} \eta^2 - \frac{\alpha (r+1)^2}{4r} \int_{\partial M} u_\alpha^{r+1} \eta^2 \\
 & \leq \frac{\xi_\alpha (r+1)^2}{4r} \int_{\partial M} u_\alpha^{q-1+r} \eta^2 + C(\delta, r) \left\{ \int_M u_\alpha^{r+1} + \int_{\partial M} u_\alpha^{r+1} \right\}.
 \end{aligned}$$

Using (1.3), the fact that  $r+1 \leq q < p$ , and the Sobolev embedding theorems, we know that

$$\begin{aligned}
 & \int_M u_\alpha^{r+1} + \int_{\partial M} u_\alpha^{r+1} \\
 (2.8) \quad & \leq C(r, M) \left\{ \int_M |\nabla_g u_\alpha|^2 + \int_{\partial M} u_\alpha^2 \right\}^{(r+1)/2} \\
 & \leq C(r, M).
 \end{aligned}$$

Consequently,

$$(2.9) \quad \int_M |\nabla_g (u_\alpha^{(r+1)/2} \eta)|^2 \leq \frac{\xi_\alpha (r+1)^2}{4r} \int_{\partial M} u_\alpha^{q-1+r} \eta^2 + C(\delta, r, M).$$

Applying Hölder's inequality and then Proposition 1.1 to  $u = u_\alpha^{(r+1)/2} \eta$  gives, for all  $\varepsilon > 0$ ,

$$\begin{aligned}
 (2.10) \quad & \int_{\partial M} u_\alpha^{q-1+r} \eta^2 \\
 & \leq \left( \int_{B_\delta(y_\alpha) \cap \partial M} u_\alpha^q \right)^{(q-2)/q} \left( \int_{\partial M} (u_\alpha^{(r+1)/2} \eta)^q \right)^{2/q}
 \end{aligned}$$

$$\begin{aligned} &\leq \left( \int_{B_\delta(y_\alpha) \cap \partial M} u_\alpha^q \right)^{(q-2)/q} (S + \varepsilon) \left\| \nabla_g (u_\alpha^{(r+1)/2} \eta) \right\|_{2,M}^2 \\ &\quad + B(\varepsilon) \left( \int_{B_\delta(y_\alpha) \cap \partial M} u_\alpha^q \right)^{(q-2)/q} \left\| u_\alpha^{(r+1)/2} \eta \right\|_{2,M}^2. \end{aligned}$$

Using the fact that  $\beta < 1$ , we now fix some  $r \in (1, q-1]$  and  $\varepsilon > 0$  satisfying

$$\frac{S + \varepsilon}{S - \varepsilon} \beta^{(q-2)/q} \frac{(r+1)^2}{4r} \leq 1 - \varepsilon.$$

Consequently, in view of (2.3), we have for  $\alpha$  large

$$\xi_\alpha (S + \varepsilon) \beta^{(q-2)/q} \frac{(r+1)^2}{4r} \leq 1 - \varepsilon.$$

Combining (2.9), (2.10), (2.8), (2.5), and the above, we obtain

$$(2.11) \quad \int_M |\nabla_g (u_\alpha^{(r+1)/2} \eta)|^2 \leq C(\delta, r, \varepsilon, \beta, M).$$

It follows from the Sobolev embedding theorems, (2.8), and (2.11) that

$$\begin{aligned} &\int_{B_{\delta/2}(y_\alpha) \cap \partial M} u_\alpha^{(r+1)q/2} \\ (2.12) \quad &\leq \int_{\partial M} (u_\alpha^{(r+1)/2} \eta)^q \\ &\leq C(M) \left\{ \left\| \nabla_g (u_\alpha^{(r+1)/2} \eta) \right\|_{2,M}^q + \left\| u_\alpha^{(r+1)/2} \eta \right\|_{2,M}^q \right\} \\ &\leq C(\delta, r, \beta, \varepsilon, M). \end{aligned}$$

Since  $(r+1)q/2 > q$ , we can derive (2.6) from (2.4), (2.12), and Hölder's inequality. Lemma 2.2 is thereby established.  $\blacksquare$

PROOF OF PROPOSITION 2.1: For  $x \in \partial M$ , we define  $\delta_{x,\alpha} > 0$  by

$$(2.13) \quad \int_{B_{\delta_{x,\alpha}}(x) \cap \partial M} u_\alpha^q = \frac{1}{2}.$$

Clearly,  $\inf_{x \in \partial M} \delta_{x,\alpha} > 0$ . We pick  $\bar{x}_\alpha \in \partial M$  satisfying

$$(2.14) \quad \delta_{\bar{x}_\alpha, \alpha} \leq 2 \inf_{x \in \partial M} \delta_{x,\alpha}.$$

We claim that  $\{\bar{x}_\alpha\}$  satisfies the property stated in the proposition. Suppose the contrary; then there exists some  $\delta > 0$ ,  $0 < \beta < 1$ , and a sequence of  $\alpha \rightarrow \infty$  such that

$$\int_{B_\delta(\bar{x}_\alpha) \cap \partial M} u_\alpha^q \leq \beta.$$

This, according to Lemma 2.2, implies

$$\lim_{\alpha \rightarrow \infty} \int_{B_{\delta/2}(\bar{x}_\alpha) \cap \partial M} u_\alpha^q = 0.$$

Therefore, in view of (2.13), we have for large  $\alpha$

$$\delta_{\bar{x}_\alpha, \alpha} \geq \frac{\delta}{2}.$$

This, together with (2.14), yields for large  $\alpha$

$$(2.15) \quad \delta_{x, \alpha} \geq \frac{\delta}{4}, \quad \forall x \in \partial M.$$

Clearly,  $\bigcup_{x \in \partial M} B_{\delta/8}(x)$  is an open cover of  $\partial M$ . Due to the compactness of  $\partial M$ , there exist  $x_1, \dots, x_m \in \partial M$  such that  $\partial M \subset \bigcup_{i=1}^m B_{\delta/8}(x_i)$ . We see from (2.15) and (2.13) that

$$\int_{B_{\delta/4}(x_i) \cap \partial M} u_\alpha^q \leq \frac{1}{2}, \quad 1 \leq i \leq m.$$

We can then apply Lemma 2.2 with  $\delta$  replaced by  $\delta/4$ ,  $\beta = 1/2$ , and  $y_\alpha = x_i$  to conclude that

$$\lim_{\alpha \rightarrow \infty} \int_{\partial M} u_\alpha^q \leq \lim_{\alpha \rightarrow \infty} \sum_{i=1}^m \int_{B_{\delta/8}(x_i) \cap \partial M} u_\alpha^q = 0.$$

This contradicts (1.3). Proposition 2.1 is thus established.  $\blacksquare$

Let  $x_\alpha \in \overline{M}$  be some maximum point of  $u_\alpha$ , that is,

$$u_\alpha(x_\alpha) = \max_{\overline{M}} u_\alpha.$$

It follows from the maximum principle that  $x_\alpha \in \partial M$  unless  $u_\alpha$  is identically equal to a constant. It is easy to see from (1.2) and (1.3) that  $u_\alpha$  is not identically equal to a constant for  $\alpha$  large. Therefore,  $x_\alpha \in \partial M$  for large

$\alpha$ . Set  $\mu_\alpha = u_\alpha(x_\alpha)^{-2/(n-2)}$ . Since  $\frac{\partial_g u_\alpha}{\partial \nu}(x_\alpha) \geq 0$ , we see from (2.2) that  $\alpha u_\alpha(x_\alpha) \leq \xi_\alpha u_\alpha(x_\alpha)^{q-1}$ , that is,

$$(2.16) \quad \alpha \mu_\alpha \leq \xi_\alpha \leq C.$$

It follows that

$$\lim_{\alpha \rightarrow \infty} \mu_\alpha = 0.$$

Let  $(y^1, \dots, y^{n-1}, y^n)$  denote some geodesic normal coordinates given by the exponential map  $\exp_{x_\alpha}$  with  $\frac{\partial}{\partial y^n}$  being the unit inner normal of  $M$  at  $y = 0$ . In this coordinate system, the metric  $g$  is given by  $g_{ij}(y) dy^i dy^j$ . For suitably small  $\delta_1 > 0$  (independent of  $\alpha$ ), we define  $v_\alpha$  in a neighborhood of  $z = 0$  by

$$v_\alpha(z) = u_\alpha(x_\alpha)^{-1} u_\alpha(\exp_{x_\alpha}(\mu_\alpha z)), \quad z \in O_\alpha \subset \mathbb{R}^n,$$

where

$$(2.17) \quad O_\alpha = \left\{ z \in \mathbb{R}^n : |z| < \frac{\delta_1}{\mu_\alpha}, \exp_{x_\alpha}(\mu_\alpha z) \in M \right\}.$$

We write  $\partial O_\alpha = \Gamma_\alpha^1 \cup \Gamma_\alpha^2$ , where

$$\begin{aligned} \Gamma_\alpha^1 &= \{z \in \partial O_\alpha : \exp_{x_\alpha}(\mu_\alpha z) \in \partial M\}, \\ \Gamma_\alpha^2 &= \{z \in \partial O_\alpha : \exp_{x_\alpha}(\mu_\alpha z) \in M\}. \end{aligned}$$

It follows from (2.2) that  $v_\alpha$  satisfies

$$(2.18) \quad \begin{cases} -\Delta_{g_\alpha} v_\alpha = 0 & \text{in } O_\alpha \\ \frac{\partial_{g_\alpha} v_\alpha}{\partial \nu} = \xi_\alpha v_\alpha^{q-1} - \alpha \mu_\alpha v_\alpha & \text{on } \Gamma_\alpha^1 \\ v_\alpha(0) = 1, \quad 0 \leq v_\alpha \leq 1, & \end{cases}$$

where  $g_\alpha$  denotes the metric on  $O_\alpha$  given by  $g_\alpha = g_{ij}(\mu_\alpha z) dz^i dz^j$ . It follows from (2.18), (2.16), and standard elliptic estimates (see, e.g., [17]) that for all  $R > 1$ ,

$$(2.19) \quad \|v_\alpha\|_{C^3(B_R \cap \overline{O}_\alpha)} \leq C(R), \quad \forall \alpha \geq 1.$$

Notice that because  $v_\alpha(0) = 1$ , we know from (2.19) that

$$(2.20) \quad \begin{cases} \int_{B_1(0) \cap \Gamma_\alpha^1} v_\alpha^q ds_{g_\alpha} \geq 1/C > 0 \\ \int_{B_1(0) \cap \Gamma_\alpha^1} v_\alpha^2 ds_{g_\alpha} \geq 1/C > 0. \end{cases}$$

It follows from the first inequality in (2.20) and Proposition 2.1 that

$$(2.21) \quad \lim_{\alpha \rightarrow \infty} |x_\alpha - \bar{x}_\alpha| = 0.$$

By change of variables, we have

$$\alpha \|u_\alpha\|_{2,\partial M}^2 \geq \alpha \int_{B_{\mu_\alpha}(x_\alpha) \cap \partial M} u_\alpha^2 = \alpha \mu_\alpha \int_{B_1 \cap \Gamma_\alpha^1} v_\alpha^2 ds_{g_\alpha}.$$

We derive from (2.4), the second inequality in (2.20), and the above that

$$(2.22) \quad \lim_{\alpha \rightarrow \infty} \alpha \mu_\alpha = 0.$$

It follows from (2.19) that there exists  $v \in C^2(\overline{\mathbb{R}}_+^n)$  such that along some subsequence,

$$(2.23) \quad \lim_{\alpha \rightarrow \infty} \|v_\alpha - v\|_{C^2(B_R \cap \overline{O}_\alpha)} = 0, \quad \forall R > 0,$$

where  $B_R = \{z \in \mathbb{R}^n : |z| < R\}$ . Clearly, in view of (2.18), (2.22), and (2.3),  $v$  satisfies

$$(2.24) \quad \begin{cases} \Delta v = 0 & \text{in } \mathbb{R}_+^n \\ \frac{\partial v}{\partial \nu} = \frac{1}{S} v^{q-1} & \text{on } \partial \mathbb{R}_+^n \\ v(0) = 1, \quad 0 < v \leq 1. \end{cases}$$

It follows from our earlier work [26] that

$$(2.25) \quad v(z', z_n) = \left( \frac{(n-2)^2 S^2}{|z'|^2 + (z_n + (n-2)S)^2} \right)^{(n-2)/2},$$

where  $z' = (z_1, \dots, z_{n-1})$ .

Due to the uniqueness of the limit function  $v$ , we know that (2.23) holds for the full limit  $\alpha \rightarrow \infty$ .

**PROPOSITION 2.3** *For  $\delta_1 = \delta_1(M, g) > 0$  small enough,*

$$\lim_{\alpha \rightarrow \infty} \int_{\Gamma_\alpha^1} |v_\alpha - v|^q = 0.$$

**PROOF:** Multiplying (2.24) by  $v$  and integrating by parts, we have

$$\int_{\mathbb{R}_+^n} |\nabla v|^2 = \frac{1}{S} \int_{\partial \mathbb{R}_+^n} v^q.$$

We also know that  $v$  is a minimum of (0.1), namely,

$$S \int_{\mathbb{R}^n_+} |\nabla v|^2 = \left( \int_{\partial\mathbb{R}^n_+} v^q \right)^{2/q}.$$

Thus,

$$(2.26) \quad \int_{\partial\mathbb{R}^n_+} v^q = 1.$$

It follows from (2.21) and Proposition 2.1 that

$$(2.27) \quad \lim_{\alpha \rightarrow \infty} \int_{\Gamma_\alpha^1} v_\alpha^q = \lim_{\alpha \rightarrow \infty} \int_{B_{\delta_1}(x_\alpha) \cap \partial M} u_\alpha^q = 1.$$

It is easy to see from (2.26) and the explicit form of  $v$  in (2.25) that

$$\lim_{\alpha \rightarrow \infty} \int_{\Gamma_\alpha^1} v^q = 1.$$

Therefore, for all  $\varepsilon > 0$ , there exists  $R = R(\varepsilon) > 1$  such that for  $\alpha$  large,

$$(2.28) \quad \int_{\Gamma_\alpha^1 \cap B_R} v^q > 1 - \varepsilon, \quad \int_{\Gamma_\alpha^1 \setminus B_R} v^q < 2\varepsilon.$$

Consequently, using the strong convergence of  $v_\alpha$  to  $v$  given in (2.23), we have, for  $\alpha$  large, that

$$(2.29) \quad \int_{\Gamma_\alpha^1 \cap B_R} |v_\alpha - v|^q < \varepsilon, \quad \int_{\Gamma_\alpha^1 \cap B_R} v_\alpha^q > 1 - 2\varepsilon.$$

We derive from (2.27) and the second inequality in (2.29) that for large  $\alpha$ ,

$$(2.30) \quad \int_{\Gamma_\alpha^1 \setminus B_R} v_\alpha^q \leq 3\varepsilon.$$

Combining the first inequality in (2.29), (2.30), and the second inequality in (2.28), we have for large  $\alpha$  that

$$\begin{aligned} \int_{\Gamma_\alpha^1} |v_\alpha - v|^q &\leq \int_{\Gamma_\alpha^1 \cap B_R} |v_\alpha - v|^q + 2^q \int_{\Gamma_\alpha^1 \setminus B_R} v_\alpha^q + 2^q \int_{\Gamma_\alpha^1 \setminus B_R} v^q \\ &\leq (1 + 2^{q+3})\varepsilon. \end{aligned}$$

Proposition 2.3 follows immediately. ■

Recall that the conformal Laplacian operator  $L_g$  and the conformal boundary operator  $B_g$  are given by (see, e.g., [15])

$$\begin{cases} L_g\psi = \Delta_g\psi - a(n)R_g\psi \\ B_g\psi = \frac{\partial_g\psi}{\partial\nu} + b(n)H_g\psi, \end{cases}$$

where  $a(n) = \frac{n-2}{4(n-1)}$ ,  $b(n) = \frac{n-2}{2}$ ,  $R_g$  is the scalar curvature of  $M$ , and  $H_g$  is the mean curvature of  $\partial M$  with respect to the inner normal of  $\partial M$  (e.g., the unit ball in  $\mathbb{R}^n$  has positive mean curvature).

Let  $\varphi$  be some  $C^2$ -positive function on  $\overline{M}$ , and let  $\hat{g} = \varphi^{4/(n-2)}g$ . It is well-known (see, e.g., [15]) that for all  $\psi \in H^1(M)$ ,

$$(2.31) \quad \begin{cases} L_{\hat{g}}(\psi/\varphi) = \varphi^{-(n+2)/(n-2)}L_g(\psi) & \text{in } M \\ B_{\hat{g}}(\psi/\varphi) = \varphi^{-n/(n-2)}B_g(\psi) & \text{on } \partial M. \end{cases}$$

Rewrite (2.2) as

$$(2.32) \quad \begin{cases} \Delta_g u_\alpha = 0 & \text{in } M \\ \frac{\partial_g u_\alpha}{\partial\nu} + b(n)H_g u_\alpha \\ \quad = \xi_\alpha u_\alpha^{q-1} - \alpha u_\alpha + b(n)H_g u_\alpha & \text{on } \partial M. \end{cases}$$

Setting  $w_\alpha = u_\alpha/\varphi$ , it follows from (2.31) that

$$(2.33) \quad \begin{cases} \Delta_g u_\alpha - a(n)R_g u_\alpha \\ \quad = \varphi^{(n+2)/(n-2)}(\Delta_{\hat{g}} w_\alpha - a(n)R_{\hat{g}} w_\alpha) & \text{in } M \\ \frac{\partial_g u_\alpha}{\partial\nu} + b(n)H_g u_\alpha \\ \quad = \varphi^{n/(n-2)}\left(\frac{\partial_g w_\alpha}{\partial\nu} + b(n)H_{\hat{g}} w_\alpha\right) & \text{on } \partial M. \end{cases}$$

We will choose an appropriate  $\varphi = \varphi_\alpha$  and then apply the Moser iteration technique to show that  $w_\alpha$  is bounded above by some constant independent of  $\alpha$ . Without loss of generality, we assume  $(M, g)$  is a smooth, bounded open set of a slightly larger Riemannian manifold  $(\widetilde{M}, \widetilde{g})$ . Let  $\gamma$  be the geodesic in  $\widetilde{M}$  with  $\gamma(0) = x_\alpha$ ,  $\gamma'(0) = \nu$ . Set  $P_\alpha = \gamma(t_\alpha \mu_\alpha)$  with  $t_\alpha = (n-2)/\xi_\alpha$ . Let  $(y^1, \dots, y^{n-1}, y^n)$  be some geodesic normal coordinate system of  $T_{P_\alpha} \widetilde{M}$  with  $\frac{\partial}{\partial y^n} = -\gamma'(t_\alpha \mu_\alpha)$ ,  $\exp_{P_\alpha} : T_{P_\alpha} \widetilde{M} \rightarrow \widetilde{M}$  denoting the exponential map, and  $g_{ij}(y) = \langle \frac{\partial}{\partial y^i}, \frac{\partial}{\partial y^j} \rangle$  denoting the metric of  $\widetilde{M}$ , with  $g_{ij}(0) = \delta_{ij}$  and  $\Gamma_{ij}^k(0) = 0$ , where  $\Gamma_{ij}^k$  is the Christoffel symbol. We define  $G_{P_\alpha}$  by

$$\begin{cases} -\Delta_g G_{P_\alpha} = n(n-2)\omega_n \delta_{P_\alpha} & \text{in } \widetilde{M} \\ G_{P_\alpha} = 0 & \text{on } \partial \widetilde{M}, \end{cases}$$

where  $\omega_n$  is the volume of the unit ball in  $\mathbb{R}^n$ . It follows from Appendix B that

$$G_{P_\alpha} \circ \exp_{P_\alpha}(y) = |y|^{2-n} + E(y),$$

where  $E(y)$  satisfies

$$(2.34) \quad |y|^{n-3}|E(y)| + |y|^{n-2}|\nabla_g E(y)| \leq C(\delta_1), \quad \forall |y| \leq \delta_1.$$

Define  $\varphi_\alpha : \overline{M} \rightarrow \mathbb{R}$  by

$$\varphi_\alpha = t_\alpha^{n-2} \mu_\alpha^{(n-2)/2} G_{P_\alpha}.$$

Clearly,  $\varphi_\alpha$  satisfies

$$-\Delta_g \varphi_\alpha = 0 \quad \text{in } M.$$

**PROPOSITION 2.4** *There exists some constant  $C$  depending only on  $(M, g)$  such that for all  $\alpha \geq 1$ ,*

$$u_\alpha \leq C\varphi_\alpha \quad \text{on } M.$$

**PROOF:** We only need to prove the proposition for  $\alpha$  large. Set  $w_\alpha = u_\alpha/\varphi_\alpha$  and  $\hat{g} = \varphi_\alpha^{4/(n-2)}g$ . Equation (2.33) holds in  $M$  for  $\varphi = \varphi_\alpha$ . Setting  $\psi = \varphi = \varphi_\alpha$  in (2.31), we have

$$(2.35) \quad \begin{cases} -a(n)[R_{\hat{g}} - R_g \varphi_\alpha^{-4/(n-2)}] = 0 & \text{in } M \\ \frac{\partial_{\hat{g}} \varphi_\alpha}{\partial \nu} + b(n)H_g \varphi_\alpha = b(n)H_{\hat{g}} \varphi_\alpha^{n/(n-2)} & \text{on } \partial M. \end{cases}$$

Combining (2.32), (2.33), and (2.35), we have

$$(2.36) \quad \begin{cases} \Delta_{\hat{g}} w_\alpha = 0 & \text{in } M, \\ \frac{\partial_{\hat{g}} w_\alpha}{\partial \nu} = \xi_\alpha w_\alpha^{q-1} - (\alpha \varphi_\alpha^{-2/(n-2)} + \frac{\partial_g \varphi_\alpha}{\partial \nu} \varphi_\alpha^{-n/(n-2)}) w_\alpha & \text{on } \partial M. \end{cases}$$

We need the following lemma to simplify (2.36):

**LEMMA 2.5** *For  $\alpha$  large,*

$$\alpha \varphi_\alpha^{-2/(n-2)} + \frac{\partial_g \varphi_\alpha}{\partial \nu} \varphi_\alpha^{-n/(n-2)} \geq 0 \quad \text{on } \partial M.$$

**PROOF:** Clearly Lemma 2.5 is equivalent to

$$(2.37) \quad -\frac{\partial_g \varphi_\alpha}{\partial \nu} \leq \alpha \varphi_\alpha \quad \text{on } \partial M.$$

Let  $0 < \delta_2 \ll \delta_1$ . It is clear from the proof how small we need  $\delta_2$  to be. It is independent of  $\alpha$ . Notice that  $G_{P_\alpha}$  is bounded below by some positive

constant independent of  $\alpha$  in  $M \setminus B_{\delta_2}(x_\alpha)$ ; also, the absolute values of its first derivatives are bounded above by some constant independent of  $\alpha$  in the same region. It is clear that (2.37) holds in  $M \setminus B_{\delta_2}(x_\alpha)$  for large  $\alpha$ .

In the  $y$ -coordinate,  $\partial M$  near  $x_\alpha$  is given by

$$y^n = t_\alpha \mu_\alpha + f(y'), \quad |y'| \leq \delta_1,$$

with  $f(0) = 0$ ,  $|\nabla_g^2 f(y')| \leq C(\delta_1)$ ,  $\forall |y'| \leq \delta_1$ . By the choice of coordinates,  $\frac{\partial}{\partial y^n}$  is orthogonal to the tangent space of  $\widetilde{M}$  at  $x_\alpha$ , which is spanned by  $\frac{\partial}{\partial y^i} + \frac{\partial f}{\partial y^i}(0) \frac{\partial}{\partial y^n}$ ,  $1 \leq i \leq n-1$ . Consequently, for  $1 \leq i \leq n-1$ ,

$$\frac{\partial f}{\partial y^i}(0) = -\frac{g_{in}}{g_{nn}} = O(\mu_\alpha^2).$$

It follows from the mean value theorem that

$$(2.38) \quad \begin{cases} f(y') = O(|y'|^2 + \mu_\alpha^2 |y'|) \\ \frac{\partial f}{\partial y^i}(y') = O(|y'| + \mu_\alpha^2), \quad 1 \leq i \leq n-1. \end{cases}$$

Therefore,

$$(2.39) \quad \mu_\alpha \leq C|y|, \quad \forall |y| \leq \delta_1, \quad y^n = t_\alpha \mu_\alpha + f(y').$$

It is not difficult to see that at  $\exp_{P_\alpha}(y', t_\alpha \mu_\alpha + f(y')) \in \partial M$ ,

$$(2.40) \quad \begin{aligned} \nu &= \sum_{i=1}^{n-1} \frac{\partial f}{\partial y^i} \frac{\partial}{\partial y^i} - \frac{\partial}{\partial y^n} + O(|y'|^2 + \mu_\alpha^2), \\ \frac{\partial \varphi_\alpha}{\partial y^i} &= -(n-2)t_\alpha^{n-2} \mu_\alpha^{(n-2)/2} y^i |y|^{-n} \\ &\quad + t_\alpha^{n-2} \mu_\alpha^{(n-2)/2} \frac{\partial E}{\partial y^i}(y), \quad 1 \leq i \leq n. \end{aligned}$$

It is easy to see from the definition of  $\varphi_\alpha$  and (2.34) that

$$(2.41) \quad \varphi_\alpha(y) \geq C^{-1} \mu_\alpha^{(n-2)/2} |y|^{2-n}, \quad |y| \leq \delta_2.$$

It follows that for all  $|y| \leq \delta_2$ ,  $y^n = t_\alpha \mu_\alpha + f(y')$ , we have

$$(2.42) \quad \begin{aligned} \frac{\partial_g \varphi_\alpha}{\partial \nu} &= \nabla_g \varphi_\alpha \cdot \nu = \sum_{l,i=1}^n g^{li} \frac{\partial \varphi_\alpha}{\partial y^i} \frac{\partial}{\partial y^l} \cdot \nu \\ &= -\frac{\partial \varphi_\alpha}{\partial y^n} + O(|y| |\nabla \varphi_\alpha|). \end{aligned}$$

Also, from (2.34), (2.40), and (2.41), we have that

$$(2.43) \quad |y| |\nabla \varphi_\alpha| \leq C \varphi_\alpha(y), \quad \forall |y| \leq \delta_2, \quad y^n = t_\alpha \mu_\alpha + f(y').$$

Combining (2.42) and (2.43), we have

$$(2.44) \quad \frac{\partial \varphi_\alpha}{\partial \nu} \geq -\frac{\partial \varphi_\alpha}{\partial y^n} - C \varphi_\alpha(y), \quad \forall |y| \leq \delta_1, \quad y^n = t_\alpha \mu_\alpha + f(y').$$

It follows from (2.34) and (2.41) that

$$\mu_\alpha^{(n-2)/2} |\nabla E(y)| \leq C \varphi_\alpha(y).$$

Using (2.38) and the above, we have, for  $|y| \leq \delta_1$ ,  $y^n = t_\alpha \mu_\alpha + f(y')$ , that

$$\begin{aligned} -\frac{\partial \varphi_\alpha}{\partial y^n} &\geq (n-2)t_\alpha^{n-2} \mu_\alpha^{(n-2)/2} y^n |y|^{-n} - C \varphi_\alpha(y) \\ &= (n-2)t_\alpha^{n-2} \mu_\alpha^{(n-2)/2} (t_\alpha \mu_\alpha) |y|^{-n} \\ &\quad + (n-2)t_\alpha^{n-2} \mu_\alpha^{(n-2)/2} f(y') |y|^{-n} - C \varphi_\alpha(y) \\ &\geq (n-2)t_\alpha^{n-2} \mu_\alpha^{(n-2)/2} (t_\alpha \mu_\alpha) |y|^{-n} - C \varphi_\alpha(y) \\ &\geq -C \varphi_\alpha(y). \end{aligned}$$

Lemma 2.5 follows from (2.44) and the above since, as pointed out earlier, (2.37) easily holds in  $M \setminus B_{\delta_2}(x_\alpha)$  for large  $\alpha$ .  $\blacksquare$

It follows from Lemma 2.5 and (2.36) that  $w_\alpha$  satisfies

$$(2.45) \quad \begin{cases} \Delta_{\hat{g}} w_\alpha = 0 & \text{in } M \\ \frac{\partial_{\hat{g}} w_\alpha}{\partial \nu} \leq \xi_\alpha w_\alpha^{q-1} & \text{on } \partial M. \end{cases}$$

Let  $\eta$  be some smooth, nonnegative cutoff function. Multiplying (2.45) by  $w_\alpha^k \eta^2$  for  $k > 1$  and integrating by parts, we obtain

$$\int_M \nabla_{\hat{g}} w_\alpha \nabla_{\hat{g}} (w_\alpha^k \eta^2) dv_{\hat{g}} \leq \xi_\alpha \int_{\partial M} w_\alpha^{q-1+k} \eta^2 ds_{\hat{g}}.$$

Here and in the following,  $C$  denotes some constant independent of  $\alpha$ . As in (2.7), we have

$$\begin{aligned} & \int_M \nabla_{\hat{g}} w_\alpha \nabla_{\hat{g}} (w_\alpha^k \eta^2) dv_{\hat{g}} \\ &= \frac{4k}{(k+1)^2} \int_M |\nabla_{\hat{g}}(w_\alpha^{(k+1)/2} \eta)|^2 dv_{\hat{g}} + \frac{k-1}{(k+1)^2} \int_M w_\alpha^{k+1} \Delta_{\hat{g}}(\eta^2) dv_{\hat{g}} \\ & \quad - \frac{4k}{(k+1)^2} \int_M w_\alpha^{k+1} |\nabla_{\hat{g}} \eta|^2 dv_{\hat{g}} - \frac{k-1}{(k+1)^2} \int_{\partial M} w_\alpha^{k+1} \frac{\partial_{\hat{g}}(\eta^2)}{\partial \nu} ds_{\hat{g}}. \end{aligned}$$

We deduce from the last two formulae that

$$\begin{aligned} (2.46) \quad & \int_M |\nabla_{\hat{g}}(w_\alpha^{(k+1)/2} \eta)|^2 dv_{\hat{g}} \\ & \leq -\frac{k-1}{4k} \int_M w_\alpha^{k+1} \Delta_{\hat{g}}(\eta^2) dv_{\hat{g}} + \int_M w_\alpha^{k+1} |\nabla_{\hat{g}} \eta|^2 dv_{\hat{g}} \\ & \quad + \frac{k-1}{4k} \int_{\partial M} w_\alpha^{k+1} \frac{\partial_{\hat{g}}(\eta^2)}{\partial \nu} ds_{\hat{g}} + \frac{\xi_\alpha (k+1)^2}{4k} \int_{\partial M} w_\alpha^{q-1+k} \eta^2 ds_{\hat{g}}. \end{aligned}$$

We still need the following lemma to start the Moser iteration process:

**LEMMA 2.6** *There exists some  $0 < \delta_0 \ll 1$ ,  $s_0 > q$ , and  $C > 1$  independent of  $\alpha$  such that*

$$(2.47) \quad \int_{\partial M \setminus B_{\mu_\alpha/\delta_0}(x_\alpha)} w_\alpha^{s_0} ds_{\hat{g}} \leq C.$$

**PROOF:** For all  $\varepsilon > 0$ , it follows from Proposition 2.1 and Proposition 2.3 that there exists  $0 < \delta_0 = \delta_0(\varepsilon) < 1$  such that

$$\int_{\partial M \setminus B_{\mu_\alpha/\delta_0}(x_\alpha)} w_\alpha^q ds_{\hat{g}} = \int_{\partial M \setminus B_{\mu_\alpha/\delta_0}(x_\alpha)} u_\alpha^q ds_g < \varepsilon.$$

Since  $\hat{g}^{ij} \sim \mu_\alpha^2 \delta^{ij}$  in  $B_{2\mu_\alpha/\delta_0}(x_\alpha) \setminus B_{\mu_\alpha/(4\delta_0)}(x_\alpha)$ , we can choose  $\eta$  to be some cutoff function satisfying

$$\begin{cases} \eta(x) = 1, \quad d(x_\alpha, x) \geq \mu_\alpha/\delta_0; \\ |\nabla_{\hat{g}} \eta| + |\nabla_{\hat{g}}^2 \eta| \leq C. \end{cases} \quad \eta(x) = 0, \quad d(x_\alpha, x) \leq \mu_\alpha/(2\delta_0)$$

We also take some  $1 < k \leq q - 1$ . It follows from (2.46) and Theorem A.1 in Appendix A that

$$\begin{aligned}
& \int_M |\nabla_{\hat{g}}(w_{\alpha}^{(k+1)/2}\eta)|^2 dv_{\hat{g}} \\
& \leq C(k, \delta_0) + \frac{\xi_{\alpha}(k+1)^2}{4k} \int_{\partial M} w_{\alpha}^{q-1+k} \eta^2 ds_{\hat{g}} \\
& \leq C(k, \delta_0) + \frac{\xi_{\alpha}(k+1)^2}{4k} \left( \int_{\partial M} (w_{\alpha}^{(k+1)/2}\eta)^q ds_{\hat{g}} \right)^{2/q} \left( \int_{\partial M} w_{\alpha}^q ds_{\hat{g}} \right)^{(q-2)/q} \\
& \leq C(k, \delta_0) + C\varepsilon^{(q-2)/q} \int_M |\nabla_{\hat{g}}(w_{\alpha}^{(k+1)/2}\eta)|^2 dv_{\hat{g}}.
\end{aligned}$$

Taking  $\varepsilon > 0$  small, we have

$$\int_M |\nabla_{\hat{g}}(w_{\alpha}^{(k+1)/2}\eta)|^2 dv_{\hat{g}} \leq C.$$

It follows from Theorem A.1 in Appendix A that

$$\int_{\partial M} (w_{\alpha}^{(k+1)/2}\eta)^q ds_{\hat{g}} \leq \left( \int_M |\nabla_{\hat{g}}(w_{\alpha}^{(k+1)/2}\eta)|^2 dv_{\hat{g}} \right)^{q/2} \leq C.$$

Lemma 2.6 is established. ■

**REMARK 2.7** Without loss of generality, we can assume that  $\delta_0$  in Lemma 2.6 is small enough so that  $B_{\mu_{\alpha}/\delta_0}(x_{\alpha}) \subset B_{4\mu_{\alpha}/\delta_0}(P_{\alpha})$ .

Set, for  $\delta = \delta_0/10$ ,

$$R_i = \mu_{\alpha} \frac{\left(2 - \frac{1}{2^{i-1}}\right)}{\delta}, \quad i = 1, 2, 3, \dots$$

Clearly

$$(2.48) \quad B_{\mu_{\alpha}/\delta_0}(x_{\alpha}) \subset B_{R_i}(P_{\alpha}) \quad \forall i.$$

Recall that for  $\mu_{\alpha}/\delta < |y| \leq 2\mu_{\alpha}/\delta$ ,

$$(2.49) \quad \frac{\mu_\alpha^{(2-n)/2}}{C} \leq \varphi_\alpha(y) \leq C\mu_\alpha^{(2-n)/2}, \quad C^{-1}\mu_\alpha^{-2}g \leq \hat{g} \leq C\mu_\alpha^{-2}g.$$

We can choose some smooth cutoff function  $\eta_i$  satisfying

$$\begin{cases} \eta_i(y) = 1, \quad |y| > R_{i+1}; \\ \eta_i(y) = 0, \quad |y| < R_i \\ |\nabla_{\hat{g}}\eta_i| \leq C2^i, \quad |\nabla_{\hat{g}}^2\eta_i| \leq C4^i. \end{cases}$$

Taking  $\eta = \eta_i$  in (2.46), we have

$$\begin{aligned} (2.50) \quad & \int_M |\nabla_{\hat{g}}(w_\alpha^{(k+1)/2}\eta_i)|^2 dv_{\hat{g}} \\ & \leq C4^i \int_{M \setminus B_{R_i}(P_\alpha)} w_\alpha^{k+1} dv_{\hat{g}} + C2^i \int_{\partial M \setminus B_{R_i}(P_\alpha)} w_\alpha^{k+1} ds_{\hat{g}} \\ & \quad + \frac{C(k+1)^2}{k} \int_{\partial M \setminus B_{R_i}(P_\alpha)} w_\alpha^{q-1+k} ds_{\hat{g}}. \end{aligned}$$

It follows from (2.48), (2.49), and Theorem A.1 in Appendix A that

$$\begin{aligned} (2.51) \quad & \left[ \int_{M \setminus B_{R_i}(P_\alpha)} (w_\alpha^{(k+1)/2}\eta_i)^p dv_{\hat{g}} \right]^{2/p} \\ & \leq C \int_{M \setminus B_{R_i}(P_\alpha)} |\nabla_{\hat{g}}(w_\alpha^{(k+1)/2}\eta_i)|^2 dv_{\hat{g}}, \end{aligned}$$

$$\begin{aligned} (2.52) \quad & \left[ \int_{\partial M \setminus B_{R_i}(P_\alpha)} (w_\alpha^{(k+1)/2}\eta_i)^q ds_{\hat{g}} \right]^{2/q} \\ & \leq C \int_{M \setminus B_{R_i}(P_\alpha)} |\nabla_{\hat{g}}(w_\alpha^{(k+1)/2}\eta_i)|^2 dv_{\hat{g}}. \end{aligned}$$

Using (2.50), we can derive from (2.51) and (2.52) that

$$\begin{aligned} (2.53) \quad & \left[ \int_{M \setminus B_{R_{i+1}}(P_\alpha)} w_\alpha^{(k+1)p/2} dv_{\hat{g}} \right]^{2/p} \\ & \leq C4^i \int_{M \setminus B_{R_i}(P_\alpha)} w_\alpha^{k+1} dv_{\hat{g}} + C2^i \int_{\partial M \setminus B_{R_i}(P_\alpha)} w_\alpha^{k+1} ds_{\hat{g}} \\ & \quad + \frac{C(k+1)^2}{k} \int_{\partial M \setminus B_{R_i}(P_\alpha)} w_\alpha^{q-1+k} ds_{\hat{g}}, \end{aligned}$$

and

$$(2.54) \quad \begin{aligned} & \left[ \int_{\partial M \setminus B_{R_{i+1}}(P_\alpha)} w_\alpha^{(k+1)q/2} ds_{\hat{g}} \right]^{2/q} \\ & \leq C 4^i \int_{M \setminus B_{R_i}(P_\alpha)} w_\alpha^{k+1} dv_{\hat{g}} + C 2^i \int_{\partial M \setminus B_{R_i}(P_\alpha)} w_\alpha^{k+1} ds_{\hat{g}} \\ & + \frac{C(k+1)^2}{k} \int_{\partial M \setminus B_{R_i}(P_\alpha)} w_\alpha^{q-1+k} ds_{\hat{g}}. \end{aligned}$$

Set  $r_0 = s_0/(q-2)$  where  $s_0$  is given in Lemma 2.6. It follows from (2.47) and Hölder's inequality that

$$(2.55) \quad \begin{aligned} & \int_{\partial M \setminus B_{R_i}(P_\alpha)} w_\alpha^{q-1+k} ds_{\hat{g}} \\ & = \int_{\partial M \setminus B_{R_i}(P_\alpha)} w_\alpha^{q-2} w_\alpha^{k+1} ds_{\hat{g}} \\ & \leq C \left( \int_{\partial M \setminus B_{R_i}(P_\alpha)} w_\alpha^{(k+1)r_0/(r_0-1)} ds_{\hat{g}} \right)^{(r_0-1)/r_0}. \end{aligned}$$

It follows from (2.53), (2.54), and (2.55) that

$$(2.56) \quad \begin{aligned} & \left[ \int_{M \setminus B_{R_{i+1}}(P_\alpha)} w_\alpha^{(k+1)p/2} dv_{\hat{g}} \right]^{2/p} + \left[ \int_{\partial M \setminus B_{R_{i+1}}(P_\alpha)} w_\alpha^{(k+1)q/2} ds_{\hat{g}} \right]^{2/q} \\ & \leq C 4^i \int_{M \setminus B_{R_i}(P_\alpha)} w_\alpha^{k+1} dv_{\hat{g}} + C 2^i \int_{\partial M \setminus B_{R_i}(P_\alpha)} w_\alpha^{k+1} ds_{\hat{g}} \\ & + \frac{C(k+1)^2}{k} \left[ \int_{\partial M \setminus B_{R_i}(P_\alpha)} w_\alpha^{(k+1)r_0/(r_0-1)} ds_{\hat{g}} \right]^{(r_0-1)/r_0}. \end{aligned}$$

By setting  $\beta = q(r_0-1)/(2r_0)$ , it is easy to see from  $s_0 > q$  that  $\beta > 1$ . Since we can take  $s_0$  close to  $q$  from the beginning, we can assume without loss of generality that  $\beta \leq p/2$ . It follows from Hölder's inequality, (2.48), and (2.49) that

$$(2.57) \quad \begin{aligned} & \left[ \int_{M \setminus B_{R_{i+1}}(P_\alpha)} w_\alpha^{(k+1)\beta} dv_{\hat{g}} \right]^{1/\beta} \\ & \leq C \left[ \int_{M \setminus B_{R_{i+1}}(P_\alpha)} w_\alpha^{(k+1)p/2} dv_{\hat{g}} \right]^{2/p} \end{aligned}$$

and

$$(2.58) \quad \int_{\partial M \setminus B_{R_i}(P_\alpha)} w_\alpha^{k+1} ds_{\hat{g}} \leq \left[ \int_{\partial M \setminus B_{R_i}(P_\alpha)} w_\alpha^{(k+1)r_0/(r_0-1)} ds_{\hat{g}} \right]^{(r_0-1)/r_0}.$$

Set  $q_0 = 2r_0/(r_0-1) < q$ ,  $q_i = \beta q_{i-1} = \beta^{i-1}q$ , and  $p_i = q_i(r_0-1)/r_0 = 2\beta^i$ , where  $i \geq 1$ . Taking  $k = p_i - 1$  ( $i \geq 1$ ) in (2.56) and using (2.57) and (2.58), we obtain

$$\begin{aligned} & \|w_\alpha\|_{p_{i+1}, M \setminus B_{R_{i+1}}(P_\alpha)}^{p_i} + \|w_\alpha\|_{q_{i+1}, \partial M \setminus B_{R_{i+1}}(P_\alpha)}^{p_i} \\ & \leq \left( C4^i + \frac{Cp_i^2}{(p_i-1)} \right) \left( \|w_\alpha\|_{p_i, M \setminus B_{R_i}(P_\alpha)}^{p_i} + \|w_\alpha\|_{q_i, \partial M \setminus B_{R_i}(P_\alpha)}^{p_i} \right). \end{aligned}$$

Since  $\beta > 1$ , we have  $a^\beta + b^\beta \leq (a+b)^\beta$  for all  $a, b \geq 0$ . It follows that

$$\begin{aligned} & \left( \|w_\alpha\|_{p_{i+1}, M \setminus B_{R_{i+1}}(P_\alpha)}^{p_{i+1}} + \|w_\alpha\|_{q_{i+1}, \partial M \setminus B_{R_{i+1}}(P_\alpha)}^{p_{i+1}} \right)^{1/p_{i+1}} \\ & \leq \left( \|w_\alpha\|_{p_{i+1}, M \setminus B_{R_{i+1}}(P_\alpha)}^{p_i} + \|w_\alpha\|_{q_{i+1}, \partial M \setminus B_{R_{i+1}}(P_\alpha)}^{p_i} \right)^{1/p_i} \\ (2.59) \quad & \leq \left( C4^i + \frac{Cp_i^2}{p_i-1} \right)^{1/p_i} \left( \|w_\alpha\|_{p_i, M \setminus B_{R_i}(P_\alpha)}^{p_i} \right. \\ & \quad \left. + \|w_\alpha\|_{q_i, \partial M \setminus B_{R_i}(P_\alpha)}^{p_i} \right)^{1/p_i}. \end{aligned}$$

It is easy to see that

$$\left( C4^i + \frac{Cp_i^2}{p_i-1} \right)^{1/p_i} \leq [C(4^i + 2\beta^i)]^{1/(2\beta^i)} \leq C^{1/(2\beta^i)}(4 + \beta)^{i/(2\beta^i)}.$$

Thus

$$\prod_{i=1}^{\infty} \left( C4^i + \frac{Cp_i^2}{p_i-1} \right)^{1/p_i} \leq C < \infty.$$

It follows that

$$\begin{aligned} \|w_\alpha\|_{p_{i+1}, M \setminus B_{R_{i+1}}(P_\alpha)} & \leq C \left( \|w_\alpha\|_{p_1, M \setminus B_{R_1}(P_\alpha)}^{2\beta} + \|w_\alpha\|_{q_1, \partial M \setminus B_{R_1}(P_\alpha)}^{2\beta} \right)^{1/(2\beta)} \\ & \leq C. \end{aligned}$$

Sending  $i$  to  $\infty$ , we have

$$(2.60) \quad \|w_\alpha\|_{L^\infty(M \setminus B_{2\mu_\alpha/\delta}(P_\alpha))} \leq C(\delta).$$

It is easy to see that inside  $B_{2\mu_\alpha/\delta}(P_\alpha)$ ,  $|y| \leq C\mu_\alpha$ . Therefore, it follows from (2.41) that  $\forall y \in B_{2\mu_\alpha/\delta}(P_\alpha) : \varphi_\alpha(y) \geq C^{-1}\mu_\alpha^{-(n-2)/2}$ . It follows that for all  $y \in B_{2\mu_\alpha/\delta}(P_\alpha)$ ,

$$(2.61) \quad w_\alpha = \frac{u_\alpha}{\varphi_\alpha} \leq C\mu_\alpha^{(n-2)/2}u_\alpha = Cu_\alpha/u_\alpha(x_\alpha) \leq C.$$

Proposition 2.4 follows from (2.60) and (2.61).  $\blacksquare$

### 3 Balance Checking via Pohozaev Identity

In this section, we derive a contradiction by using the Pohozaev identity to do a balance checking in a ball centered at  $x_\alpha$  of radius  $1/\alpha$ . The upper bound obtained in Section 2 plays a crucial role. For  $n = 3$ , it is subtler since we need to obtain an appropriate lower bound of  $u_\alpha$  in order to reach a contradiction. This lower bound is obtained in this section by use of the maximum principle.

By choosing an appropriate coordinate system centered at  $x_\alpha$ , we can assume without loss of generality that  $x_\alpha = 0$ ,  $g_{ij}(0) = \delta_{ij}$ ,  $B_1^+(0) \subset M$ , and  $\{(x', 0) : |x'| < 1\} \subset \partial M$ .

Let  $R_\alpha = 1/(\alpha\mu_\alpha)$ ,  $h_\alpha = g_{ij}(\mu_\alpha x) dx^i dx^j$  in  $B_{10R_\alpha}^+(0)$ , and

$$\bar{v}_\alpha(x) = \mu_\alpha^{(n-2)/2}u_\alpha(\mu_\alpha x) \quad \text{for } x \in B_{10R_\alpha}^+(0).$$

It follows from (2.22) and (2.2) that  $R_\alpha \rightarrow \infty$  as  $\alpha \rightarrow \infty$ , and  $\bar{v}_\alpha$  satisfies

$$(3.1) \quad \begin{cases} \Delta_{h_\alpha} \bar{v}_\alpha = 0 & \text{in } B_{10R_\alpha}^+(0) \\ \frac{\partial_{h_\alpha} \bar{v}_\alpha}{\partial \nu} = \xi_\alpha \bar{v}_\alpha^{q-1} - \alpha \mu_\alpha \bar{v}_\alpha & \text{on } \{(x', 0) : |x'| < 10R_\alpha\} \\ \bar{v}_\alpha(0) = 1, \quad 0 < \bar{v}_\alpha \leq 1. & \end{cases}$$

Clearly

$$(3.2) \quad |h_\alpha^{ij}(x) - \delta^{ij}| \leq C|\mu_\alpha x|, \quad |\Gamma_{ij}^k(x)| \leq C\mu_\alpha \quad \text{in } B_{10R_\alpha}^+(0),$$

where  $\Gamma_{ij}^k$  is the Christoffel symbol of  $h_\alpha$  and  $C$  is, as always, some constant independent of  $\alpha$ .

As explained in Section 2,

$$(3.3) \quad \lim_{\alpha \rightarrow \infty} \|\bar{v}_\alpha - v\|_{C^3(\overline{B_R^+(0)})} = 0 \quad \forall R > 0,$$

where  $v$  is the function defined in  $\mathbb{R}_+^n$  given in (2.25). It is not difficult to see from Proposition 2.4 that

$$(3.4) \quad \bar{v}_\alpha(x) \leq \frac{C}{1+|x|^{n-2}} \quad \text{for } x \in \bar{B}_{10R_\alpha}^+(0).$$

We need some further estimates on  $\bar{v}_\alpha$ .

**PROPOSITION 3.1** *For all  $\alpha \geq 1$ ,  $x \in B_{R_\alpha}^+(0)$ , we have*

$$|\nabla \bar{v}_\alpha(x)| \leq \frac{C}{1+|x|^{n-1}}, \quad |\nabla^2 \bar{v}_\alpha(x)| \leq \frac{C}{1+|x|^n},$$

where  $|\nabla^2 \bar{v}_\alpha| = \sum_{i,j=1}^n |\partial^2 \bar{v}_\alpha / \partial x^i \partial x^j|$  and  $C$  is some constant independent of  $\alpha$  and  $x$ .

**PROOF:** It follows from (3.3) that

$$|\nabla \bar{v}_\alpha(x)| < C, \quad |\nabla^2 \bar{v}_\alpha(x)| \leq C \quad \text{in } B_1^+(0).$$

So we only need to show Proposition 3.1 for  $|x| > 1$ . For all  $x_0 \in B_{R_\alpha}^+(0) \setminus B_1^+(0)$ , set  $R = |x_0|$ ,  $\tilde{u}(x) = R^{n-2} \bar{v}_\alpha(Rx)$ , and  $\tilde{g}_{ij}(x) = (h_\alpha)_{ij}(Rx)$ . It follows from (3.1) that

$$\begin{cases} \Delta_{\tilde{g}} \tilde{u} = 0 & \text{in } B_5^+ \setminus B_{1/5}^+ \\ \frac{\partial_{\tilde{g}} \tilde{u}}{\partial \nu} = R^{n-1} (\xi_\alpha \bar{v}_\alpha^{q-1}(Rx) - \alpha \mu_\alpha \bar{v}_\alpha(Rx)) & \text{on } \{(x', 0) : \frac{1}{5} < |x'| < 5\}. \end{cases}$$

On  $\{(x', 0) : \frac{1}{5} < |x'| < 5\}$ , we derive from (3.4) that

$$\left| \frac{\partial_{\tilde{g}} \tilde{u}}{\partial \nu} \right| = \left| R^{n-1} (\xi_\alpha \bar{v}_\alpha^{q-1}(Rx) - \alpha \mu_\alpha \bar{v}_\alpha(Rx)) \right| \leq CR^{-1} + C\alpha\mu_\alpha R \leq C.$$

It follows from standard elliptic estimates that for some  $0 < \beta < 1$ ,

$$(3.5) \quad \|\tilde{u}\|_{C^\beta(\overline{B_4^+ \setminus B_{1/4}^+})} \leq C.$$

Rewriting the boundary condition of  $\tilde{u}$  as

$$\frac{\partial_{\tilde{g}} \tilde{u}}{\partial \nu} = \xi_\alpha R^{-1} \tilde{u}^{q-1} - \alpha \mu_\alpha R \tilde{u}$$

and noticing (see (3.5))

$$\|\xi_\alpha R^{-1} \tilde{u}^{q-1} - \alpha \mu_\alpha R \tilde{u}\|_{C^\beta(\overline{B_4^+ \setminus B_{1/4}^+})} \leq C,$$

we have, by standard elliptic estimates, that

$$(3.6) \quad \|\nabla_{\tilde{g}} \tilde{u}\|_{C^\beta(\overline{B_3^+ \setminus B_{1/3}^+})} \leq C.$$

Therefore

$$|\nabla_{h_\alpha} \bar{v}_\alpha(x_0)| = R^{1-n} \left| \nabla_{\tilde{g}} \tilde{u} \left( \frac{x_0}{R} \right) \right| \leq C |x_0|^{1-n} \leq \frac{C}{(1 + |x_0|^{n-1})},$$

which gives us the gradient estimate.

Also, from (3.5) and (3.6), we know for some  $0 < \beta' < \beta$ ,

$$\|\xi_\alpha R^{-1} \tilde{u}^{q-1} - \alpha \mu_\alpha R \tilde{u}\|_{C^{1,\beta'}(\overline{B_3^+ \setminus B_{1/3}^+})} \leq C.$$

Thus by standard elliptic estimates

$$\|\nabla_{\tilde{g}}^2 \tilde{u}\|_{C^{\beta'}(\overline{B_2^+ \setminus B_{1/2}^+})} \leq C,$$

which gives us

$$|\nabla_{h_\alpha}^2 \bar{v}_\alpha(x_0)| = R^{-n} \left| \nabla_{\tilde{g}}^2 \tilde{u} \left( \frac{x_0}{R} \right) \right| \leq C |x_0|^{-n} \leq \frac{C}{(1 + |x_0|^n)}.$$

We have, in view of (3.2), established Proposition 3.1. ■

For  $n = 3$ , we need to obtain an appropriate lower bound of  $\bar{v}_\alpha$ . Clearly one can also obtain lower bounds for  $n \geq 4$  by the same method, but since we do not need it for the application in this paper, we omit it.

**PROPOSITION 3.2** *For  $n = 3$  and  $\alpha$  large enough,*

$$\bar{v}_\alpha(x) \geq \frac{1}{C(1 + |x|)} \quad \forall x \in \overline{B_{R_\alpha^{1/4}}^+(0)},$$

where  $C > 0$  is some constant independent of  $\alpha$ .

PROOF: In view of (3.3), we only need to prove the above estimate for  $|x| > 20$ . In the following,  $\alpha$  is always assumed to be suitably large. Let  $\bar{x} = (0, \dots, 0, 1)$  and

$$G_\alpha(x) = \frac{1}{|x - \bar{x}|} - \frac{1}{R_\alpha^{1/2}|x - \bar{x}|^{1/2}} \quad \text{in } B_{R_\alpha^{1/3}}(\bar{x}) \setminus B_2(\bar{x}).$$

It is easy to see that

$$\frac{1}{2|x - \bar{x}|} \leq G_\alpha(x) \leq \frac{2}{|x - \bar{x}|} \quad \text{in } B_{R_\alpha^{1/3}}(\bar{x}) \setminus B_2(\bar{x}).$$

From (3.2) we have that in  $B_{R_\alpha^{1/3}}^+(\bar{x}) \setminus B_2(\bar{x})$ ,

$$\begin{aligned} \Delta_{h_\alpha} \left( -\frac{1}{R_\alpha^{1/2}|x - \bar{x}|^{1/2}} \right) &\geq \frac{1}{CR_\alpha^{1/2}|x - \bar{x}|^{5/2}}, \\ \left| \Delta_{h_\alpha} \left( \frac{1}{|x - \bar{x}|} \right) \right| &\leq \frac{C\mu_\alpha}{|x - \bar{x}|^2}, \end{aligned}$$

where  $B_{R_\alpha^{1/3}}^+(\bar{x}) = \{x \in \mathbb{R}_+^n : |x - \bar{x}| < R_\alpha^{1/3}\}$ . It follows that  $\Delta_{h_\alpha} G_\alpha \geq 0$ .

Also, it follows from (3.2) that for all  $x = (x', 0)$ ,  $1 < |x'| < R_\alpha^{1/3}$ ,

$$\begin{aligned} \frac{\partial_{h_\alpha}}{\partial \nu} \left( \frac{1}{|x - \bar{x}|} \right) &\leq -\frac{1}{C|x - \bar{x}|^3}, \\ \left| \frac{\partial_{h_\alpha}}{\partial \nu} \left( \frac{1}{R_\alpha^{1/2}|x - \bar{x}|^{1/2}} \right) \right| &\leq \frac{C}{R_\alpha^{1/4}|x - \bar{x}|^3}. \end{aligned}$$

Therefore, using (3.4), we have, for all  $x = (x', 0)$ ,  $1 < |x'| < R_\alpha^{1/3}$ ,

$$-\alpha\mu_\alpha \bar{v}_\alpha - \frac{\partial_{h_\alpha}}{\partial \nu}(G_\alpha) \geq -\frac{C}{R_\alpha(1+|x|)} + \frac{1}{C|x - \bar{x}|^3} > 0.$$

We will use the maximum principle on  $A = \{x \in \mathbb{R}_+^n : 10 < |x - \bar{x}| < R_\alpha^{1/3}\}$ . Let  $\Sigma_1 = \partial A \cap \{x_n = 0\}$ ,  $\Sigma_2 = \partial A \cap \{|x - \bar{x}| = 10\}$ , and  $\Sigma_3 = \partial A \cap \{|x - \bar{x}| = R_\alpha^{1/3}\}$ . Choose  $0 < \tau < 1$  small enough such that  $\tau G_\alpha \leq \bar{v}_\alpha$  on  $\Sigma_2$ . Define  $H_\alpha = \tau G_\alpha - \max_{\Sigma_3}(\tau G_\alpha)$ ; then

$$\begin{cases} \Delta_{h_\alpha}(\bar{v}_\alpha - H_\alpha) \leq 0 & \text{in } A \\ \bar{v}_\alpha - H_\alpha \geq 0 & \text{on } \Sigma_2 \cup \Sigma_3 \\ \frac{\partial_{h_\alpha}(\bar{v}_\alpha - H_\alpha)}{\partial \nu} > 0, & \text{on } \Sigma_1. \end{cases}$$

It follows from the maximum principle that

$$\bar{v}_\alpha \geq H_\alpha \quad \text{in } A.$$

Consequently, for all  $x \in B_{R_\alpha^{1/4}}^+(0) \setminus B_{10}^+(\bar{x})$ ,

$$\bar{v}_\alpha(x) \geq H_\alpha(x) \geq \frac{C\tau}{|x - \bar{x}|} - \frac{C\tau}{R_\alpha^{1/3}} \geq \frac{C\tau}{2|x - \bar{x}|}.$$

Proposition 3.2 is established.  $\blacksquare$

For convenience, throughout the rest of this section we set  $\Gamma_1 = \partial B_{R_\alpha}^+(0) \cap \{(x', 0) : x' \in \mathbb{R}^{n-1}\}$  and  $\Gamma_2 = \partial B_{R_\alpha}^+(0) \cap \{(x', x_n) : x_n > 0\}$ . We always use  $dV$  for the volume element of the standard Euclidean metric,  $dS$  for the surface element of the standard Euclidean metric,  $\nu$  for the unit outer normal vector of the corresponding surface with respect to the specified metrics, and  $\cdot$  for the inner product under the standard Euclidean metric. The balance checking via the Pohozaev identity will be performed in  $B_{R_\alpha}^+(0)$ .

The following identity can easily be verified (see [22]):

$$2\Delta\bar{v}_\alpha(\nabla\bar{v}_\alpha \cdot x) = \operatorname{div}[2(\nabla\bar{v}_\alpha \cdot x)\nabla\bar{v}_\alpha - |\nabla\bar{v}_\alpha|^2 x] + (n-2)|\nabla\bar{v}_\alpha|^2.$$

It follows that

$$\begin{aligned} (3.7) \quad & \int_{B_{R_\alpha}^+} \Delta\bar{v}_\alpha(\nabla\bar{v}_\alpha \cdot x) dV - \frac{n-2}{2} \int_{B_{R_\alpha}^+} |\nabla\bar{v}_\alpha|^2 dV \\ &= \frac{1}{2} \int_{B_{R_\alpha}^+} \operatorname{div}[2(\nabla\bar{v}_\alpha \cdot x)\nabla\bar{v}_\alpha - |\nabla\bar{v}_\alpha|^2 x] dV. \end{aligned}$$

Integrating by parts, we have

$$\begin{aligned} & \frac{1}{2} \int_{B_{R_\alpha}^+} \operatorname{div}[2(\nabla\bar{v}_\alpha \cdot x)\nabla\bar{v}_\alpha - |\nabla\bar{v}_\alpha|^2 x] dV \\ &= \frac{1}{2} \int_{\partial B_{R_\alpha}^+} [2(\nabla\bar{v}_\alpha \cdot x)(\nabla\bar{v}_\alpha \cdot \nu) - |\nabla\bar{v}_\alpha|^2(x \cdot \nu)] dS. \end{aligned}$$

It is easy to check that

$$(\nabla\bar{v}_\alpha \cdot x)(\nabla\bar{v}_\alpha \cdot \nu) = \begin{cases} \left(\frac{\partial\bar{v}_\alpha}{\partial\nu}\right)^2 |x| & \text{on } \Gamma_2 \\ \left(\sum_{i=1}^{n-1} x_i \frac{\partial\bar{v}_\alpha}{\partial x_i}\right) \frac{\partial\bar{v}_\alpha}{\partial\nu} & \text{on } \Gamma_1, \end{cases}$$

and

$$x \cdot \nu = \begin{cases} |x| & \text{on } \Gamma_2 \\ 0 & \text{on } \Gamma_1. \end{cases}$$

Therefore

$$\begin{aligned} (3.8) \quad & \frac{1}{2} \int_{B_{R_\alpha}^+} \operatorname{div}[2(\nabla \bar{v}_\alpha \cdot x) \nabla \bar{v}_\alpha - |\nabla \bar{v}_\alpha|^2 x] dV \\ &= \int_{\Gamma_1} \left( \sum_{i=1}^{n-1} x_i \frac{\partial \bar{v}_\alpha}{\partial x_i} \right) \frac{\partial \bar{v}_\alpha}{\partial \nu} + \int_{\Gamma_2} \left[ |x| \left( \frac{\partial \bar{v}_\alpha}{\partial \nu} \right)^2 - \frac{|x|}{2} |\nabla \bar{v}_\alpha|^2 \right] dS \\ &= \int_{\Gamma_1} \left( \sum_{i=1}^{n-1} x_i \frac{\partial \bar{v}_\alpha}{\partial x_i} \right) \frac{\partial \bar{v}_\alpha}{\partial \nu} + \int_{\Gamma_2} \frac{|x|}{2} \left[ \left( \frac{\partial \bar{v}_\alpha}{\partial \nu} \right)^2 - |\partial_{\tan} \bar{v}_\alpha|^2 \right] dS, \end{aligned}$$

where  $\partial_{\tan}$  denotes the tangential differentiation on  $\Gamma_2$ .

On the other hand,

$$\int_{B_{R_\alpha}^+} |\nabla \bar{v}_\alpha|^2 = - \int_{B_{R_\alpha}^+} \bar{v}_\alpha \Delta \bar{v}_\alpha + \int_{\partial B_{R_\alpha}^+} \bar{v}_\alpha \frac{\partial \bar{v}_\alpha}{\partial \nu}.$$

Therefore,

$$\begin{aligned} & \int_{B_{R_\alpha}^+} \Delta \bar{v}_\alpha (\nabla \bar{v}_\alpha \cdot x) dV - \frac{n-2}{2} \int_{B_{R_\alpha}^+} |\nabla \bar{v}_\alpha|^2 dV \\ &= \int_{B_{R_\alpha}^+} \Delta \bar{v}_\alpha (\nabla \bar{v}_\alpha \cdot x) dV + \frac{n-2}{2} \int_{B_{R_\alpha}^+} \Delta \bar{v}_\alpha \bar{v}_\alpha dV \\ & \quad - \frac{n-2}{2} \int_{\partial B_{R_\alpha}^+} \bar{v}_\alpha \frac{\partial \bar{v}_\alpha}{\partial \nu}. \end{aligned}$$

Combining (3.7), (3.8), and the above, we have

$$\begin{aligned} (3.9) \quad & \int_{B_{R_\alpha}^+} \Delta \bar{v}_\alpha (\nabla \bar{v}_\alpha \cdot x) dV + \frac{n-2}{2} \int_{B_{R_\alpha}^+} \bar{v}_\alpha \Delta \bar{v}_\alpha dV \\ &= J(R_\alpha, \bar{v}_\alpha) + I(R_\alpha, \bar{v}_\alpha), \end{aligned}$$

where

$$\begin{aligned} J(R_\alpha, \bar{v}_\alpha) &= \frac{1}{2} \int_{\Gamma_2} \left\{ \left| \frac{\partial \bar{v}_\alpha}{\partial \nu} \right|^2 |x| - |\partial_{\tan} \bar{v}_\alpha|^2 |x| + (n-2) \frac{\partial \bar{v}_\alpha}{\partial \nu} \bar{v}_\alpha \right\} dS, \\ I(R_\alpha, \bar{v}_\alpha) &= \frac{1}{2} \int_{\Gamma_1} \left\{ 2 \left( \sum_{i=1}^{n-1} x_i \frac{\partial \bar{v}_\alpha}{\partial x_i} \right) \frac{\partial \bar{v}_\alpha}{\partial \nu} + (n-2) \frac{\partial \bar{v}_\alpha}{\partial \nu} \bar{v}_\alpha \right\} dS. \end{aligned}$$

Replacing  $\Delta \bar{v}_\alpha$  in (3.9) by

$$\Delta \bar{v}_\alpha = \Delta_{h_\alpha} \bar{v}_\alpha - (h_\alpha^{ij} - \delta^{ij}) \partial_{ij} \bar{v}_\alpha + h_\alpha^{ij} \Gamma_{ij}^k \partial_k \bar{v}_\alpha,$$

we have

$$\begin{aligned} & - \int_{B_{R_\alpha}^+} (x^i \partial_i \bar{v}_\alpha) \Delta_{h_\alpha} \bar{v}_\alpha dV - \frac{n-2}{2} \int_{B_{R_\alpha}^+} \bar{v}_\alpha \Delta_{h_\alpha} \bar{v}_\alpha dV \\ & + \int_{B_{R_\alpha}^+} (x^k \partial_k \bar{v}_\alpha) (h_\alpha^{ij} - \delta^{ij}) \partial_{ij} \bar{v}_\alpha dV - \int_{B_{R_\alpha}^+} (x^l \partial_l \bar{v}_\alpha) (h_\alpha^{ij} \Gamma_{ij}^k \partial_k \bar{v}_\alpha) dV \\ & + \frac{n-2}{2} \int_{B_{R_\alpha}^+} \bar{v}_\alpha (h_\alpha^{ij} - \delta^{ij}) \partial_{ij} \bar{v}_\alpha dV - \frac{n-2}{2} \int_{B_{R_\alpha}^+} \bar{v}_\alpha (h_\alpha^{ij} \Gamma_{ij}^k) \partial_k \bar{v}_\alpha dV \\ & = -J(R_\alpha, \bar{v}_\alpha) - I(R_\alpha, \bar{v}_\alpha), \end{aligned}$$

where  $x^i \partial_i \bar{v}_\alpha = \sum_{i=1}^n x_i \partial_i \bar{v}_\alpha$ , and so on, here and in the discussion below. So far we have not used the equation of  $\bar{v}_\alpha$ . Now we use equation (3.1) satisfied by  $\bar{v}_\alpha$  and obtain

$$(3.10) \quad A(h_\alpha, \bar{v}_\alpha) = -J(R_\alpha, \bar{v}_\alpha) - I(R_\alpha, \bar{v}_\alpha)$$

where

$$\begin{aligned} & A(h_\alpha, \bar{v}_\alpha) \\ & = \int_{B_{R_\alpha}^+} (x^k \partial_k \bar{v}_\alpha) (h_\alpha^{ij} - \delta^{ij}) \partial_{ij} \bar{v}_\alpha dV - \int_{B_{R_\alpha}^+} (x^l \partial_l \bar{v}_\alpha) (h_\alpha^{ij} \Gamma_{ij}^k \partial_k \bar{v}_\alpha) dV \\ & + \frac{n-2}{2} \int_{B_{R_\alpha}^+} \bar{v}_\alpha (h_\alpha^{ij} - \delta^{ij}) \partial_{ij} \bar{v}_\alpha dV - \frac{n-2}{2} \int_{B_{R_\alpha}^+} \bar{v}_\alpha (h_\alpha^{ij} \Gamma_{ij}^k) \partial_k \bar{v}_\alpha dV. \end{aligned}$$

Using (3.2), we have

$$\begin{aligned} (3.11) \quad A(h_\alpha, \bar{v}_\alpha) & = O\left( \int_{B_{R_\alpha}^+} \mu_\alpha |x|^2 |\nabla \bar{v}_\alpha| |\nabla^2 \bar{v}_\alpha| dV \right) \\ & + O\left( \int_{B_{R_\alpha}^+} \mu_\alpha |x| |\nabla \bar{v}_\alpha|^2 dV \right) \\ & + O\left( \int_{B_{R_\alpha}^+} \mu_\alpha |x| \bar{v}_\alpha |\nabla^2 \bar{v}_\alpha| dV \right) \\ & + O\left( \int_{B_{R_\alpha}^+} \mu_\alpha \bar{v}_\alpha |\nabla \bar{v}_\alpha| dV \right) \end{aligned}$$

We simplify  $I(R_\alpha, \bar{v}_\alpha)$  by using the equation of  $\bar{v}_\alpha$  (3.1). It is easy to see from (3.2) that

$$\frac{\partial_{h_\alpha} \bar{v}_\alpha}{\partial \nu} = \frac{\partial \bar{v}}{\partial \nu} + O(\mu_\alpha |x'| |\nabla \bar{v}_\alpha|) \quad \text{on } \Gamma_1.$$

It follows that

$$\begin{aligned} (3.12) \quad & 2I(R_\alpha, \bar{v}_\alpha) \\ &= \int_{\Gamma_1} \left\{ 2 \left( \sum_{i=1}^{n-1} x_i \frac{\partial \bar{v}_\alpha}{\partial x_i} \right) \frac{\partial_{h_\alpha} \bar{v}_\alpha}{\partial \nu} + (n-2) \frac{\partial_{h_\alpha} \bar{v}_\alpha}{\partial \nu} \bar{v}_\alpha \right\} dS \\ & \quad + O \left( \int_{\Gamma_1} [\mu_\alpha |x'|^2 |\nabla \bar{v}_\alpha|^2 + \mu_\alpha |x'| \bar{v}_\alpha |\nabla \bar{v}_\alpha|] dS \right). \end{aligned}$$

Using the boundary condition of  $\bar{v}_\alpha$  in (3.1), we have

$$\begin{aligned} & \int_{\Gamma_1} \left\{ 2 \left( \sum_{i=1}^{n-1} x_i \frac{\partial \bar{v}_\alpha}{\partial x_i} \right) \frac{\partial_{h_\alpha} \bar{v}_\alpha}{\partial \nu} + (n-2) \frac{\partial_{h_\alpha} \bar{v}_\alpha}{\partial \nu} \bar{v}_\alpha \right\} dS \\ &= \int_{\Gamma_1} \left\{ 2(\xi_\alpha \bar{v}_\alpha^{q-1} - \alpha \mu_\alpha \bar{v}_\alpha) \left( \sum_{i=1}^{n-1} x_i \frac{\partial \bar{v}_\alpha}{\partial x_i} \right) \right. \\ & \quad \left. + (n-2)(\xi_\alpha \bar{v}_\alpha^{q-1} - \alpha \mu_\alpha \bar{v}_\alpha) \bar{v}_\alpha \right\} dS \\ &= -\frac{2(n-1)}{q} \int_{\Gamma_1} \xi_\alpha \bar{v}_\alpha^q dS + (n-1) \alpha \mu_\alpha \int_{\Gamma_1} \bar{v}_\alpha^2 dS \\ & \quad + \frac{2}{q} \int_{\partial \Gamma_1} \xi_\alpha \bar{v}_\alpha^q |x| dS - \int_{\partial \Gamma_1} \alpha \mu_\alpha \bar{v}_\alpha^2 |x| dS \\ & \quad + (n-2) \int_{\Gamma_1} \xi_\alpha \bar{v}_\alpha^q dS - (n-2) \alpha \mu_\alpha \int_{\Gamma_1} \bar{v}_\alpha^2 dS \\ &= \alpha \mu_\alpha \int_{\Gamma_1} \bar{v}_\alpha^2 dS + \frac{2}{q} \int_{\partial \Gamma_1} \xi_\alpha \bar{v}_\alpha^q |x| dS - \int_{\partial \Gamma_1} \alpha \mu_\alpha \bar{v}_\alpha^2 |x| dS. \end{aligned}$$

Thus

$$\begin{aligned} (3.13) \quad & I(R_\alpha, \bar{v}_\alpha) \\ &= \frac{\alpha \mu_\alpha}{2} \int_{\Gamma_1} \bar{v}_\alpha^2 dS + O \left( \int_{\partial \Gamma_1} (\bar{v}_\alpha^q + \alpha \mu_\alpha \bar{v}_\alpha^2) |x| dS \right) \\ & \quad + O \left( \int_{\Gamma_1} [\mu_\alpha |x'|^2 |\nabla \bar{v}_\alpha|^2 + \mu_\alpha |x'| \bar{v}_\alpha |\nabla \bar{v}_\alpha|] dS \right). \end{aligned}$$

Clearly

$$(3.14) \quad J(R_\alpha, \bar{v}_\alpha) = O\left(\int_{\Gamma_2} (|x| |\nabla \bar{v}_\alpha|^2 + \bar{v}_\alpha |\nabla \bar{v}_\alpha|) dS\right).$$

In view of all the above estimates, we rewrite (3.10) as the following Pohozaev-type identity:

$$\begin{aligned}
 & \alpha \mu_\alpha \int_{\Gamma_1} \bar{v}_\alpha^2 dS \\
 &= O\left(\int_{B_{R_\alpha}^+} \mu_\alpha |x|^2 |\nabla \bar{v}_\alpha| |\nabla^2 \bar{v}_\alpha| dV\right) + O\left(\int_{B_{R_\alpha}^+} \mu_\alpha |x| |\nabla \bar{v}_\alpha|^2 dV\right) \\
 &+ O\left(\int_{B_{R_\alpha}^+} \mu_\alpha |x| \bar{v}_\alpha |\nabla^2 \bar{v}_\alpha| dV\right) + O\left(\int_{B_{R_\alpha}^+} \mu_\alpha \bar{v}_\alpha |\nabla \bar{v}_\alpha| dV\right) \\
 (3.15) \quad &+ O\left(\int_{\Gamma_2} (|x| |\nabla \bar{v}_\alpha|^2 + \bar{v}_\alpha |\nabla \bar{v}_\alpha|) dS\right) \\
 &+ O\left(\int_{\partial \Gamma_1} (\bar{v}_\alpha^q + \alpha \mu_\alpha \bar{v}_\alpha^2) |x| dS\right) \\
 &+ O\left(\int_{\Gamma_1} [\mu_\alpha |x'|^2 |\nabla \bar{v}_\alpha|^2 + \mu_\alpha |x'| \bar{v}_\alpha |\nabla \bar{v}_\alpha|] dS\right).
 \end{aligned}$$

We will derive a contradiction from (3.15) by showing that the left-hand side is much larger than the right-hand side for  $\alpha$  large.

LEMMA 3.3 *For  $n \geq 3$ , there exists some constant  $C > 0$  independent of  $\alpha$  such that  $\int_{\Gamma_1} \bar{v}_\alpha^2 dS > 1/C$  for all  $\alpha \geq 1$ . Moreover, for  $n = 3$ ,  $\int_{\Gamma_1} \bar{v}_\alpha^2 dS \geq (\log R_\alpha)/C$  for all  $\alpha \geq 1$ .*

PROOF: We only need to prove the lemma for large  $\alpha$ . It follows easily from (3.3) that

$$\int_{\Gamma_1} \bar{v}_\alpha^2 dS \geq \frac{1}{C}.$$

For  $n = 3$ , it follows from Proposition 3.2 that

$$\int_{\Gamma_1} \bar{v}_\alpha^2 dS \geq \frac{1}{C} \int_{\partial \mathbb{R}_+^3 \cap B_{R_\alpha^{1/4}}} \left(\frac{1}{1+|x|}\right)^2 dS \geq \frac{\log R_\alpha}{C}.$$

■

LEMMA 3.4 *The following estimates hold:*

$$\begin{aligned}
\int_{\partial\Gamma_1} \bar{v}_\alpha^q |x| dS &\leq CR_\alpha^{1-n} = C(\alpha\mu_\alpha)^{n-1}, \\
\int_{\partial\Gamma_1} \alpha\mu_\alpha \bar{v}_\alpha^2 |x| dS &\leq \alpha\mu_\alpha R_\alpha^{3-n}, \\
\int_{\Gamma_1} (\mu_\alpha |x'|^2 |\nabla \bar{v}_\alpha|^2 + \mu_\alpha |x'| \bar{v}_\alpha |\nabla \bar{v}_\alpha|) dS &\leq \begin{cases} C\mu_\alpha \log R_\alpha & n=3 \\ C\mu_\alpha & n \geq 4, \end{cases} \\
\int_{\Gamma_2} (|x| |\nabla \bar{v}_\alpha|^2 + \bar{v}_\alpha |\nabla \bar{v}_\alpha|) dS &\leq C(\alpha\mu_\alpha)^{n-2}, \\
\int_{B_{R_\alpha}^+} (\mu_\alpha |x|^2 |\nabla \bar{v}_\alpha| |\nabla^2 \bar{v}_\alpha| + \mu_\alpha |x| |\nabla \bar{v}_\alpha|^2) dV &\leq \begin{cases} C\mu_\alpha \log R_\alpha & n=3 \\ C\mu_\alpha & n \geq 4, \end{cases} \\
\int_{B_{R_\alpha}^+} (\mu_\alpha |x| \bar{v}_\alpha |\nabla^2 \bar{v}_\alpha| + \mu_\alpha \bar{v}_\alpha |\nabla \bar{v}_\alpha|) dV &\leq \begin{cases} C\mu_\alpha \log R_\alpha & n=3 \\ C\mu_\alpha & n \geq 4, \end{cases}
\end{aligned}$$

PROOF: These estimates follow easily from (3.4), Proposition 3.1, and some elementary calculations.  $\blacksquare$

PROOF OF THEOREM 0.1: We draw a contradiction from (3.15) by using Lemmas 3.3 and 3.4 because the left-hand side is clearly much larger than the right-hand side in (3.15) as  $\alpha$  tends to infinity.  $\blacksquare$

## Appendix A

Let  $(M, g)$  be a smooth, compact Riemannian manifold of dimension  $n$  ( $n \geq 3$ ) with boundary. In this appendix we present some weighted Sobolev embedding inequalities that should be well-known. We include a proof for completeness.

**THEOREM A.1** *There exists some constant  $C = C(M, g)$  such that for all  $x_0 \in \overline{M}$ ,  $\mu > 0$ ,  $u \in H^1(M)$ ,  $u(x) = 0 \forall d(x_0, x) < \mu$ , we have*

$$\left( \int_M \frac{|u(x)|^p}{d(x_0, x)^{2n}} dv_g \right)^{2/p} \leq C \int_M \frac{|\nabla_g u(x)|^2}{d(x_0, x)^{2n-4}} dv_g$$

and

$$\left( \int_{\partial M} \frac{|u(x)|^q}{d(x_0, x)^{2n-2}} ds_g \right)^{2/q} \leq C \int_M \frac{|\nabla_g u(x)|^2}{d(x_0, x)^{2n-4}} dv_g,$$

where  $d(x_0, x)$  denotes the distance between  $x_0$  and  $x$ .

Theorem A.1 in the case  $x_0 \in \partial M$  follows immediately from Lemma A.2 and Lemma A.4 below. The general case can be proved in a similar way.

LEMMA A.2 *For  $n \geq 3$ , there exists some constant  $C = C(n) > 0$  such that for all  $u \in H^1(B_1^+(0))$ ,  $u \equiv 0$  in an open neighborhood of  $x = 0$ , we have*

$$\left( \int_{B_1^+(0)} \frac{|u(x)|^p}{|x|^{2n}} dx \right)^{2/p} \leq C \int_{B_1^+(0)} \frac{|\nabla u(x)|^2}{|x|^{2n-4}} dx$$

and

$$\left( \int_{|x'| < 1} \frac{|u(x', 0)|^q}{|x'|^{2n-2}} dx' \right)^{2/q} \leq C \int_{B_1^+(0)} \frac{|\nabla u(x)|^2}{|x|^{2n-4}} dx,$$

where  $x = (x', x_n)$ ,  $p = 2n/(n-2)$ ,  $q = 2(n-1)/(n-2)$ ,  $B_1^+(0) = \{x \in \mathbb{R}^n : |x| < 1, x_n > 0\}$ .

PROOF: It follows from the hypothesis that for some  $\mu = \mu(u) > 0$ ,  $u(x) = 0 \forall |x| < \mu$ ,  $x_n > 0$ . Consider

$$v(y) = u(y/|y|^2), \quad |y| > 1, \quad y_n > 0.$$

Clearly

$$v(y) = 0 \quad \forall |y| > 1/\mu, \quad y_n > 0,$$

and, for some  $C(n) > 0$ ,

$$\begin{aligned} \int_{B_1^+(0)} \frac{|u(x)|^p}{|x|^{2n}} dx &= C(n) \int_{\{|y| \geq 1, y_n \geq 0\}} |v(y)|^p dy, \\ \int_{B_1^+(0)} \frac{|\nabla u(x)|^2}{|x|^{2n-4}} dx &= C(n) \int_{\{|y| \geq 1, y_n \geq 0\}} |\nabla v(y)|^2 dy, \\ \int_{|x'| < 1} \frac{|u(x', 0)|^q}{|x'|^{2n-2}} dx' &= C(n) \int_{|y'| > 1} |v(y', 0)|^q dy'. \end{aligned}$$

It follows from standard Sobolev embedding theorems (with appropriate extensions of  $v$  to  $|y| < 1$ ) that

$$\begin{aligned} \left( \int_{\{|y| \geq 1, y_n \geq 0\}} |v(y)|^p dy \right)^{2/p} + \left( \int_{|y'| > 1} |v(y', 0)|^q dy' \right)^{2/q} \\ \leq C(n) \int_{\{|y| \geq 1, y_n \geq 0\}} |\nabla v(y)|^2 dy. \end{aligned}$$

Lemma A.2 follows immediately. ■

The following corollary is immediate:

**COROLLARY A.3** *There exist some constants  $\delta = \delta(M, g) > 0$  and  $C = C(M, g) > 0$  such that for all  $x_0 \in \partial M$ ,  $u \in H^1(M)$ ,  $u \equiv 0$  in an open neighborhood of  $x_0$  we have*

$$\begin{aligned} \left( \int_{B_\delta^+(x_0)} \frac{|u(x)|^p}{d(x_0, x)^{2n}} dv_g \right)^{2/p} + \left( \int_{\partial M \cap \overline{B_\delta^+(x_0)}} \frac{|u(x)|^q}{d(x_0, x)^{2n-2}} ds_g \right)^{2/q} \\ \leq C \int_{B_\delta^+(x_0)} \frac{|\nabla_g u(x)|^2}{d(x_0, x)^{2n-4}} dv_g, \end{aligned}$$

where  $B_\delta^+(x_0) = \{x \in M : d(x_0, x) < \delta\}$ .

**LEMMA A.4** *For  $\delta > 0$ , there exists  $C = C(M, g, \delta) > 0$  such that for all  $x_0 \in \overline{M}$ ,  $u \in H^1(M \setminus B_{\delta/2}^+(x_0))$ , we have*

$$\begin{aligned} \left( \int_{M \setminus B_\delta^+(x_0)} |u|^p dv_g \right)^{2/p} + \left( \int_{\partial M \setminus \overline{B_\delta^+(x_0)}} |u|^q ds_g \right)^{2/q} \\ \leq C \left\{ \int_{M \setminus B_{\delta/2}^+(x_0)} |\nabla_g u|^2 dv_g + \int_{B_\delta^+(x_0) \setminus B_{\delta/2}^+(x_0)} u^2 dv_g \right\}. \end{aligned}$$

**PROOF:** Suppose the contrary of Lemma A.4, namely, that for some  $\delta > 0$ , there exists a sequence of points  $\{x_i\} \in \overline{M}$ ,  $\{u_i\} \in H^1(M \setminus B_{\delta/2}^+(x_i))$ , satisfying

$$(A.1) \quad \left( \int_{M \setminus B_\delta^+(x_i)} |u_i|^p dv_g \right)^{2/p} + \left( \int_{\partial M \setminus \overline{B_\delta^+(x_i)}} |u_i|^q ds_g \right)^{2/q} = 1$$

and

$$\int_{M \setminus B_{\delta/2}^+(x_i)} |\nabla_g u_i|^2 dv_g + \int_{B_\delta^+(x_i) \setminus B_{\delta/2}^+(x_i)} u_i^2 dv_g < \frac{1}{i}.$$

It follows that

$$\|u_i\|_{H^1(M \setminus B_{\delta/2}^+(x_i))} \leq C$$

and

$$(A.2) \quad \lim_{i \rightarrow \infty} \left\{ \int_{M \setminus B_{\delta/2}^+(x_i)} |\nabla_g u_i|^2 dv_g + \int_{B_\delta^+(x_i) \setminus B_{\delta/2}^+(x_i)} u_i^2 dv_g \right\} = 0.$$

After passing to some subsequence, we have that  $u_i$  converges weakly to  $u$  in  $H^1(M \setminus B_{\delta/2}^+(x_i))$ . In view of (A.2),  $u \equiv 0$ . It follows from the compact embedding of  $H^1$  into  $L^2$  that

$$\int_{M \setminus B_{\delta/2}^+(x_i)} u_i^2 \rightarrow 0.$$

This, together with (A.2), yields

$$\|u_i\|_{H^1(M \setminus B_{\delta/2}^+(x_i))} \rightarrow 0$$

which contradicts (A.1) because of the Sobolev embedding theorems. Lemma A.4 is established.  $\blacksquare$

## Appendix B

For  $n \geq 3$ , let  $B_1$  denote the unit ball in  $\mathbb{R}^n$ , and let  $g = g_{ij}(x)dx^i dx^j$  be a  $C^2$ -metric on  $B_1$  with  $g_{ij}(0) = \delta_{ij}$  and  $\frac{\partial g_{ij}}{\partial x_k}(0) = 0$  for all  $1 \leq i, j, k \leq n$ . Let  $K \in L^\infty(B_1)$ .

**PROPOSITION B.1** *There exists some constant  $\tau_1 > 0$  depending only on  $n$ ,  $\|g_{ij}\|_{C^2(B_1)}$ , and  $\|K\|_{L^\infty(B_1)}$  such that for all  $0 < \tau \leq \tau_1$ , there exists some function  $G(y) = |y|^{2-n} + E(y)$  satisfying*

$$(B.1) \quad \begin{cases} -\Delta_g G + K(y)G = n(n-2)\omega_n\delta_0 & \text{in } B_\tau \setminus \{0\} \\ G = 0 & \text{on } \partial B_\tau, \end{cases}$$

where  $\omega_n$  is the volume of the unit ball in  $\mathbb{R}^n$  and  $E$  satisfies the following: For all  $0 < \varepsilon < 1$ , there exists some constant  $C(\varepsilon)$  depending only on  $\varepsilon$ ,  $n$ ,  $\|g_{ij}\|_{C^2(B_1)}$ , and  $\|K\|_{L^\infty(B_1)}$  such that

$$|y|^{n-4+\varepsilon}|E(y)| + |y|^{n-3+\varepsilon}|\nabla E(y)| \leq C(\varepsilon) \quad \forall y \in B_\tau, n \geq 4,$$

and

$$|y|^{\varepsilon-1}|E(y) - E(0)| + |y|^\varepsilon|\nabla E(y)| \leq C(\varepsilon) \quad \forall y \in B_\tau, n = 3.$$

**REMARK B.2** *In fact, such  $G$  is unique.*

**PROOF:** The existence of  $\tau_1 > 0$  is well-known; see, for example, [7]. Clearly (B.1) is equivalent to

$$(B.2) \quad \begin{cases} -\Delta_g E + K(y)E = O(|y|^{2-n}) & \text{in } B_\tau \\ E = -\tau^{2-n} & \text{on } \partial B_\tau. \end{cases}$$

Since  $|y|^{2-n} \in L^r(B_\tau)$  for all  $r < n/(n-2)$ , it is well-known that (B.2) has a unique solution  $E \in W^{2,r}(B_\tau)$ . It follows from the Sobolev embedding theorems that

$$\|E\|_{L^s(B_\tau)} \leq \begin{cases} C(s) & \forall s < \frac{n}{(n-4)}, n \geq 5, \\ C(s) & \forall s < \infty, n = 4, \end{cases}$$

and

$$(B.3) \quad \|E\|_{C^{1-\varepsilon}(B_\tau)} \leq C(\varepsilon) \quad \forall 0 < \varepsilon < 1, n = 3.$$

For  $0 < r \leq \tau/5$ ,  $x \in A_0 = \{x : \frac{1}{5} \leq |x| \leq 5\}$ , set

$$E_1(x) = \begin{cases} r^{n-2}E(rx) & n \geq 4 \\ r^{n-2}(E(rx) - E(0)) & n = 3. \end{cases}$$

Then  $E_1$  satisfies

$$-\Delta_h E_1(x) + K(rx)r^2 E_1(x) = O(r^2) \quad x \in A_0,$$

where  $|O(r^2)| \leq Cr^2$  with  $C$  independent of  $r$  and  $h = h_{ij}(x)dx^i dx^j = g_{ij}(rx)dx^i dx^j$ . For  $n \geq 4$ , for all  $0 < \varepsilon < 1$ , we can choose some  $s_1 = s_1(\varepsilon) < n/(n-4)$  such that

$$\|E_1\|_{L^{s_1}(A_0)} \leq r^{2-\varepsilon} \|E\|_{L^{s_1}(B_\tau)} \leq C(\varepsilon)r^{2-\varepsilon}.$$

Using the equation of  $E_1$  and applying the bootstrap method finite times (using the  $L^p$  theory of elliptic equations and the Sobolev embedding theorems, see e.g., [17]), we have

$$|E_1(x)| + |\nabla E_1(x)| \leq C(\varepsilon)r^{2-\varepsilon}, \quad \frac{1}{2} \leq |x| \leq 2.$$

Consequently,

$$|y|^{n-2}|E(y)| + |y|^{n-1}|\nabla E(y)| \leq C(\varepsilon)|y|^{2-\varepsilon}, \quad |y| \leq \frac{\tau}{5}.$$

This establishes Proposition B.1 in the case  $n \geq 4$ . For  $n = 3$ , we know from (B.3), for all  $0 < \varepsilon < 1$ , that

$$|E_1(x)| \leq C(\varepsilon)r^{2-\varepsilon} \quad \forall x \in A_0.$$

It follows from the equation of  $E_1$  and standard elliptic estimates that

$$|\nabla E_1(x)| \leq C(\varepsilon)r^{2-\varepsilon} \quad \forall \frac{1}{2} \leq |x| \leq 2.$$

Consequently,

$$|y| |E(y) - E(0)| + |y|^2 |\nabla E(y)| \leq C(\varepsilon) |y|^{2-\varepsilon}, \quad |y| \leq \frac{\tau}{5}.$$

Proposition B.1 is thus established. ■

**Acknowledgement.** We thank H. Brezis for his interest in the work as well as for informing us about the work of Adimurthi and Yadava [1] while we were working on this paper. Part of this work was completed while the first author was visiting the Courant Institute; he would like to express his thanks to L. Nirenberg for the arrangement and his kind hospitality. The first author appreciates the support from the Alfred Sloan Foundation and from National Science Foundation Grant DMS-9401815.

### Bibliography

- [1] Adimurthi and Yadava, S. L., *Some remarks on Sobolev type inequalities*, Calc. Var. Partial Differential Equations 2, 1994, pp. 427–442.
- [2] Atkinson, F. V., and Peletier, L. A., *Elliptic equations with nearly critical growth*, J. Differential Equations 70, 1987, pp. 349–365.
- [3] Aubin, T., *Espaces de Sobolev sur les variétés Riemanniennes*, Bull. Sci. Math. (2) 100, 1976, pp. 149–173.
- [4] Aubin, T., *Problèmes isopérimétriques et espaces de Sobolev*, J. Differential Geometry 11, 1976, pp. 573–598.
- [5] Aubin, T., *Nonlinear Analysis on Manifolds. Monge-Ampère Equations*, Grundlehren der Mathematischen Wissenschaften [Fundamental Principles of Mathematical Science] No. 252, Springer-Verlag, New York–Berlin, 1982.
- [6] Beckner, W., *Sharp Sobolev inequalities on the sphere and the Moser-Trudinger inequality*, Ann. of Math. (2) 138, 1993, pp. 213–242.
- [7] Berestycki, H., Nirenberg, L., and Varadhan, S. R. S., *The principal eigenvalue and maximum principle for second-order elliptic operators in general domains*, Comm. Pure Appl. Math. 47, 1994, pp. 47–92.
- [8] Brezis, H., and Lieb, E. H., *Sobolev inequalities with remainder terms*, J. Funct. Anal. 62, 1985, pp. 73–86.
- [9] Brezis, H., and Nirenberg, L., *Positive solutions of nonlinear elliptic equations involving critical Sobolev exponents*, Comm. Pure Appl. Math. 36, 1983, pp. 437–477.
- [10] Brezis, H., and Peletier, L. A., *Asymptotics for elliptic equations involving critical growth*, pp. 149–192 in: *Partial Differential Equations and the Calculus of Variations, Vol. I*, Progress in Nonlinear Differential Equations and Their Applications No. 1, Birkhäuser Boston, Boston, 1989.
- [11] Carleson, L., and Chang, S. Y. A., *On the existence of an extremal function for an inequality of J. Moser*, Bull. Sci. Math. 110, 1986, pp. 113–127.

- [12] Chang, S.-Y. A., Gursky, M. J., and Yang, P. C., *The scalar curvature equation on 2- and 3-spheres*, Calc. Var. Partial Differential Equations 1, 1993, pp. 205–229.
- [13] Cherrier, P., *Problèmes de Neumann non linéaires sur les variétés Riemanniennes*, J. Funct. Anal. 57, 1984, pp. 154–206.
- [14] Escobar, J. F., *Uniqueness theorems on conformal deformation of metrics, Sobolev inequalities, and an eigenvalue estimate*, Comm. Pure Appl. Math. 43, 1990, pp. 857–883.
- [15] Escobar, J. F., *Conformal deformation of a Riemannian metric to a scalar flat metric with constant mean curvature on the boundary*, Ann. of Math. (2) 136, 1992, pp. 1–50.
- [16] Escobar, J. F., *Sharp constant in a Sobolev trace inequality*, Indiana Univ. Math. J. 37, 1988, pp. 687–698.
- [17] Gilbarg, D., and Trudinger, N. S., *Elliptic Partial Differential Equations of Second Order*, 2nd ed., Grundlehren der Mathematischen Wissenschaften [Fundamental Principles of Mathematical Sciences] No. 224, Springer-Verlag, Berlin–New York, 1983.
- [18] Han, Z.-C., *Asymptotic approach to singular solutions for nonlinear elliptic equations involving critical Sobolev exponent*, Ann. Inst. H. Poincaré Anal. Non Linéaire 8, 1991, pp. 159–174.
- [19] Hebey, E., *Optimal Sobolev inequalities on complete Riemannian manifolds with Ricci curvature bounded below and positive injectivity radius*, Amer. J. Math. 118, 1996, pp. 291–300.
- [20] Hebey, E., and Vaugon, M., *The best constant problem in the Sobolev embedding theorem for complete Riemannian manifolds*, Duke Math. J. 79, 1995, pp. 235–279.
- [21] Hebey, E., and Vaugon, M., *Meilleures constantes dans le théorème d'inclusion de Sobolev*, Ann. Inst. H. Poincaré Anal. Non Linéaire 13, 1996, pp. 57–93.
- [22] Kazdan, J. L., and Warner, F. W., *Remarks on some quasi-linear elliptic equations*, Comm. Pure Appl. Math. 28, 1975, pp. 567–597.
- [23] Li, Y.Y., *The Nirenberg problem in a domain with boundary*, Topol. Methods Nonlinear Anal. 6, 1995, pp. 309–329.
- [24] Li, Y.Y., *Prescribing scalar curvature on  $S^n$  and related problems, I*, J. Differential Equations 120, 1995, pp. 319–410.
- [25] Li, Y.Y., *Prescribing scalar curvature on  $S^n$  and related problems, II. Existence and compactness*, Comm. Pure Appl. Math. 49, 1996, pp. 541–597.
- [26] Li, Y.Y., and Zhu, M., *Uniqueness theorems through the method of moving spheres*, Duke Math. J. 80, 1995, pp. 383–417.
- [27] Lieb, E. H., *Existence and uniqueness of the minimizing solution of Choquard's nonlinear equation*, Studies in Appl. Math. 57, 1976/77, pp. 93–105.
- [28] Lieb, E. H., *Sharp constants in the Hardy-Littlewood-Sobolev and related inequalities*, Ann. of Math. 118, 1983, pp. 349–374.
- [29] Lions, P.-L., *The concentration-compactness principle in the calculus of variations, The limit case, II*, Rev. Mat. Iberoamericana 1, 1985, pp. 145–121.
- [30] Moser, J., *A sharp form of an inequality by N. Trudinger*, Indiana Univ. Math. J. 20, 1970/71, pp. 1077–1092.
- [31] Rey, O., *Proof of two conjectures of H. Brézis and L. A. Peletier*, Manuscripta Math. 65, 1989, pp. 19–37.
- [32] Schoen, R., *On the number of constant scalar curvature metrics in a conformal class*, pp. 311–320 in: *Differential Geometry*, Pitman Monographs and Surveys in Pure and Applied Mathematics No. 52, Longman Scientific & Technical, Harlow, 1991.

- [33] Schoen, R. and Zhang, D., *Prescribed scalar curvature on the  $n$ -sphere*, Calc. Var. Partial Differential Equations 4, 1996, pp. 1–25.
- [34] Talenti, G., *Best constant in Sobolev inequality*, Ann. Mat. Pura Appl. (4) 110, 1976, pp. 353–372.
- [35] Trudinger, N., *On imbeddings into Orlicz spaces and some applications*, J. Math. Mech. 17, 1967, pp. 473–483.
- [36] Zhang, D., *New results on geometric variational problems*, Thesis, Stanford University, 1990.

YANYAN LI

Department of Mathematics  
Rutgers University  
New Brunswick, NJ 08903  
E-mail:  
yyli@math.rutgers.edu

MEIJUN ZHU

Department of Mathematics  
University of British Columbia  
Vancouver, BC  
CANADA V6T 1Z2  
E-mail: mzhu@math.ubc.ca

Received April 1996.