

Problems 28, 31, 37 from Section 3.5 of the book.

*Hint to Problem 28(a):* Use Proposition 3.13(a).

**Additional problem 1.** Let  $(x, y)$  and  $(r, \theta)$  be the Cartesian and the polar coordinates in the plane  $\mathbb{R}^2$ , respectively, and let  $\mathbf{0}$  be the origin,  $(x, y) = (0, 0)$ . For  $n \in \mathbb{N}$  define the numbers

$$a_n = \frac{1}{10^n} \left( 1 - \frac{1}{10^n} \right), \quad b_n = \frac{1}{10^n}$$

and the sets  $D_n = \{(x, y) \in \mathbb{R}^2 : r \in [a_n, b_n)\}$ . Each  $D_n$  is an *annulus* (plural: *annuli*), i.e., the area between two concentric circles:  $D_n = B(b_n, \mathbf{0}) \setminus B(a_n, \mathbf{0})$ . Note that  $D_n$  are disjoint. Define the function  $f : \mathbb{R}^2 \rightarrow \mathbb{R}$  as follows:

$$f(x, y) = \sum_{n \in \mathbb{N}} \chi_{D_n}(x, y),$$

i.e.,  $f(x, y)$  is 1 if  $(x, y)$  belongs to some annulus  $D_n$  and 0 otherwise. Note that  $f(\mathbf{0}) = 0$ .

(a) Is the function  $f$  continuous at  $\mathbf{0}$ ?

(b) Show that the Lebesgue measure (i.e., the area) of  $D_n$  is smaller than  $\frac{2\pi}{10^{3n}}$ .

(c) Let  $(x, y) \in D_n$ , i.e.,  $r \in [a_n, b_n)$ . Prove that the average  $(A_r f)(\mathbf{0})$  of  $f$  over the ball  $B(r, \mathbf{0})$  decreases with  $n$  as  $\frac{C}{10^n}$  for some constant  $C > 0$ .

*Solution:*

$$\begin{aligned} (A_r f)(\mathbf{0}) &= \frac{1}{m(B(r, \mathbf{0}))} \int_{B(r, \mathbf{0})} f \, dm \leq \frac{1}{m(B(a_n, \mathbf{0}))} \int_{B(b_n, \mathbf{0})} f \, dm \\ &= \frac{1}{\pi a_n^2} \sum_{j=n}^{\infty} m(D_j) < \frac{10^{2n}}{\pi \left(1 - \frac{1}{10^{2n}}\right)^2} \sum_{j=n}^{\infty} \frac{2\pi}{10^{3j}} \\ &\leq \frac{2 \cdot 10^{2n}}{\pi} \frac{2\pi}{10^{3n}} \sum_{k=0}^{\infty} \frac{1}{10^{3k}} = \frac{C}{10^n}. \end{aligned}$$

(d) Let  $(x, y)$  be a point in the area between  $D_n$  and  $D_{n+1}$ , i.e, let the distance from  $(x, y)$  to  $\mathbf{0}$  be  $r \in [b_{n+1}, a_n)$ . Give an upper bound on the average  $(A_r f)(\mathbf{0})$  of  $f$  over the ball  $B(r, \mathbf{0})$  in terms of  $n$  similarly to the bound in part (c).

(e) Based on the bounds in parts (c) and (d), what can you conclude about the behavior of the averages  $(A_r f)(\mathbf{0})$  as  $r \rightarrow 0$ ? How about the Hardy-Littlewood maximal function  $(Hf)(\mathbf{0})$ ? Is the point  $\mathbf{0}$  in the Lebesgue set of  $f$ ?

**Food for thought.** Problems 27, 29 from Section 3.5 of the book.

**Food for thought.** Read Examples 1, 2, 14, 15, 16, 17 from Chapter 8 of *Counterexamples in Analysis* by Gelbaum and Olmsted, and think about the properties of the Cantor function and the measure associated with it.