

Regularity of critical invariant circles of non-twist maps

A. Apte

Department of Physics, University of Texas at Austin

*1 University Station C1500, Austin, TX 78712-0262**

Rafael de la Llave

Department of Mathematics, The University of Texas at Austin

1 University Station C1200, Austin, TX 78712-0257†

Nikola P. Petrov

Department of Mathematics and Michigan Center for Theoretical Physics

University of Michigan

Ann Arbor, MI 48109-1109‡

(Dated: October 6, 2004)

We study critical invariant circles of several noble rotation numbers at the edge of breakdown for area preserving maps of the cylinder which violate the twist conditions.

These circles admit essentially unique parameterizations by rotational coordinates. We present a high accuracy computation of about 10^7 Fourier coefficients. This allows us to compute the regularity of the conjugating maps and show that, to the extent of the precision, it only depends on the tail of the continued fraction expansion.

PACS numbers: 05.10.Cc 02.30.Nw 02.30.-f 47.53.+n

Keywords: non-twist maps, renormalization group, regularity of functions, Fourier analysis

*Electronic address: apte@physics.utexas.edu

†Electronic address: llave@math.utexas.edu

‡Electronic address: npetrov@umich.edu

I. INTRODUCTION

Area-preserving maps of a cylinder have been studied for several decades both as low-dimensional models of physical systems and as interesting mathematical problems. Until recently, most of these studies have dealt with *twist* maps which are the maps that satisfy a certain non-degeneracy condition. Lately, maps that violate this condition locally in phase space have been of great interest. These *nontwist* maps occur in various physical systems such as, e.g., the study of magnetic field lines in toroidal plasma devices with reversed shear profile, channel flows, and other physical applications (see, e.g., [1–8]).

Non-twist maps present new phenomena which are not observed in twist maps. Some of these phenomena (reconnection, bifurcation, meandering tori, etc.) are discussed in [8–14]. It should also be remarked that non-twist maps appear naturally in iterations of twist maps. This is because of the fact that, even though the twist property has dynamical consequences, it is not preserved by composition or by a change of variables.

These maps are also of mathematical interest because many important results about maps, e.g., KAM and Aubry-Mather theory, depend on twist condition. Even though Aubry-Mather variational theory for twist maps does not generalize for non-twist maps, there is an analogue of KAM theory. In [15, 16] it was suggested that one can continue shearless circles with a Diophantine rotation number provided that one adjusts a second parameter; this was proved rigorously in [17]. Furthermore, in [15, 16] it was observed numerically that if the curve is continued, one reaches a critical value where the circles cannot be continued anymore. The circle that cannot be continued – henceforth referred to as the critical circle (or circle at breakup) – enjoys remarkable self-similar properties which are universal (that is, they do not seem to depend on the map). The calculations of critical shearless invariant tori in [16] were extended and refined in [18–20].

As we will show later, a natural way of representing the circle is by giving some parameterization in which the motion becomes a rotation. This parameterization is essentially unique. The study of the Fourier spectrum reveals several properties of the function. One such property to measure is the regularity of the parameterization. The study of the regularity of functions – which is crucial in KAM theory – has been studied extensively in the mathematical literature. Efficient algorithms for problems very similar to those considered here were developed in [21].

In this paper, we analyze further the properties of the critical invariant tori computed in [20] by using techniques developed in [22]. Since the new calculations in [20] allow us to compute accurately orbits of length several millions, we have a very accurate computation of the critical invariant circle.

We have found that the functions corresponding to the critical shearless circle saturate several bounds in harmonic analysis, so that the regularity is easy to ascertain. In the cases we have examined, the regularity of the critical circles is the same for different rotation numbers with the same tail of the continued fraction expansions. This is consistent with the predictions of universality that follow from the renormalization group picture.[23]

The paper is organized as follows. Section II introduces the basic terminology of the nontwist maps. In Sec. III, we describe the convenient parameterization used to study these regularities and its symmetry properties. The numerical techniques are briefly explained in Sec. IV. We discuss the results in Sec. V.

II. PRELIMINARIES

The *standard nontwist map* $F : \mathbb{T}^1 \times \mathbb{R} \rightarrow \mathbb{T}^1 \times \mathbb{R}$ is given by

$$y' = y - b \sin(2\pi x), \quad x' = x + a(1 - y'^2) \pmod{1}, \quad (1)$$

where $(x, y) \in \mathbb{T}^1 \times \mathbb{R}$ and a, b are parameters. This map violates the *twist condition*

$$\frac{\partial x'}{\partial y} \neq 0, \quad (2)$$

along the curve $y = b \sin(2\pi x)$ in the phase space. An *orbit* of the map is the set of points $\{(x_i, y_i)\}_{i=-\infty}^{\infty}$ such that $(x_{i+1}, y_{i+1}) = F(x_i, y_i)$. An orbit is *periodic* of least period n if $x_{i+n} = x_i$, $y_{i+n} = y_i$, and n is the smallest such integer. We also consider the lift of the nontwist map $\tilde{F} : \mathbb{R} \times \mathbb{R} \rightarrow \mathbb{R} \times \mathbb{R}$ given by (1) by removing the $\pmod{1}$. We denote by $\{(\tilde{x}_i, \tilde{y}_i)\}$ an orbit of the lift map corresponding to the orbit $\{(x_i, y_i)\}$ of the map. The *winding* (or *rotation*) *number* ω of an orbit is defined as

$$\omega = \lim_{i \rightarrow \infty} \frac{\tilde{x}_i}{i}, \quad (3)$$

if the limit exists. This number is independent of the lift taken. The winding number of a periodic orbit is rational; an orbit for which winding number doesn't exist is chaotic. A set

homeomorphic to a circle that is mapped to itself by F is called an *invariant circle*. It was already proved by Poincaré that all the points in a circle have the same winding number. The winding number of orbits belonging to these circles is generically irrational.

Invariant circles that are homotopically nontrivial form barriers to transport, i.e., points on one side cannot cross to the other side. (We will abbreviate homotopically nontrivial invariant circles to just invariant circles.) When the perturbation increases, some of the circles cease to exist leaving behind invariant sets homeomorphic to Cantor sets. The breakup of these circles (in both twist and nontwist maps) has been studied extensively.

In contrast to twist maps, the orbits of the standard nontwist map are generally found in pairs – two or none – for any winding number. Also, for the standard nontwist map, the rotation numbers are bounded from above. The orbit with the extremal winding number occurs in the region in phase space where the twist condition fails. The curve in parameter space along which there is only one orbit of winding number ω is called the ω -*bifurcation curve* and the orbit itself is called the *shearless*. The rotation number of the shearless orbit is an extremum of all the rotation numbers of orbits in the map.

Remark 1. For the integrable map $b = 0$, the circle with rotation number a is shearless. Thus, the parameter values $(a, 0)$ belong to a -bifurcation curve (i.e., to the ω -bifurcation curve for $\omega = a$). It was shown in [17] that if a is Diophantine, then there is an analytic a -bifurcation curve for small perturbations.

Shearless invariant circles are present in the phase space region where the twist condition fails. The breakup of shearless circles with noble winding numbers (with a continued fraction expansion ending in 1) has been studied extensively in [15, 18, 20]. The breakup of shearless invariant circles for nontwist time periodic Hamiltonian flows has been analyzed in [19, 24].

The critical circles for the three winding numbers

$$\begin{aligned}\omega^{(1)} &= [0, 1, 1, 1, \dots] = 1/\gamma, \\ \omega^{(2)} &= [0, 2, 1, 1, 1, \dots] = 1/\gamma^2, \\ \omega^{(3)} &= [0, 2, 2, 1, 1, 1, \dots] = \gamma^2/(1 + 2\gamma^2),\end{aligned}\tag{4}$$

where γ is the golden mean, were studied in [20]. Here $\omega = [a_0, a_1, a_2, \dots]$ denotes the




FIG. 1: Critical circles with winding numbers $\omega^{(1)}$, $\omega^{(2)}$, and $\omega^{(3)}$ of (4).

continued fraction expansion [25] which is defined by

$$\omega = [a_0, a_1, a_2, \dots] := a_0 + \frac{1}{a_1 + \frac{1}{a_2 + \dots}}. \quad (5)$$

The parameter values for which there is a critical circle was found [20] using Greene's residue criterion [26]. This method uses the observation that the invariant circle can be approximated by periodic orbits of winding numbers equal to the convergents obtained by truncating the continued fraction expansion. The residue criterion relates the stability of these periodic orbits with the breakup (loss of stability) of the invariant circle. A rigorous partial justification of this procedure for nontwist maps is presented in [17].

In the case of the standard nontwist map, the ω -bifurcation curve for irrational ω is also approximated by the numerically obtained limit of bifurcation curves of the periodic orbits. This limit shows a scaling behavior which further increases the accuracy of these calculations, for details of which we refer to [20]. The parameter values for which the above circles are critical are as follows:

$$\begin{aligned} (a_c, b_c)^{(1)} &= (0.686049108, 0.74249354915518), \\ (a_c, b_c)^{(2)} &= (0.425160543, 0.92446364703553), \\ (a_c, b_c)^{(3)} &= (0.45297741955, 0.84582913999445). \end{aligned} \quad (6)$$

These critical circles, which are shown in Fig. 1, are not smooth, and they show a self-similar structure [20]. Our aim here is to study numerically the smoothness of these critical circles and details of their fine scale structure using Fourier analysis. We will first describe the method used to parameterize invariant circles of nontwist maps.

III. THE PARAMETERIZATION FUNCTION

In contrast with the situation for twist maps, homotopically non-trivial invariant circles for non-twist maps can fail to be graphs. Indeed, our computations show that the critical shearless invariant circles are not graphs in any smooth coordinate system. A convenient way to describe circles which are not graphs is to find a parametric representation. In our case, it is convenient to choose this parameterization in such a way that, moreover, turns the dynamics into a rotation. Such parameterization functions are often called “hull functions,” and they have been used extensively for computation of periodic orbits and invariant circles. For example, they have been used as the basis for KAM theory in [27].

Therefore, the main object of our study will be a continuous map $K : \mathbb{T}^1 \rightarrow \mathbb{T}^1 \times \mathbb{R}$ whose range is the invariant circle we are seeking to describe and such that

$$F \circ K(\theta) = K(\theta + \omega) . \quad (7)$$

We will denote the components of K by $K_x : \mathbb{T}^1 \rightarrow \mathbb{T}^1$ and $K_y : \mathbb{T}^1 \rightarrow \mathbb{R}$.

If we set $K(0)$ to be a point on the circle, applying (7) we obtain

$$F^n(K(0)) = K(n\omega) . \quad (8)$$

Hence, it is very easy to compute numerically the function K at points in the set $\{\theta = n\omega\}_{n=0}^{\infty}$. Since we are assuming that $\omega \notin \mathbb{Q}$, this is a dense set on the circle and therefore the continuous function K is determined. For the numerical applications, it is quite fortunate that $\{n\omega\}_{n=0}^N$ is very uniformly distributed in the torus.

The argument above shows that the continuous solutions of (7) are unique once we fix a point. More precisely, if K, \hat{K} solve (7), then there is a σ such that

$$\hat{K}(\theta) = K(\theta + \sigma) . \quad (9)$$

Indeed, since $\hat{K}(0)$ is in the circle, it can be written as $K(\sigma)$. Then, applying (7), we obtain $\hat{K}(n\omega) = K(n\omega + \sigma)$. Hence, (9) holds for a dense set.

We will also find it convenient to study lifts \tilde{F} of the map F and lifts \tilde{K} of the function K . In general, a map $\tilde{K} : \mathbb{R} \rightarrow \mathbb{R} \times \mathbb{R}$ is a lift of a map $K : \mathbb{T}^1 \rightarrow \mathbb{T}^1 \times \mathbb{R}$ when the map \tilde{K} becomes the map K upon identifying the angle variables (the independent variable and the first coordinate of the range) that differ by an integer.




FIG. 2: The functions $\tilde{K}_x(\theta) - \theta$ (left) and $\tilde{K}_y(\theta)$ (right) for the three winding numbers from (4). The symmetry properties (Lemma 1) can be seen very clearly. The graphs of $\tilde{K}_x(\theta) - \theta$ have been shifted by a constant for clarity.

Of course, a lift corresponds to only one map K , but a map K has infinitely many lifts. If \tilde{K} is a lift of K , all the other lifts are given by $\tilde{K}(\theta + \ell) + (n, 0)$ where $\ell, n \in \mathbb{Z}$.

The lifts of a map K in the annulus satisfy

$$\tilde{K}(\theta + 1) = \tilde{K}(\theta) + (n, 0) . \quad (10)$$

The number n measures how many turns are experienced by the angle variable in the range of \tilde{K} when the angle variable in the domain of \tilde{K} experiences one turn. We will call n the *index* of the map. In our case, we will consider only circles of index 1. It is easy to see that circles of index bigger than 1 have self-intersections, and circles of index 0 are homotopically trivial. The lift \tilde{F} of our map F of the annulus also satisfies

$$\tilde{F}(x + 1, y) = \tilde{F}(x, y) + (1, 0) .$$

Note that a consequence of (10) is that $\tilde{K}_x(\theta) - \theta$ and $\tilde{K}_y(\theta)$ are periodic functions. These are the functions we will be studying numerically. This is equivalent to studying K because if K satisfies (7), then

$$\tilde{F} \circ \tilde{K}(\tilde{\theta}) = \tilde{K}(\tilde{\theta} + \omega) . \quad (11)$$

The functions $\tilde{K}_x(\theta) - \theta$ and $\tilde{K}_y(\theta)$ for the shearless circles of the winding numbers $\omega^{(1)}$, $\omega^{(2)}$, and $\omega^{(3)}$ are shown in Fig. 2.

A. Symmetry properties

The standard non-twist map (1) is symmetric under S :

$$F \circ S = S \circ F , \quad (12)$$

where S is the map of the annulus defined by

$$S(x, y) = \left(x + \frac{1}{2}, -y \right) .$$

Note that S is an involution:

$$S^2 = \text{Id} . \quad (13)$$

This symmetry (12) implies the following properties for the functions K solving (7).

Lemma 1. *If the map F satisfies (12) and the circle is of index 1, then K satisfies*

$$S \circ K(\theta) = K(\theta + 1/2) ,$$

and any lift \tilde{K} satisfies

$$\tilde{K}_x(\theta + 1/2) = \tilde{K}_x(\theta) + 1/2 , \quad \tilde{K}_y(\theta + 1/2) = -\tilde{K}_y(\theta) . \quad (14)$$

In particular $\tilde{K}_x(\theta) - \theta$ is of period $1/2$.

Proof. We have

$$\begin{aligned} F \circ S \circ K(\theta) &= S \circ F \circ K(\theta) \\ &= S \circ K(\theta + \omega) , \end{aligned}$$

so that $S \circ K$ is also a solution of (7) and parameterizes the same circle. Hence, by the uniqueness result in (9), there exists a number σ for which

$$S \circ K(\theta) = K(\theta + \sigma) .$$

Noting that $S^2 = \text{Id}$, we conclude that

$$K(\theta + 2\sigma) = K(\theta) . \quad (15)$$

If σ were an irrational number, then (15) would imply that $K(\theta)$ is constant, which is impossible. Hence, σ is rational.

If $\sigma = n/m$ with m odd, then

$$SK(\theta) = S^m K(\theta) = K(\theta + n) = K(\theta) .$$

This implies that $-K_y(\theta) = K_y(\theta)$, so that the circle is the x axis and that $K_x(\theta) + 1/2 = K_x(\theta)$, both of which are absurd.

Hence, the only possibility is $\sigma = n/2k$ with $n, k \in \mathbb{N}$, $\gcd(n, k) = 1$, and, in particular n is odd. Then we can find $L, p \in \mathbb{N}$ such that $nL - pk = 1$. Hence, iterating (15) L times, we obtain

$$K(\theta) = S^{2L} K(\theta) = K\left(\theta + \frac{nL}{k}\right) = K\left(\theta + \frac{1}{k}\right) .$$

This will imply that the index of the circle is k . Since we are considering circles of index 1, we choose $k = 1$, and the result is proved. \square

B. Reversibility

The map (1) is reversible, which implies antisymmetry properties of the parameterization function. We follow the notation of [15].

Reversibility means that

$$I_0 \circ F = F^{-1} \circ I_0 , \quad I_0^2 = \text{Id} , \quad (16)$$

where I_0 is a map of the annulus. Then it is easy to see that $I_1 = F \circ I_0$ also satisfies

$$I_1 \circ F = F^{-1} \circ I_1 , \quad I_1^2 = \text{Id} . \quad (17)$$

The map F is just a composition of these two involutions:

$$F = I_1 \circ I_0 . \quad (18)$$

For the map (1), we can take

$$I_0(x, y) = (-x, y - b \sin(2\pi x)) , \quad I_1(x, y) = (-x + a(1 - y^2), y) .$$

Now we investigate the implications of reversibility for the function K . If K satisfies (7), we also have $F^{-1} \circ K(\theta) = K(\theta - \omega)$ or, denoting $L(\theta) = K(-\theta)$ we have

$$F^{-1} \circ L(\theta) = L(\theta + \omega) . \quad (19)$$

Also, composing (7) with I_0 and using (16), we get

$$I_0 \circ K(\theta + \omega) = F^{-1} \circ I_0 \circ K(\theta) .$$

Thus, $I_0 \circ K$ and L satisfy the same equation, and the argument leading to (9) implies that

$$I_0 \circ K(\theta) = L(\theta + \sigma) = K(-\theta - \sigma) .$$

for some σ . Clearly, σ depends on the I_0 and on the origin of coordinates we have chosen for θ .

If we change the origin of coordinates in θ by $\sigma/2$, we obtain that

$$I_0 \circ K(\theta) = K(-\theta) . \tag{20}$$

Such K will be said to be *antisymmetric* with respect to I_0 . Note that if (20) holds, then $I_0 \circ K(0) = K(0)$, so that the antisymmetry is evident if we choose $K(0)$ on the symmetry line (fixed point set) of I_0 .

Composing (20) with I_1 , we obtain

$$\begin{aligned} K(\theta + \omega) &= F \circ K(\theta) = I_1 \circ I_0 \circ K(\theta) \\ &= I_1 \circ K(-\theta) . \end{aligned} \tag{21}$$

So, we see that if K is antisymmetric with respect to I_0 , a translation in θ by $\omega/2$ will make it antisymmetric with respect to I_1 . In this case, $K(0)$ belongs to the symmetry line for I_1 .

For the case of the standard non-twist map, we obtain using (21) for the x component and (20) for the y component that

$$\begin{aligned} K_x(-\theta) &= -K_x(\theta) \\ K_y(-\theta) &= K_y(\theta + \omega) . \end{aligned} \tag{22}$$

Of course, the other two identities that we obtain from (20) and (21) are equivalent to (22) and (7). One consequence of (22) is that the Fourier series of $K(\theta) - \theta$ is simply a sine series or equivalently, the phases of the complex Fourier transform (which is what we implemented numerically) are either $\pi/2$ or $3\pi/2$.

The equations (22) refer to parameterizations with the $K(0)$ on the symmetry line. In our calculations, we have chosen to take $K(0)$ to be the *indicator point* [28] which is known for sure to belong to the shearless curve. In this coordinates, we can see that the phases of

the Fourier coefficients should be $|k|\sigma + r_k$ where r_k is either $\pi/2$ or $3\pi/2$. This provides an independent check of the calculations of the Fourier coefficients and we discuss it in detail in Sec. V A.

We could use (14) to compute only the even (resp. odd) Fourier coefficients of $K_x(\theta) - \theta$ [resp. $K_y(\theta)$] and carry out the analysis only for them. We have chosen not to hardwire this symmetry into the calculations of Fourier spectra. In this way, the size of the odd (resp. even) coefficients gives us an estimate of the numerical error in our computations.

IV. COMPUTING THE REGULARITY OF THE PARAMETERIZATION FUNCTIONS

In this section, we will briefly describe the methods we used to compute the regularity of the parameterization functions. A much more extensive discussion can be found in [21], to which we refer for further details and for references to the mathematical literature.

In this paper, we will only use the two methods that were found to be the most accurate in [21], namely the so-called ‘‘Continuous Littlewood-Paley’’ and the ‘‘Difference methods’’. Notably, we do not consider the methods based on wavelets that were implemented in [21]. We refer to [21] for a comparison of various numerical methods.

A. Some mathematical results

Following the standard mathematical notation, we will express the regularity of the functions considered by saying that the function belongs to certain spaces in a family indexed by a number. We will use the following definition.

Definition 1. *For $\alpha = n + \beta$ with $n \in \mathbb{Z}$, $\beta \in (0, 1)$, we say that the function K is in $\Lambda_\alpha(\mathbb{T})$ when K is n times differentiable and*

$$|D^n K(\theta) - D^n K(\tilde{\theta})| \leq C|\theta - \tilde{\theta}|^\beta \tag{23}$$

for some constant C .

The above definition does not cover the case when α is an integer. It is a fact of life that many of the characterizations we will use become delicate when α is an integer. Fortunately, for the applications here, we do not need to consider the case when α is an integer.

The following results can be found in [29, Ch. 5, Lemma 5] and [30], respectively.

Theorem 1 (CLP). *The function K is in $\Lambda_\alpha(\mathbb{T})$ if and only if for some $\eta \geq 0$ there exists a $C > 0$ such that for any $t > 0$*

$$\left\| \left(\frac{\partial}{\partial t} \right)^\eta e^{-t\sqrt{-\Delta}} K \right\|_{L^\infty(\mathbb{T})} \leq C t^{\alpha-\eta} . \quad (24)$$

Remark 2. If the above result holds for some $\eta \geq 0$, then it holds for all $\eta \geq 0$.

Remark 3. The operator $e^{-t\sqrt{-\Delta}}$ is the Poisson kernel, which appears very often in the study of boundary value problems.

Theorem 2 (FD). *Let \mathcal{D}_θ^n be the finite difference operator,*

$$(\mathcal{D}_\theta^n K)(\theta) := \sum_{j=0}^n (-1)^j \binom{n}{j} K(\theta + (n-2j)\theta) .$$

Then $K \in \Lambda_\alpha(\mathbb{T})$ if and only if $\exists C > 0$ such that $\forall \theta \in \mathbb{T}$

$$\|\mathcal{D}_\theta^n K\|_{L^\infty(\mathbb{T})} \leq C |\theta|^\alpha . \quad (25)$$

Remark 4. The mathematical theory only requires (24) and (25) to be upper bounds. Nevertheless, we note that in our case, the bounds become identities for a significant range of parameters. The fact that the upper bounds are saturated is very possibly a consequence of the fact that, because of the renormalization picture, the functions we are considering are very self-similar. The same effect was also observed for the functions considered in [21]. (See also the discussion in Sec. III B.)

B. Some remarks on the numerical implementation

Both numerical methods considered here are made easier if the function K is specified at equally spaced points. The computation described in Section III, however, gives the function values for $\{\theta = n\omega\}_{n=0}^M$. Hence, the first step is to compute the function on an evenly spaced grid of size 2^{-N} . This is done using a cubic interpolation spline. Here N is chosen such that $2^N \approx M$.

To use Theorem 2 numerically, we note that substituting $\theta = 2^{-j}$ and $n = 1$ in Equation (25) gives

$$\log_2 \|\mathcal{D}_{2^{-j}}^1 K\|_{L^\infty(\mathbb{T})} \leq \text{const} - \alpha j . \quad (26)$$

Thus, the finite difference method simply evaluates the difference operators for various j , and we fit a straight line to a logarithmic plot of $\|\mathcal{D}_{2^{-j}}^1 K\|_{L^\infty(\mathbb{T})}$ versus j .

For the CLP method (Theorem 1), the most effective way to compute the Poisson kernel is to observe that it is diagonal in Fourier series. Specifically, if

$$K(\theta) = \sum_{n \in \mathbb{Z}} \hat{K}_n e^{-2\pi i n \theta} , \quad (27)$$

then

$$\left(\frac{\partial}{\partial t}\right)^\eta e^{-t\sqrt{-\Delta}} K(\theta) = \sum_{n \in \mathbb{Z}} (-2\pi|n|)^\eta \hat{K}_n e^{-2\pi t|n|} e^{-2\pi i n \theta} . \quad (28)$$

Using the Fast Fourier transform – which becomes possible once we have the function interpolated in an equally spaced grid – it is very efficient to compute the Fourier coefficients, multiply them by the appropriate coefficients, compute the inverse Fourier transform and then take the supremum over θ . By repeating the computation for several values of t and then fitting a straight line in a double logarithmic plot of $\left\| \left(\frac{\partial}{\partial t}\right)^\eta e^{-t\sqrt{-\Delta}} K \right\|_{L^\infty(\mathbb{T})}$ versus t , we obtain a good estimation of the regularity.

In the application of the CLP method, it is convenient to choose η in such a way that the power in t that results in the decay is as close to zero as possible. The reason is that the coefficients which are very close to zero are strongly affected by round-off error in the calculations.

Remark 5. In [21] it was argued that the CLP method was numerically more stable while the difference method was more susceptible to numerical error. Hence, in this paper, we will obtain the regularities from the CLP method and use the FD method as a confirmation.

V. RESULTS AND DISCUSSION

A. Visual observations of the Fourier spectrum

As mentioned earlier, the critical circles have self-similar structure which is manifested in the self-similarity of the parameterization functions. We can see this by looking at their Fourier spectra (the modulus of the Fourier coefficients). Fig. 3 shows the spectra of $K_x(\theta) - \theta$ for the three frequencies studied. We see that the Fourier spectra are periodic with a period of $6 \log_{10} \gamma$. Similar structure is also observed for the spectra of $K_y(\theta)$. We note that the peaks in the spectra are observed at values that are the denominators of the truncations




FIG. 3: Plot of $\log_{10}(n|\hat{K}_n|)$ vs. $\log_{10} n$ where \hat{K}_n are the Fourier coefficients of $K_x(\theta) - \theta$ for $\omega^{(1)}$ (left), $\omega^{(2)}$ (center), and $\omega^{(3)}$ (right). The abscissas marked here are the denominators of continued fraction convergents for that specific frequency.

of the continued fraction expansion of the corresponding winding number, with every sixth denominator showing a major peak while those in between show minor peaks. Specifically, the peaks occur at Fibonacci numbers only for the first two winding numbers.

Remark 6. In the case of twist maps, the period of such a plot is $\log_{10} \gamma$. The factor of 6 for the nontwist maps is related to the fact that the critical nontwist map belongs to a period-12 cycle of the renormalization group operator whereas the critical twist map is a fixed point of that operator.

As we noted in Sec. III B, the functions $K_x(\theta) - \theta$ and $K_y(\theta)$ can be made antisymmetric with appropriate choice of initial conditions, i.e., values of these functions at $\theta = 0$. These initial conditions (the points of intersection of the critical shearless circle with the symmetry lines) are known with a limited accuracy whereas the indicator points that we used as initial conditions are guaranteed to lie on the shearless circle. Indeed, in our calculations, the odd (resp. even) Fourier coefficients of $K_x(\theta) - \theta$ [resp. $K_y(\theta)$] are smaller when using the indicator points (10^{-11} to 10^{-10}) than when using the symmetry lines (10^{-10} to 10^{-9}).

Since we chose the indicator point as initial conditions, the phases of Fourier spectra lie on straight lines. The slope of these lines is $2\pi y_0$ where y_0 is the point of intersection of the shearless circle with the symmetry line of I_0 while the intercept is $\pi/2$ or $3\pi/2$.

To check the numerical accuracy of the Fourier coefficients, we also calculated the spectra after choosing the initial conditions to make the function $K_x(\theta) - \theta$ antisymmetric, i.e., choosing $K_x(0)$ to be a fixed point of I_0 . Figure 4 shows the phases for all $n < 2 \times 10^6$ (left panel) and also for n for which $\log_{10} |\hat{K}_n| > -9.5$ (right panel). We see that the phases are




FIG. 4: The phases of the Fourier transform of $K_x(\theta) - \theta$ with $K_x(0)$ chosen to be on the symmetry line of I_0 . Left panel shows all the phases while the right panel shows phases for n for which $\log_{10} \hat{K}_n > -9.5$.

indeed $\pi/2$ or $3\pi/2$ for all $n < 1.6 \times 10^6$ with $\log_{10} |\hat{K}_n| > -9.5$.

B. Regularity using CLP method

We compared results from different grid sizes to check for robustness of the methods used here. Figure 5 shows the plots of $\log_{10} \left\| \left(\frac{\partial}{\partial t} \right)^2 e^{-t\sqrt{-\Delta}} (\tilde{K}(\theta) - \theta) \right\|_{L^\infty(\mathbb{T})}$ versus $\log_{10} t$ computed with 2^{-N} number of points for $N = 24$ (boxes) and $N = 22$ (circles). In order to assess the accuracy, we also calculated the above norms for $N = 24$ by setting all the Fourier coefficients \hat{K}_n with magnitude less than 10^{-7} or 10^{-9} to zero, shown in Figure 5 with pluses and crosses respectively. We see that the plot for $N = 22$ compares very well with that for $N = 24$ with all $|\hat{K}_n| < 10^{-9}$ set to zero. We also see that for $\log_{10} t > -5$, all the calculations give essentially the same results.

The observations in Figure 5 make it apparent that there is a substantial range of values of $\log_{10} t$ for which (24) is close to a equality. (See also Figure 6.) This range is limited on one side (of small $\log_{10} t$) by the numerical error (round-off and truncation of Fourier series) and on the other side (of larger $\log_{10} t$) by the universal self-similar structure which is only true asymptotically.

Once we have identified the range where (24) is an equality, a good measure of the regularity is obtained by the slope of the linear fit within that range. We will see in Sec. V C




FIG. 5: Comparing the CLP method for different grid sizes 2^{-N} for $N = 24$ (boxes) and $N = 22$ (circles). We also show the results from the runs with Fourier coefficients of magnitude less than 10^{-9} (pluses) or 10^{-7} (crosses) set to zero.




FIG. 6: The plots of $\log_{10} \left\| \left(\frac{\partial}{\partial t} \right)^2 e^{-t\sqrt{-\Delta}} (\tilde{K}(\theta) - \theta) \right\|_{L^\infty(\mathbb{T})}$ versus $\log_{10} t$ for $K_x(\theta) - \theta$ (left) and $K_y(\theta)$ (right) used to calculate the regularity using the CLP method.

that the deviations from the linear fit show a universal behavior. Thus, a systematic way of choosing the range of fit is where the universality applies. Table I shows the regularities found using such ranges. The errors shown are the standard deviations of the linear regression.

	η	$\omega^{(1)}$	$\omega^{(2)}$	$\omega^{(3)}$
$K_x(\theta) - \theta$	1	0.700 ± 0.005	0.718 ± 0.004	0.698 ± 0.005
	2	0.679 ± 0.006	0.656 ± 0.005	0.671 ± 0.005
	3	0.647 ± 0.018	0.647 ± 0.018	0.636 ± 0.015
	4	0.687 ± 0.007	0.669 ± 0.007	0.676 ± 0.007
	5	0.683 ± 0.008	0.667 ± 0.009	0.672 ± 0.009
$K_y(\theta)$	1	0.724 ± 0.007	0.728 ± 0.003	0.730 ± 0.006
	2	0.700 ± 0.006	0.692 ± 0.006	0.692 ± 0.006
	3	0.645 ± 0.015	0.622 ± 0.015	0.629 ± 0.015
	4	0.690 ± 0.008	0.675 ± 0.008	0.680 ± 0.008
	5	0.690 ± 0.005	0.702 ± 0.006	0.698 ± 0.005

TABLE I: Regularities of the parameterization functions for three frequencies using the CLP method with different η . The error shown is the standard error from the linear regression. The ranges of $\log_{10} t$ were chosen so that corrections to the linear behavior are periodic, as explained in detail in Section V C.

Increasing the number of values of t at which the norm is calculated obviously decreases the standard deviation of the linear fit. Thus the errors shown in Table II can be made smaller or larger. Of course, choosing different ranges of $\log_{10} t$ for the linear fit give slightly different regularities, depending on whether the jumps in the deviation from the linear behavior (as seen in, e.g., Figure 7) are included or not. Thus, the errors shown in Table II include these systematic errors from all the five values of η used here.

Remark 7. The three frequencies considered here are all noble numbers. As a consequence, the invariant circles with rotation numbers $\omega^{(2)}$ and $\omega^{(3)}$ for the standard nontwist map are equivalent to circles with rotation number $\omega^{(1)}$ for maps which are obtained from the standard nontwist map by renormalization. The fact that the regularities we find here are the same for all the circles is equivalent to saying that they are the same for circles of rotation number $\omega^{(1)}$ for different maps. Hence, the fact that the regularities for different frequencies are seen to be the same is an evidence of universality of critical noble invariant circles.

C. Universal corrections to the CLP bound

If the inequalities (24) were identically saturated, we would have that $R(\log_{10} t) \equiv \log_{10} \left\| \left(\frac{\partial}{\partial t} \right)^2 e^{-t\sqrt{-\Delta}} K \right\|_{L^\infty(\mathbb{T})}$ would be an affine function of $\log_{10} t$ and the plot of $R(l)$ versus l would be a straight line.

Nevertheless, it is a prediction from renormalization group theory that there are systematic deviation from the linear behavior of R . In particular, there should be a periodic oscillation.

As it was argued in [21], the deviation of $R(l)$ from an affine behavior is related to the fact that the scaling predicted by the renormalization group is not a continuous scaling but just a discrete one. The renormalization group predicts that, for sufficiently small $t > 0$ (i.e., large negative $\log_{10} t$), there are numbers a, b so that $R(l + a) \approx R(l) + b$. This leaves open the possibility that, superimposed on the linear behavior of $R(l)$ versus l , there is a periodic behavior. That is, $R(l) \approx (b/a)l + P(l)$ where $P(l)$ is a function of period a . According to the renormalization group picture, the function P should be universal. Of course, for different functions in the same universality class, the asymptotic behavior may manifest itself only for very large negative values of l .

The periodic correction to $R(l)$ is closely related to the approximate periodicity found for the Fourier coefficients $|\hat{K}_n|$ as a function of $\log_{10} |n|$ which we already observed in Figure 3. The universality of periodic function $P(l)$ is related to the fact that there is an asymptotic shape for the period of the scaled Fourier coefficients.

We have verified the existence of a universal periodic function $P(l)$ by computing the finite differences in the plot of $R(l)$ versus l . The results for $\eta = 2, 4$ for $K_x(\theta) - \theta$ and for $\eta = 1, 3$ for $K_y(\theta)$ are depicted in Figure 7. We also see that, as commented earlier, the smaller values of η are more accurate, i.e., the periodicity of these corrections is more clearly seen for smaller η .

We note that the period of the oscillations in these plots is 6 times longer than the period observed in the problems in [21]. This is due to the fact that the renormalization operator in our case has period 6 rather than a fixed point as in the cases considered in [21]. This effect is clearly visible in the period of the Fourier coefficients as seen already in Section V A.

We also note that for different ω we obtain a different phase in the periodic correction. This is due to the fact that our ω have a different entry in the first place of the continued




FIG. 7: The finite differences in the plots of $\log_{10} \left\| \left(\frac{\partial}{\partial t} \right)^2 e^{-t\sqrt{-\Delta}} K \right\|_{L^\infty(\mathbb{T})}$ versus $\log_{10} t$ for $\eta = 2, 4$ for $K_x(\theta) - \theta$ (left) and for $\eta = 1, 3$ for $K_y(\theta)$ (right). The ranges selected for getting regularities are shown by arrows. Plots for all three frequencies are shown but those for $\omega^{(2)}$ (- - -) and $\omega^{(3)}$ (- - -) have been shifted horizontally to match that of $\omega^{(1)}$ (· · ·).

fraction expansion, so that they need one more application of the renormalization to start exhibiting the periodic behavior. Hence, in the figure, we have shifted the finite differences to exhibit that, for our examples, they agree as predicted by universality.

The existence of a periodic correction to the linear dependence of R as well as its agreement in the examples studied, gives us good confidence that we are observing real effects. From a more practical point of view, it gives us the following criterion: We fit a straight line to the plot of $R(l)$ versus l only in the region where the periodicity of the correction is manifest. These regions are shown in Figure 7. All these tasks were greatly facilitated by the remarkable public domain program `xmgrace`.

D. Regularity using the finite difference method

As with the CLP method, we first compare results for different grid sizes. The plot of the $\log_2 \|\mathcal{D}_{2^{-j}}^1 K\|_{L^\infty(\mathbb{T})}$ vs. j is shown in Fig. 8 for $\tilde{K}_x(\theta) - \theta$ for the frequency $\omega^{(1)} = 1/\gamma$ for $N = 20, 22,$ and 24 . We note that there is a break at $j = 10$. The regularity obtained from the plot with $j > 10$ agrees very well with the CLP method while that obtained from all the points with $j > 5$ is lower. Since this method is affected more by round-off errors, we have used it only as a quick confirmation of results from the CLP method. The agreement between the two methods is much better for $K_y(\theta)$ probably because the round-off errors in




FIG. 8: Comparing the plots of $\log_2 \|\mathcal{D}_{2^{-j}}^1 K\|_{L^\infty(\mathbb{T})}$ versus j with 2^N points for $N = 20, 22, 24$ for $\tilde{K}_x(\theta) - \theta$ for the frequency $\omega^{(1)}$.




FIG. 9: The plots of $\log_2 \|\mathcal{D}_{2^{-j}}^1 K\|_{L^\infty(\mathbb{T})}$ vs. j for $K_x(\theta) - \theta$ (left) and $K_y(\theta)$ (right) used to calculate the regularity using the finite difference method.

iterating the map F propagate faster along the angle coordinate (x) [and hence in $K_x(\theta)$] than in the action coordinate (y) [and hence in $K_y(\theta)$].[26, Appendix C].

Figure 9 compares the plots of $\log_2 \|\mathcal{D}_{2^{-j}}^1 K\|_{L^\infty(\mathbb{T})}$ versus j for $N = 24$ for all the three frequencies for $K_x(\theta) - \theta$ and $K_y(\theta)$ with the line whose slope is the regularity obtained from the CLP method. We see that the bound in (25) is not saturated for $j < 10$ for $K_x(\theta) - \theta$.

From the regularities of the parameterization functions $K_x(\theta) - \theta$ and $K_y(\theta)$ for the three frequencies $\omega^{(1)}$, $\omega^{(2)}$, and $\omega^{(3)}$ shown in Table II, we conclude the universal regularity of $K(\theta)$ for critical shearless noble circles to be 0.69 ± 0.03 .

	$\omega^{(1)}$	$\omega^{(2)}$	$\omega^{(3)}$
$K_x(\theta) - \theta$	0.68 ± 0.02	0.67 ± 0.02	0.67 ± 0.01
$K_y(\theta)$	0.69 ± 0.02	0.68 ± 0.03	0.69 ± 0.03

TABLE II: Regularities of $K_x(\theta) - \theta$ and $K_y(\theta)$ for the three frequencies.

Acknowledgments

R.L. has been partially supported by NSF. N.P. has been partially supported by NSF grant DMS-0405903. A.A. has been partially supported by U.S. Department of Energy Contract No. DE-FG01-96ER-54346. We thank the Mathematics Department at UT-Austin for the use of computer facilities.

-
- [1] T. H. Stix, Phys. Rev. Lett. **36**, 10 (1976).
 - [2] G. A. Oda and I. L. Caldas, Chaos, Solit. and Fract. **5**, 15 (1995).
 - [3] R. Balescu, Phys. Rev. E **58**, 3781 (1998).
 - [4] W. Horton, H. B. Park, J. M. Kwon, D. Strozzi, P. J. Morrison, and D. I. Choi, Phys. Plasmas **5**, 3910 (1998).
 - [5] K. Ullmann and I. L. Caldas, Chaos, Solit. and Fract. **11**, 2129 (2000).
 - [6] D. Del-Castillo-Negrete and M.-C. Firpo, Chaos **12**, 496 (2002).
 - [7] A. Munteanu, E. Petrisor, E. García-Berro, and J. José, Commun. Nonlinear Sci. Numer. Simul. **8**, 355 (2003).
 - [8] E. Petrisor, J. H. Misguich, and D. Constantinescu, Chaos Solitons Fractals **18**, 1085 (2003).
 - [9] D. Del-Castillo-Negrete and P. J. Morrison, Phys. Fluids A **5**, 948 (1993).
 - [10] J. E. Howard and J. Humpherys, Phys. D **80**, 256 (1995).
 - [11] C. Simó, Regul. Chaotic Dyn. **3**, 180 (1998).
 - [12] G. Voyatzis, E. Meletlidou, and S. Ichtiaroglou, Chaos Solitons Fractals **14**, 1179 (2002).
 - [13] H. R. Dullin and J. D. Meiss, Chaos **13**, 1 (2003).
 - [14] J. Franks and P. Le Calvez, Ergodic Theory Dynam. Systems **23**, 111 (2003).
 - [15] D. Del-Castillo-Negrete, J. M. Greene, and P. J. Morrison, Physica D **91**, 1 (1996).

- [16] D. del Castillo-Negrete, J. M. Greene, and P. J. Morrison, *Phys. D* **100**, 311 (1997).
- [17] A. Delshams and R. de la Llave, *SIAM J. Math. Anal.* **31**, 1235 (2000).
- [18] A. Apte, A. Wurm, and P. J. Morrison, *Chaos* **13**, 421 (2003).
- [19] H. Koch and D. Gaidashev, *Nonlinearity* **17**, 1713 (2004).
- [20] A. Apte, Ph.D. thesis, Univ. of Texas at Austin (2004).
- [21] R. de la Llave and N. P. Petrov, *Experimental Math.* **11**, 219 (2001).
- [22] N. Petrov, Ph.D. thesis, Univ. of Texas at Austin (2002).
- [23] A. Apte, A. Wurm, and P. J. Morrison, *Physica D*, to appear (2003).
- [24] D. Gaidashev, *mp_arc* # 03-542 (2003).
- [25] A. Y. Khinchin, *Continued Fractions* (University of Chicago Press, Chicago, 1964), 3rd ed.
- [26] J. M. Greene, *Jour. Math. Phys.* **20**, 1183 (1979).
- [27] R. de la Llave, A. Gonzalez, A. Jorba, and J. Villanueva, *mp_arc* # 03-421 (2003).
- [28] S. Shinohara and Y. Aizawa, *Progr. of Theor. Phys. Suppl.* **139**, 527 (2000).
- [29] E. M. Stein, *Singular Integrals and Differentiability Properties of Functions* (Princeton University Press, Princeton, 1970).
- [30] S. G. Krantz, *Exposition. Math.* **1**, 193 (1983).